

**Moral Hazard and Less Invasive Medical Treatment for Coronary Artery Disease:  
An Analysis of Smoking in the National Health Interview Survey**

Jason Hockenberry

Jesse Margolis

Shin-Yi Chou

Michael Grossman\*

March 31, 2014

Abstract: Over the last several decades, numerous medical studies have compared the effectiveness of two common procedures for Coronary Artery Disease: Percutaneous Coronary Intervention (PCI) and Coronary Artery Bypass Graft (CABG). Most evidence indicates that CABG – the more invasive procedure – leads to superior long term outcomes for otherwise similar patients, though there is little consensus as to why. In this article, we propose a novel explanation: patient offsetting behavior. We hypothesize that patients who undergo the more invasive procedure, CABG, are more likely to improve their behavior – eating, exercise, smoking, and drinking – in a way that increases longevity. To test our hypothesis, we use Medicare records linked to the National Health Interview Survey to study one such behavior: smoking. We find that CABG patients are 12 percentage points more likely to quit smoking in the one-year period immediately surrounding their procedure than PCI patients, a result that is robust to numerous alternative specifications.

DRAFT – PLEASE DO NOT CITE

\* Hockenberry: Emory University and NBER (jason.hockenberry@emory.edu). Margolis: City University of New York Graduate Center (jmargolis@gc.cuny.edu). Chou: Lehigh University and NBER (syc2@lehigh.edu). Grossman: City University of New York Graduate Center, NBER, and IZA (mgrossman@gc.cuny.edu).

Acknowledgement: Research for this paper was supported by grant number 5R21 AG033876 from the National Institute on Aging and the Office of Behavioral and Social Sciences Research to the National Bureau of Economic Research. This paper was presented at seminars at the University of Chicago, the University of Illinois at Chicago, Johns Hopkins University, the University of Melbourne, the Ohio State University, and the University of Connecticut. It also was presented at the 2013 Bureau of the Census Research Data Centers Conference at the Federal Reserve Bank of Atlanta. We would like to thank the participants in those forums for helpful comments and suggestions. We also would like to thank Peter Cram for his medical insight and Sandra Decker for her advice on working with the linked NHIS/Medicare data. We are indebted to Jonathan Fisher, Research Data Center (RDC) Administrator at the New York Bureau of the Census RDC at Baruch College, and Frances McCarty, Senior Service Fellow at the National Center for Health Statistics RDC, for their assistance in helping us to gain access to and work with restricted files from the National Health Interview Survey. All errors are our own.

## **I. Introduction**

Coronary Artery Disease (CAD) is a common and deadly disease. In 2010, over 350,000 people died of CAD in the United States, making the disease responsible for roughly one in seven deaths (Murphy, Xu, and Kochanek, 2012). CAD is caused by a buildup of plaque on the arterial walls leading to the heart, resulting in reduced blood flow. If the buildup is not checked, CAD can result in an acute myocardial infarction (AMI, a.k.a. “heart attack”) due to insufficient oxygen reaching the heart.

A number of medical treatments are available to patients with CAD. First, and least invasive, is “medical management.” Medical management involves non-surgical treatment including prescription medication, lifestyle modification, and frequent monitoring. The second treatment is a revascularization procedure known as Percutaneous Coronary Intervention (PCI). A doctor (usually a cardiologist) performing PCI makes a small incision and arthroscopically inserts and inflates a balloon at the site of the lesion to expand the vessel. PCI in the modern era usually involves the placement of a wire mesh stent at the blockage site, which assists in keeping the arterial walls expanded to maintain blood flow. The PCI procedure takes approximately 60 minutes and the patient usually spends one night in the hospital.<sup>1</sup> The third and generally most invasive treatment is Coronary Artery Bypass Graft (CABG)<sup>2</sup>, a major surgical procedure that involves harvesting a section of vessel from a different area of the body (either vessels in the groin or chest wall), opening the chest cavity via a sternectomy, and connecting one healthy part of the diseased artery to another, surgically bypassing the lesion. CABG surgery takes approximately four hours and patients generally spend at least a week recovering in the hospital.<sup>3</sup>

---


<sup>1</sup> [http://www.medicinenet.com/coronary\\_artery\\_bypass\\_graft/article.htm](http://www.medicinenet.com/coronary_artery_bypass_graft/article.htm) (accessed 5/31/12)

<sup>2</sup> Less invasive CABG procedures have been in development and increasing use in recent years, though these were very infrequent during the period we examine.

<sup>3</sup> [http://www.medicinenet.com/coronary\\_angioplasty/article.htm](http://www.medicinenet.com/coronary_angioplasty/article.htm) (accessed 5/31/12)

Of the two procedures, PCI is the more recent, having been initially used in the late 1970s, over than a decade after CABG was first performed. Its use expanded rapidly upon FDA approval of the coronary stent in 1994 (Cutler and Huckman, 2003). Since the development of PCI, there have been numerous studies comparing the effectiveness of the two procedures in various populations (see Hlatky et al., 2009, Weintraub et al., 2012, and Mohr et al., 2013 for three recent studies that summarize prior research). While the results vary, the emerging consensus is that CABG patients have worse short-run outcomes than similar PCI patients – partly due to higher perioperative mortality – but better long term outcomes. Results from a large observational study (Weintraub et al., 2012) are reproduced in Figure 1, showing that the survival curve for CABG patients is initially lower than for PCI patients, but that this trend is reversed a year or more after the procedure. A meta-analyses of ten randomized controlled trials shows a similar pattern (Hlatky et al., 2009).

Figure 1 – Comparative Effectiveness: PCI vs. CABG (Weintraub et. al, 2012)



Source: Weintraub et al., 2012. N = 189,793. Figure adjusts based on inverse propensity score weighting.

While studies show that otherwise similar CABG patients have superior long run outcomes than PCI patients, there is little consensus as to why. In this paper, we propose a novel hypothesis: patient offsetting behavior. Specifically, a patient who undergoes CABG rather than PCI is more likely to change his behavior in a way that promotes good health and a longer life: he is more likely to quit smoking, begin exercising, improve his diet, and avoid excessive alcohol intake. This might happen because the more invasive nature of CABG – a patient’s heart and lungs are bypassed during the surgery, he is in the hospital for a week, has a longer post-operative recovery period, and is left with a major scar and residual pain from the sternectomy – sends a stronger signal to the patient that he has a serious health problem.

This hypothesis is consistent with a prior economic research on moral hazard, showing that individuals change their behavior when their perceived risks change. Peltzman’s (1975) study of the effects of automobile safety regulation is a classic and seminal example. He develops a model in which the legal mandate to install various safety devices on automobiles lowers the price of fast and reckless driving because it lowers the probability that the driver will die in an accident. Hence the demand for this activity rises. Empirically, he finds that the increase in this offsetting behavior (reckless driving) is so large that the regulations at issue had very little impact on highway deaths and actually increased pedestrian deaths. More recently, Dave and Kaestner (2009) investigate the impact of health insurance access on the health behaviors of the elderly, showing that access to Medicare at age 65 leads to a reduction in preventative behaviors and an increase in risky health behavior amongst the elderly. Peltzman (2011) demonstrates how medical technology breakthroughs can lead to offsetting behavior by showing that the age cohorts that benefited the most from the introduction of antibiotics experienced worse mortality rates from risky health behaviors.

In this study, we test one potential behavioral response to surgery – smoking – and see results consistent with patient offsetting behavior. Patients who undergo CABG – the more invasive procedure – are 12 percentage points more likely to quit smoking than PCI patients. Our results are robust to a number of different specifications, including a simple grouped-by-year regression using 11 observations, done in the spirit of Donald & Lang (2007).

## **II. Data**

We use individual Medicare data merged with responses from the National Health Interview Survey (NHIS). The Medicare records identify those patients who have been diagnosed with CAD and show which of them have undergone PCI or CABG, along with the exact date of each diagnoses and procedure. The Medicare data also allow us to control for disease severity and other conditions that might be correlated with procedure type and induce quitting, such as a myocardial infarction (a.k.a. “heart attack”). The NHIS provides information on smoking and quitting behavior, as well as individual characteristics.

The Medicare data are provided by the Center for Medicare & Medicaid Services (CMS). To identify CAD patients and the type of treatment they underwent, we use the Medicare Standard Analytical Files, including the Inpatient, Outpatient, Skilled Nursing Facility, Carrier, Durable Medical Equipment, Home Health Agency, and Hospice claims files. These files contain one or more records for each individual.<sup>4</sup> Each record contains the ICD-9-CM codes for all diagnoses made and procedures performed during that stay or claim. We identify CAD patients as those who have at least one diagnosis code beginning with 410, 411, 412, 413, or 414.

---

<sup>4</sup> A single record in the Inpatient file corresponds to a stay in a hospital. A single record in the Skilled Nursing Facility file corresponds to a stay in a Skilled Nursing Facility. A single record in the Outpatient file corresponds to a claim by an institutional outpatient provider (Hospital outpatient clinic, rural health clinics, etc.). A single record in the Carrier claim file corresponds to a claim by a non-institutional outpatient provider (physicians, physician assistants, etc.)

We identify PCI patients as those CAD patients with at least one procedure code beginning with 0066, 3601, 3602, 3605, or 3606. We identify CABG patients as those CAD patients with procedure codes beginning with 361.<sup>5</sup> Finally, we identify medically managed patients as those patients who have been diagnosed with CAD, but do not have a concurrent or subsequent PCI or CABG procedure.<sup>6</sup>

The NHIS is an annual survey of approximately 85,000 individuals in over 30,000 U.S. households run by the National Center for Health Statistics (NCHS), part of the Centers for Disease Control and Prevention (CDC). All participants are asked questions about their general state of health and disability. Each year, a subset of approximately 30,000 individuals is asked about their smoking habits. These respondents are asked if they have ever smoked 100 cigarettes in their life. For those who say yes, they are asked if they currently smoke every day, some days, or not at all. If they do not currently smoke, they are asked when they quit, a question they can answer in days, weeks, months, or years. We use the responses to these questions to create a synthetic panel, identifying whether a person smoked on each date prior to their NHIS interview. Each person is categorized as either an always smoker, a never smoker, or a quitter who smoked up to the day she reports quitting.<sup>7</sup>

The individual NHIS responses have been linked to Medicare data by the CDC and CMS and made available as a restricted-use dataset to researchers. The linkage is based on social security number, date of birth, and gender. To be linked, the data must match on all three fields. To date, the CDC and CMS have linked the 1994-1998 NHIS surveys to Medicare data from

---

<sup>5</sup> For both PCI and CABG, we exclude the small number of patients who do not have a concurrent or prior CAD diagnosis.

<sup>6</sup> A patient who is diagnosed with CAD before her NHIS interview date and has PCI or CABG after her NHIS interview date is counted as medically managed at the time of the NHIS survey.

<sup>7</sup> This categorization vastly over-simplifies the complexity of smoking and quitting behavior, but still allows us to investigate our key question: what is the *difference* in quitting behavior between CAD patients undergoing medical management, PCI, or CABG.

1991-2007 and the 1999-2005 NHIS surveys to Medicare data from 1999-2007. The linkage is described further in the appendix.

### **III. Initial Analysis**

In total, 12,265 NHIS respondents were linked to Medicare data and diagnosed with CAD.<sup>8</sup> Of these individuals, between the date of their diagnosis and the date of their NHIS interview, 10,713 patients were treated only with medical management, 771 patients underwent PCI but not CABG surgery, and 781 patients underwent CABG surgery.<sup>9</sup> Though our focus is on the two procedures – PCI and CABG – we include medically managed patients in all analyses for two reasons. First, due to the substantially greater number of medically managed patients, including them improves the precision of our covariate estimates (e.g. determining the impact of having a heart attack on smoking). Second, since medical management is the least invasive treatment for CAD, we might expect medically managed patients to quit at the lowest rate. This result, which we find in the data, supports our theory that changes in smoking behavior are related to treatment invasiveness. Our main finding, however – that CABG patients quit smoking at a higher rate than PCI patients – is robust to excluding medically managed patients from the analysis entirely.

Basic characteristics of the 12,265 CAD patients are shown in Table 1. Overall, when compared to patients undergoing medical management, patients who undergo a procedure (PCI or CABG) are more likely to be younger, male, and white. PCI and CABG patients appear to

---

<sup>8</sup> To be included, patients had to be diagnosed with CAD after the start of our Medicare data, but before the date of their NHIS interview (so that we have information on their smoking behavior both before and after their treatment).

<sup>9</sup> These counts are weighted by the NHIS probability weights. The unweighted totals are 10,772 medically managed patients, 723 PCI patients, and 770 CABG patients. Unweighted, 99 patients underwent both PCI and CABG surgery. These patients are included in the CABG category, because that is the more invasive treatment. Our results are robust to including them in the PCI category or excluding them altogether.



have largely similar demographic characteristics, though CABG patients are somewhat more likely to be male. When comparing medical conditions, both PCI and CABG patients are substantially more likely than medically managed patients to have had their first Acute Myocardial Infarction (AMI, a.k.a. “heart attack”) within six months of initiating treatment. A number of other comorbidities – including congestive heart failure and valvular disease – show up most frequently in CABG patients, followed by PCI patients. In some of our regression specifications, we control for the covariates shown in Table 1.

Table 1 – Characteristics by Treatment<sup>10</sup>

Demographic Characteristics				Medical Conditions			
	MM	PCI	CABG		MM	PCI	CABG
<i>Age</i>				<i>First AMI Within 6 Months of Treatment</i>			
< 55	4%	4%	2%	Yes	8%	40%	37%
55-64	7%	8%	7%	No	92%	60%	63%
65-69	24%	25%	26%	<i>% With Comorbidity Within 6 Months of Treatment</i>			
70-74	22%	25%	26%	Congestive heart failure	14%	21%	34%
75-79	21%	20%	23%	Valvular disease	11%	19%	27%
80-84	13%	13%	13%	Pulmonary circulation disorder	2%	4%	4%
85+	8%	5%	2%	Peripheral vascular disorder	10%	18%	22%
<i>Gender</i>				Paralysis	1%	1%	3%
Male	44%	53%	59%	Other neurological	3%	3%	3%
Female	56%	47%	41%	Chronic pulmonary disease	15%	12%	22%
<i>Race</i>				Diabetes w/o chronic comp.	14%	16%	17%
Asian	1%	1%	1%	Diabetes w/ chronic comp.	4%	7%	10%
Black	9%	6%	4%	Hypothyroidism	7%	6%	5%
Hispanic	5%	3%	4%	Renal failure	2%	4%	4%
White	85%	89%	90%	Liver disease	1%	1%	0%
Mult./Oth/Unknown	1%	1%	1%	Chronic Peptic ulcer disease	0%	1%	0%
<i>Education</i>				HIV and AIDS	0%	0%	0%
Elem (K-8)	21%	17%	20%	Lymphoma	0%	0%	0%
HS (non-grad); GED	19%	23%	18%	Metastatic cancer	1%	1%	1%
HS grad	29%	30%	29%	Solid tumor without metastasis	5%	5%	3%
Some col; AA deg.	18%	20%	18%	Rheumatoid arthritis	3%	2%	2%
BA degree	7%	7%	8%	Coagulation deficiency	3%	4%	9%
Grad. Degree	5%	5%	6%	Obesity	3%	8%	9%
Unknown	1%	0%	1%	Weight loss	1%	1%	2%
<i>Family Income</i>				Fluid and electrolyte disorders	10%	13%	24%
\$0 to \$9,999	19%	15%	14%	Blood loss anemia	1%	3%	3%
\$10,000 to \$19,999	25%	22%	24%	Deficiency anemias	11%	14%	23%
\$20,000 to \$35,000	20%	24%	24%	Alcohol abuse	1%	1%	1%
\$35,000 or over	18%	22%	21%	Drug abuse	0%	0%	0%
Unknown	18%	17%	17%	Psychoses	2%	2%	2%
				Depression	4%	7%	6%
				Hypertension	37%	37%	36%
<i>Count (N = 12,265)</i>	10,713	771	781	<i>Count (N = 12,265)</i>	10,713	771	781

Note: All data are weighted by the NHIS probability weights. Age is as of diagnosis (CAD) or procedure (PCI / CABG). Comorbidities based on Elixhauser et al. (1998)

Table 2 shows the smoking status of each group of respondents – medical management, PCI, and CABG – as of the date of the NHIS interview. In looking at this table, two items merit

<sup>10</sup> Results presented in this paper include all Medicare participants, regardless of age. Results excluding those under 65, available upon request, are similar.

notice. First, CABG and PCI patients are more likely to have ever smoked than medically managed patients (i.e. the percentage of respondents who never smoked is lower for CABG and PCI patients). Second, most people who have ever smoked have quit smoking by the time of the NHIS interview, a trend that is most pronounced for CABG patients. While 61.0% of CABG patients in our study smoked at some point in their life, only 9.1% smoke as of the NHIS interview date. PCI patients have a lower proportion of quitters, followed by medically managed patients.

Table 2 – Smoking Status as of NHIS Interview Date

Smoking Status as of Survey	Treatment			Total
	MM	PCI	CABG	
Current	12.2%	11.3%	9.1%	11.9%
Quit	42.6%	49.3%	52.0%	43.6%
Never Smoked	45.3%	39.5%	39.0%	44.5%
Total	100.0%	100.0%	100.0%	100.0%
<i>Count</i>	<i>10,713</i>	<i>771</i>	<i>781</i>	<i>12,265</i>

Note: This table shows the smoking status of every NHIS respondent who was diagnosed with CAD prior to their interview date. Data are weighted by the NHIS probability weights.

The data in Table 2 are consistent with the broad hypothesis in our study – patients who undergo a more invasive treatment for CAD are more likely quit smoking. However, they could also be consistent with a story in which people who undergo CABG are also more likely to quit smoking for reasons unrelated to their surgery. If our hypothesis is true, we should see that the differential quitting behavior between CABG, PCI, and medically managed patients is driven by quits that occur close to the date of the treatment.

Table 3 – Quitting Within Six Months of Treatment

	Treatment			Total
	MM	PCI	CABG	
Smoke before treatment	15.7%	15.6%	15.1%	15.7%
Smoke after treatment	14.2%	12.9%	10.5%	13.8%
Percentage Point Change	-1.6	-2.7	-4.5	-1.9
<i>Percent Change (Quit Rate)</i>	<i>10.1%</i>	<i>17.4%</i>	<i>30.1%</i>	<i>11.8%</i>
Count	10,713	771	781	12,265

Note: This table includes every NHIS respondent who was diagnosed with CAD in our data prior to their interview date. It shows their smoking status exactly six months before and exactly six months after diagnosis (CAD) or surgery (PCI/CABG). Data are weighted by the NHIS probability weights.

Table 3 focuses on those quits that take place immediately around the initiation of treatment, where the initiation of treatment is defined to be the diagnosis date for medically managed patients and the procedure date for PCI and CABG patients. The “before” period is exactly six months before the diagnosis/procedure date, while the “after” period is exactly six months after diagnosis/procedure date.<sup>11</sup> Among the 10,713 patients diagnosed with CAD who receive only medical management, approximately 15.7% smoked six months before their diagnosis and 14.2% smoked six months after their diagnosis. The quitters represent a 1.6 *percentage point* reduction in the number of smokers and a 10.1 *percent* reduction. The corresponding numbers for PCI are a 2.7 percentage point reduction and a 17.4 percent reduction. For CABG, they are a 4.5 percentage point reduction and a 30.1 percent reduction.

Further evidence is provided by Figures 2 and 3. In Figure 2, we calculate the percentage of the population smoking at twelve points in time, measured in years relative to the date of


<sup>11</sup> Creating a “quit window” around the date of diagnosis/procedure is necessary for two reasons. First, it is unlikely that many individuals quit on exactly the day their treatment began. Second, our smoking data, which are based on individuals’ recollections, are insufficiently precise to pinpoint the exact day of quitting. Our conclusions do not change with other reasonable definitions of the quit window.

diagnosis (in the case of medically managed patients) or procedure (in the case of PCI and CABG patients)<sup>12</sup>. In the CABG series, for example, the year -3.5 shows the percentage of CABG patients who were smoking exactly three and a half years prior to their procedure date. In the 10 years prior to diagnosis/procedure, the three series track each other reasonably closely. At the first point on the graph – 9.5 years before diagnosis/procedure – CABG patients are roughly one percentage point more likely than PCI patients to smoke, who are, in turn, roughly one percentage point more likely than MM patients to smoke. In the periods immediately prior to diagnosis/procedure, the three groups smoke at roughly equal rates. The differences between the three series emerge most starkly in the period immediately after diagnosis/procedure. Six months after diagnosis/procedure, CABG patients are substantially less likely to smoke than either PCI or CABG patients.

---

<sup>12</sup> Because we have data on only the most recent quit date for each individual, we assume that each smoker was smoking in all years before their quit date. Since we are using Medicare data for our analysis, most people are over 65 when they received their diagnosis or procedure, and it is unlikely that they started smoking for the first time in the ten years immediately prior. It is possible that individuals quit and restarted during this time period, and we do not distinguish them from continuous smokers.

Figure 2 – Smoking Rate by Year Relative to Diagnosis (MM) or Procedure (PCI & CABG)




Note: Figure shows the percentage of respondents who smoked at each point relative to their diagnosis/procedure. Weighted by NHIS probability weights

Figure 3 displays the same data in a different format, showing the annual quit rate for patients in each of the three treatment groups relative to the date of diagnosis/procedure. For the group that receives only medical management, roughly 5% of smokers quit each year in the nine years prior to being diagnosed, a rate that doubled to 10% during the year of their diagnosis with coronary artery disease. The PCI and CABG series show a similar trend, though they represent fewer individuals and are somewhat noisier. In the years prior to their procedure, roughly 5% of smokers quit each year, though this percentage began to rise the year before the procedure date. During the procedure year – defined to be the six month window on either side of the procedure date – the quit rate jumped to 17% for patients receiving PCI and 30% for patients receiving

CABG. In the year following diagnosis/procedure, the quit rate for all three groups dropped back to approximately 5%. Figures 2 and 3 provide reasonably compelling evidence that at least a portion of the increased quit rate for more invasive treatments observed in Table 2 is related to treatment received, and not simply a spurious correlation.

Figure 3 – Quit Rate by Year Relative to Diagnosis (MM) or Procedure (PCI & CABG)



Note: figure shows the % of remaining smokers who quit at each point relative to their diagnosis/procedure. Weighted by the NHIS probability weights

#### IV. Results

To further explore the relationship between treatment for coronary artery disease and smoking behavior, we fit two related models of quitting smoking. The first is a discrete time linear probability hazard function with 11 periods: 9 before treatment and 2 after. We fit the

model using individual data, allowing us to control for time-varying events – like CAD diagnosis or a patient’s first AMI – that may occur either prior to or concurrent with a PCI or CABG procedure. The second model is a multi-period quit function using grouped data, inspired by Donald and Lang (2007). In this model, we build a synthetic panel, grouping the individual data by period and treatment type into 33 cells (3 treatment types and 11 periods), and run difference-in-differences regressions with 11 data points.

Before presenting the results of these models, it is useful to point out the relationship between a smoking participation function and a quit function. As an identity<sup>13</sup>

$$\frac{S_t}{S_{t-1}} \equiv 1 - q_t, \quad (1)$$

where  $s_t$  and  $s_{t-1}$  are the smoking participation rates in periods  $t$  and  $t-1$ , respectively, and  $q_t$  is the quit rate in the window defined by periods  $t$  and  $t-1$ . All rates are defined as fractions and can be interpreted as probabilities at the individual level. Take natural logarithms of the identity to obtain

$$\ln \frac{S_t}{S_{t-1}} \equiv \ln(1 - q_t) \cong -q_t. \quad (2)$$

---

<sup>13</sup> Let  $S_t$  be the number of smokers in period  $t$ , and let  $S_{t-1}$  be the number of smokers in period  $t-1$ . Let  $Q_t$  be the number of quitters in period  $t$  (the number who smoke in period  $t-1$  but do not smoke in period  $t$ ). Assume as is the case in our data that there are no starters or re-starters. Then

$$S_t \equiv S_{t-1} - Q_t.$$

Divide both sides of the identity by  $N$ , the size of the population:

$$\frac{S_t}{N} \equiv \frac{S_{t-1}}{N} - \frac{Q_t}{S_{t-1}} \frac{S_{t-1}}{N}.$$

Solve the last identity for  $\frac{S_t}{\frac{S_{t-1}}{N}}$  to obtain equation (1).



The approximation in the last part of equation (2) is better for smaller values of  $q$  ( $q \leq 0.2$  is generally used as a rule of thumb). However, even for a quit rate as large as 0.3 (the largest rate in our data),  $\ln(1 - q) = -0.350$ , which is close to 0.3. Equation (2) indicates that a regression in which the first difference of the log of smoking participation is the dependent variable should have approximately the same coefficients with the signs reversed as one in which the quit rate is the dependent variable. It also suggests that it is useful to begin with a log smoking participation function to arrive at a specification of a quit function. In particular, if the log smoking participation function contains individual fixed effects, these effects are eliminated by taking first differences to obtain the quit function.

*Model 1: Discrete-time Linear Probability Hazard Model (Individual Data)*

To implement the first model, we develop a synthetic panel with 12 points in time, as shown in Figure 2. For CABG and PCI patients, there are 10 points in time prior to treatment (from 9.5 years before to 0.5 years before) and two points in time after treatment (0.5 years after and 1.5 years after). For medically managed (MM) patients, there are the same 10 points in time before diagnosis and the same two points in time after diagnosis. To focus on the key aspects of the model, we ignore the socioeconomic and demographic variables for the time being, assume a single Elixhauser comorbidity, and suppress the subscript  $i$  for an individual. Let  $f_g$  be a person-specific fixed effect. Let  $a_t$  be a dummy variable that equals 0 at each point in time before treatment for PCI or CABG and equals 1 at each point in time after treatment. Specifically,  $a_t$  equals 1 in periods 11 and 12. This variable is not relevant for MM patients (see below). Let  $d_t$  be a dummy variable that equals 0 at all points in time before diagnosis and equals 1 at all points in time after diagnosis. Let  $h_t$  be a dummy variable that equals zero before an AMI and equals 1

after an AMI. Finally, let  $e_t$  equal 0 before an Elixhauser comorbidity is reported and equals 1 thereafter.

The log smoking participation model for PCI and CABG patients ( $g = p$  or  $c$ )

$$\ln s_{gt} = f_g - \beta_g a_{gt} - \phi h_{gt} - \gamma e_{gt} - \alpha d_{gt} - \lambda t, \quad (3)$$

where we assume a linear trend in the absence of treatment. The model for MM patients is the same except that  $a_{mt}$  coincides with  $d_{mt}$ , so that we constrain  $\beta_m$  to equal zero. After pooling and taking first differences, one obtains

$$-(\ln s_t - \ln s_{t-1}) \cong q_t = \lambda + \alpha(d_t - d_{t-1}) + \beta_p p(a_t - a_{t-1}) + \beta_c c(a_t - a_{t-1}) + \phi(h_t - h_{t-1}) + \gamma(e_t - e_{t-1}). \quad (4)$$

Strictly speaking, time-invariant individual characteristics, such as formal schooling, can only be added to equation (4) by assuming that they interact with the linear trend in equation (3). Our results are not affected by allowing the trend to be nonlinear or by allowing individual characteristics to interact with the indicator for the period after treatment in addition to their interactions with a linear trend.

We fit equation (4) as a discrete time linear probability hazard model. We include only individuals who smoke at the first point in time for which we compute smoking participation (9.5 years before treatment or diagnosis), dropping everyone who never smoked or had previously quit. Each person is assigned a  $q_{it}$  variable that is equal to one in the period in which they quit and zero in all other periods. Individuals are deleted once they quit. The model in equation (4) has at most 11 observations per person corresponding to the 11 time periods in Figure 3.

Individuals who smoke in all periods are the censored observations. The first period is defined by the window starting 9.5 years before treatment and ending 8.5 years before treatment. The last period is the window from 0.5 years after treatment to 1.5 years after treatment. The key window is period 10 and spans the dates from half a year before treatment or diagnosis to half a year after. That is the only period in which  $a_t - a_{t-1}$  is equal to 1. Since there are repeat observations on all persons except those who quit in period 1, we cluster standard errors at the individual level. Standard errors that ignore clustering are, however, very similar to those that take account of it. This would be the case if the unspecified disturbance term in the log smoking participation function in equation (3) is a random walk. In that case, we eliminate serial correlation by taking first differences.

Results are shown in Table 4. Note that  $\Delta$ After stands for  $a_t - a_{t-1}$ ,  $\Delta$ Diagnosed stands for  $d_t - d_{t-1}$ , and  $\Delta$ AMI stands for  $h_t - h_{t-1}$  in the table. Column 1 shows the simplest specification, with controls only for diagnosis with CAD and treatment with either PCI or CABG. The  $\Delta$ Diagnosed indicator is equal to one in the period when MM patients, PCI patients, or CABG patients are diagnosed, whether the diagnosis occurs in the same period as the treatment or not.<sup>14</sup> Being diagnosed with CAD is associated with a 5.0 percentage point increase in the probability of quitting, on top of the typical yearly quit rate of 4.9 percent (reflected in the coefficient on the constant term). Being treated with PCI or CABG is associated with an incremental 9.0 or 22.5 percentage point increase in the quit rate, respectively. This assumes that diagnosis and treatment occur in the same period. If not, the diagnosis coefficient must be subtracted since  $\Delta$ Diagnosed equals zero for PCI and CABG patients diagnosed before treatment but equals one for all MM patients. That results in a 4.0 percentage point increase for PCI patients and a 17.5

---

<sup>14</sup> In results not shown, we find no evidence of a *differential* diagnosis effect based on treatment received. Therefore, in all results shown, we assume that there is a single diagnosis effect that does not vary by treatment.

percentage point increase for CABG patients. Since 40 percent of PCI patients and 43 percent of CABG patients are diagnosed before treatment, the average percentage point increases in the quit rates are 7.0 and 20.4, respectively.<sup>15</sup> Regardless of how these computations are made, the difference between the coefficients on PCI and CABG treatments is 13.5 percentage points and is significant at the  $\alpha = 0.05$  level.

Table 4 – Regression Results for Discrete Time Linear Probability Hazard Model

	(1)	(2)	(3)	(4)	(5)
Constant	0.049*** (0.001)	0.049*** (0.001)	0.047*** (0.001)	0.046*** (0.011)	-0.050 (0.032)
PCI * ΔAfter	0.090** (0.038)	0.054 (0.040)	0.034 (0.040)	0.033 (0.039)	0.026 (0.042)
CABG * ΔAfter	0.225*** (0.045)	0.195*** (0.045)	0.160*** (0.044)	0.155*** (0.044)	0.141*** (0.046)
ΔDiagnosed	0.050*** (0.008)	0.042*** (0.008)	0.025*** (0.008)	0.025*** (0.008)	0.004 (0.026)
ΔAMI		0.083*** (0.023)	0.075*** (0.024)	0.076*** (0.024)	0.076*** (0.024)
CABG * ΔAfter - PCI * ΔAfter	0.135**	0.141**	0.126**	0.122**	0.115**
Elixhauser			X	X	X
Demographics				X	X
Year, Period, Age, Yrs Smoked					X
Observations	26,658	26,658	26,658	26,658	26,658
Individuals	3,065	3,065	3,065	3,065	3,065

Robust standard errors, clustered at the individual level, in parentheses. Regressions are weighted by NHIS probability weights. ΔAMI is an indicator for a patient having his first AMI during a particular period. Specifications 3 and 4 include 29 dummy variables for the Elixhauser commorbidity conditions first diagnosed during a particular period. In specification 4, demographic controls include gender, race, education dummies, and income category dummies (including a dummy for missing income data). In specification 5, we also include year dummies, period dummies, age and age squared, and a variable indicating the number of years a person had been smoking as of the baseline period (9.5 years before treatment). "P-Value: PCI \* ΔAfter = CABG \* ΔAfter" is the p-value on a test of equality of the PCI \* ΔAfter and CABG \* ΔAfter coefficients. \* significant at the 0.10 level; \*\* significant at the 0.05 level; \*\*\* significant at the 0.01 level.

<sup>15</sup> For MM patients, the increase in the quit rate is 0.0501. Since 60 percent of PCI patients are diagnosed in the same period as treatment, the average predicted increase in the quit rate for these patients is  $0.0897 + 0.60 * 0.0501 = 0.1198$ . The difference between that increase and the increase for MM patients is 0.0697 or 7.0 percentage points. Since 57 percent of CABG patients are diagnosed in the same period as treatment, the average predicted increase is  $0.2250 + 0.57 * 0.0501 = 0.2536$ . The difference between that increase and the increase for MM patients is 0.2035 or 20.4 percentage points.

In columns 2-5 of Table 4, we add additional control variables that may be associated with both selection into treatment and the propensity to quit smoking. In column 2, we add a control for a patient's first AMI. In column 3, we further control for the Elixhauser comorbidity conditions. In column 4, we add controls for demographic characteristics. And, in column 5, we add controls for the period, the calendar year, an individual's age, and the number of years he has smoked as of the baseline period. An AMI proves to be a strong predictor of quitting smoking, increasing the predicted probability of quitting by between 7.5 and 8.3 percentage points, depending on the specification. Moreover, once we add AMI, the estimated size of the PCI coefficient declines and is no longer statistically significant at conventional levels. In our final specification, we find that patients undergoing PCI are 2.6 percentage points more likely to quit smoking, though given the imprecision of our estimate, we cannot rule out the possibility of no association.<sup>16</sup> While the coefficient on CABG also declines, it remains large and strongly statistically significant, both in comparison to medically managed patients and PCI patients. In our fifth specification, undergoing CABG surgery in the same period in which CAD was diagnosed is associated with a roughly 14.1 percentage point increase in the probability of quitting smoking. The increased quit rate associated with CABG is substantially larger than the increase associated with less invasive treatments for CAD, and approximately twice the increase associated with an AMI. Moreover, in every specification, the difference between the CABG and PCI coefficients is consistently large – ranging from 0.115 to 0.141 – and significant at the 0.05 level.

---

<sup>16</sup> As discussed above, this estimate is for a PCI patient who was diagnosed and treated in the same period.

### *Model 2: Multi-Period Quit Model Using Grouped Data*

To illustrate that our results are not sensitive to a flexible specification of period effects and to account for clustering of disturbance terms by group and period at the individual level, we aggregate the data into 11 periods for each of the three groups of patients. There are 33 cells in the aggregate sample. We obtain two quit series. The first, shown in Panel A of Table 5, is unadjusted for covariates. It is identical to the quit rate that appears in Figure 3. The second quit series, shown in Panel B of Table 5, adjusts for effects due to diagnosis, AMI, and Elixhauser comorbidities. It is obtained from the individual data by estimating a discrete time hazard function for the probability of quitting that includes 11 period dummies interacted with each of three treatment dummies (one for CABG, one for PCI, and one for MM), and the diagnosis, AMI, and Elixhauser variables defined in equation (4). The 33 coefficients associated with the period-treatment interactions are quit rates by group and period adjusted for the effects of the last three variables just mentioned.<sup>17</sup>

---

<sup>17</sup> We do not adjust for demographic and socioeconomic characteristics since the inclusion of these characteristics has a very minor impact on the estimates in Table 5.

Table 5 – Grouped Data on Smoking Participation and Quit Rate

A. Unadjusted Quit Rate

Period	Quit Rate			Difference in Quit Rate			$\Delta$ After
	(1)	(2)	(3)	(4)	(5)	(6)	
	CABG	PCI	MM	CABG - PCI	CABG - MM	PCI - MM	
-9	6.2%	4.1%	5.1%	2.2%	1.2%	-1.0%	0
-8	5.1%	6.0%	4.2%	-1.0%	0.8%	1.8%	0
-7	4.2%	5.7%	4.3%	-1.5%	-0.1%	1.4%	0
-6	2.7%	3.9%	4.6%	-1.2%	-2.0%	-0.8%	0
-5	4.5%	7.7%	4.4%	-3.2%	0.1%	3.3%	0
-4	4.5%	1.7%	5.1%	2.8%	-0.6%	-3.4%	0
-3	8.1%	5.4%	4.6%	2.6%	3.5%	0.9%	0
-2	7.5%	3.9%	5.4%	3.6%	2.1%	-1.5%	0
-1	11.4%	9.0%	5.5%	2.4%	6.0%	3.5%	0
0	30.1%	17.4%	10.1%	12.7%	20.0%	7.3%	1
1	7.0%	7.8%	5.3%	-0.7%	1.8%	2.5%	0

B. Adjusted Quit Rate

Period	Adjusted Quit Rate			Difference in Adj. Quit Rate			$\Delta$ After
	CABG	PCI	MM	CABG - PCI	CABG - MM	PCI - MM	
-9	6.2%	4.1%	5.1%	2.2%	1.2%	-1.0%	0
-8	5.1%	6.0%	4.2%	-1.0%	0.8%	1.8%	0
-7	4.2%	5.7%	4.3%	-1.5%	-0.1%	1.4%	0
-6	2.6%	3.8%	4.6%	-1.2%	-2.0%	-0.8%	0
-5	4.4%	7.6%	4.4%	-3.1%	0.1%	3.2%	0
-4	4.4%	1.3%	5.1%	3.1%	-0.6%	-3.8%	0
-3	7.7%	5.4%	4.5%	2.3%	3.2%	0.9%	0
-2	6.8%	3.6%	5.2%	3.2%	1.5%	-1.7%	0
-1	10.1%	8.0%	5.1%	2.2%	5.0%	2.9%	0
0	22.6%	10.6%	8.4%	12.1%	14.3%	2.2%	1
1	5.6%	6.7%	4.1%	-1.1%	1.6%	2.6%	0

Note: Data are weighted by the NHIS probability weights. Adjusted quit rate adjusts for diagnosis, AMI, and Elixhauser comorbidities.

In the spirit of Donald and Lang (2007), we use this data to perform simple difference-in-differences regressions with 11 observations. To illustrate the model that we estimate, consider a log smoking participation function for two groups ( $g = c$  or  $p$ ,  $c = \text{CABG}$ ,  $p = \text{PCI}$ ):

$$\ln s_{gt} = \mu + \rho c + \beta c a_t + 11 \text{ period dummies} + \varepsilon_{gt}. \quad (5)$$

Here  $a_t$ , as defined in equation (3), is an indicator that equals 1 in each of the two periods after treatment and  $\varepsilon_{gt}$  is the error term. Take the difference between each group in a given period to eliminate the intercept ( $\mu$ ) and the period dummies. Then take first differences to eliminate the group effect ( $\rho$ ):

$$\ln s_{ct} - \ln s_{ct-1} - (\ln s_{pt} - \ln s_{pt-1}) \cong q_{ct} - q_{pt} = \beta c(a_t - a_{t-1}) + \text{error}. \quad (6)$$

Equation (6) is a regression forced through the origin with 11 observations. The dependent variable is the difference between the quit rate of CABG patients and the quit rate of PCI patients in each period. The independent variable,  $(a_t - a_{t-1})$ , equals 1 in the window spanning the period from 6 months before treatment to six months after treatment (period 10) and equals 0 in each of the other 10 periods or quit windows.

This approach has a number of attractive features. First, aggregation accounts for clustering in the disturbance term in an individual-level log smoking participation or quit equation by group and period. Second, if the error term in equation (5) is a random walk, then serial correlation is eliminated once first differences are taken. Third, the regression specified by equation (6) implicitly controls for a full set of period effects. Finally, by focusing on the



difference in the quit rates in each period, we are asking whether this difference during the treatment period is sufficiently unusual compared to past and future values that it is unlikely to have arisen by chance. If the quit rates in the two series normally track one another but do not during the treatment year, we would expect that there is something unusual about the treatment year. On the other hand, if the quit rates in the two series often diverge wildly, then a substantial divergence in the treatment year might simply be due to chance.

Six aggregate quit regressions are contained in Table 6. The three in the top row employ the unadjusted quit series, while the three in the bottom row employ the adjusted quit series. In column 1, the dependent variable is the difference between the CABG and PCI quit rates; in column 2, it is the difference between the CABG and MM rates; and in column 3, it is the difference between the PCI and MM rates. Three separate regressions are obtained for each series because of evidence that the residual variance is not the same for each dependent variable.<sup>18</sup> To be consistent with the notation in Table 4, the variable  $a_t - a_{t-1}$  is termed  $\Delta$ After in the table.

---

<sup>18</sup> Consider the following two regressions

$$q_{pt} - q_{pt-1} - (q_m - q_{mt-1}) = \beta_p(a_t - a_{t-1})$$

$$q_{ct} - q_{ct-1} - (q_{mt} - q_{mt-1}) = \beta_c(a_t - a_{t-1}),$$

where m denotes medical management. Estimates of  $\beta_p$ ,  $\beta_c$ , and  $\beta_c - \beta_p$  could be obtained from a pooled regression of the form

$$q_{gt} - q_{gt-1} - (q_m - q_{mt-1}) = \beta_c c(a_t - a_{t-1}) + \beta_p(1 - c)(a_t - a_{t-1}).$$

We do not follow that approach because the residual variance in the first regression is not equal to the corresponding variance in the second regression.

Table 6 – Quit Rate Regression with Grouped Data

	(1) CABG - PCI	(2) CABG - MM	(3) PCI - MM
$\Delta$ After (no adjustments)	0.127** (0.050)	0.200*** (0.044)	0.073 (0.047)
$\Delta$ After (with adjustments)	0.120** (0.052)	0.142*** (0.042)	0.022 (0.051)
N	11	11	11

Note: Each cell represents the coefficient on  $\Delta$ After from a separate regression, with the standard error in parentheses. The dependent variable is the quit rate for one group of patients minus the quit rate for another. The independent variable is a dummy for the treatment year ( $\Delta$ After). In the top row, the quit rate is unadjusted. In the bottom row, the quit rate is adjusted for diagnosis, AMI, and Elixhauser comorbidities, based on an individual regression. All quit rates are calculated using the NHIS probability weights. Regressions are forced through the origin and weighted by the inverse of the square root of the variance of the difference between period-specific quit rates from individual level regressions with 33 group period interactions. \* significant at the 0.10 level; \*\* significant at the 0.05 level; \*\*\* significant at the 0.01 level.

Focusing on the first column of Table 6, one sees that the difference between the CABG quit rate and the PCI quit rate in treatment period is between 12 and 13 percentage points, regardless of whether one uses the raw data or adjusts the quit rate for covariates. In the second column, we see that the difference in the quit rate between CABG and MM patients drops when one adjusts for covariates, but remains large (14.2 percentage points) and strongly statistically significant. In the final column, we see that while the unadjusted quit rate for PCI patients is 7.3 percentage points larger than for MM patients, once one adjusts for a patient’s CAD diagnosis and other medical conditions, the difference between the two quit rates is substantially smaller and not statistically distinguishable from zero.

## V. Conclusion

Coronary Artery Disease is a frequently occurring and deadly disease. There are several common treatments – including medical management, PCI, and CABG – and each has benefits and costs associated with it. In this paper, we have examined one previously unexplored benefit of more invasive treatment: those who undergo a procedure, particularly the more invasive CABG surgery, are more likely to quit smoking. In our preferred regression model, we estimate that CAD patients who undergo CABG are 12 percentage points more likely to quit smoking in the one-year window surrounding their surgery than patients who undergo PCI. During the same one-year window, CABG patients are 14 percentage points more likely to quit smoking than medically managed patients. These results are robust to a number of alternative specifications.

While we do not have data on behaviors other than smoking, we suspect that patients undergoing more invasive surgery are also more likely to improve their diet, limit excessive consumption of alcohol, and (when recommended) exercise more. Taken together, these behavioral responses may offset the inherent risks of a more invasive surgery and help explain why the longer term outcomes for CABG patients exceed those of similar patients receiving PCI. Our findings also highlight the importance of emphasizing healthier behavior to those patients who have less invasive medical treatment.

## VI. References

- Cutler, David M. and Robert S. Huckman (2003). "Technological development and medical productivity: the diffusion of angioplasty in New York state." *Journal of Health Economics*, 22 (2): 187-217.
- Donald, Stephan G and Kevin Lang (2007). "Inference with Difference-in-Differences and Other Panel Data." *The Review of Economics and Statistics*, 89(2): 221–233
- Dave, Dhaval M. and Robert Kaestner (2009). "Health Insurance and Ex Ante Moral Hazard: Evidence from Medicare." *International Journal of Health Care Finance and Economics*, 9 (4), 367-390.
- Elixhauser Anne, Claudia Steiner, D. Robert Harris and Rosanna M. Coffey. "Comorbidity Measures for Use with Administrative Data." *Med Care* 1998 36:8-27.
- Hlatky, M.A., Boothroyd, D.B., Bravata, D.M., et al. (2009). "Coronary artery bypass surgery compared with percutaneous coronary interventions for multivessel disease: a collaborative analysis of individual patient data from ten randomised trials." *Lancet* 373: 1190–97
- Mohr, Friedrich W. Marie-Claude Morice, A. Pieter Kappetein et al., (2013). Coronary artery bypass graft surgery versus percutaneous coronary intervention in patients with three-vessel disease and left main coronary disease: 5-year follow-up of the randomised, clinical SYNTAX trial." *Lancet* 381: 629–38.
- Murphy SL, Xu JQ, Kochanek KD (2012). "Deaths: Preliminary Data for 2010." *National Vital Statistics Reports*; vol 60 no 4. Hyattsville, MD: National Center for Health Statistics
- Peltzman, Sam. (1975). "The Effects of Automobile Safety Regulation." *Journal of Political Economy* 83: 677-725.
- Peltzman, Sam. (2011). "Offsetting Behavior, Medical Breakthroughs, and Breakdowns." *Journal of Human Capital*, 5(3): 302-341.
- Weintraub, William S., Maria V. Grau-Sepulveda, Jocelyn M. Weiss, et al. (2012). "Comparative Effectiveness of Revascularization Strategies." *New England Journal of Medicine* 366: 1467-76

## Appendix – NHIS / Medicare Data Linkage

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
<b>NHIS 1994-1998</b>																	
NHIS				1994	1995	1996	1997	1998									
Medicare	1991	1992	1993	1994	1995	1996	1997	1998									
<b>NHIS 1999-2005</b>																	
NHIS									1999	2000	2001	2002	2003	2004	2005		
Medicare									1999	2000	2001	2002	2003	2004	2005		

Note: the Medicare years labeled on the chart are potentially useful for our study because they represent a Medicare record that is linked to a later NHIS interviews. Medicare records linked to earlier NHIS interviews provide no information on quitting behavior after CAD treatment.

For those respondents who were diagnosed with CAD prior to their NHIS interview date, we have the ability to look at their smoking behavior before and after their diagnosis. For the subset of CAD patients who underwent PCI or CABG, we can also look at their smoking behavior before and after their procedure. For example, for individuals interviewed in 1994 who had PCI, we can look at their smoking behavior before and after their procedure *only if they underwent PCI between 1991 and 1994* (and within 1994, only if their procedure was before the date of the NHIS interview). If a person had PCI before 1991, then we have no record of their procedure. If a person had PCI after 1994, then we have no record of their smoking behavior *after* their procedure.

Each person in the linked dataset, therefore, has a “diagnosis window” within which they must be diagnosed with CAD to be included in our study. The longest window is for a person who was interviewed in 1998 – he will be included in our study if he was enrolled in Medicare and diagnosed with CAD at any point between 1991 and 1998. The shortest window is for a person who was interviewed in early 1999 – he will be included only if he was enrolled in Medicare and diagnosed with CAD on an earlier date in 1999 than the date of his interview.

Figure A1 – Characteristics by Treatment for Regression Sample

Demographic Characteristics				Medical Conditions			
	MM	PCI	CABG		MM	PCI	CABG
<i>Age</i>				<i>First AMI Within 6 Months of Treatment</i>			
< 55	11%	13%	6%	Yes	9%	51%	40%
55-64	15%	17%	15%	No	91%	49%	60%
65-69	31%	32%	35%	<i>% With Comorbidity Within 6 Months of Treatment</i>			
70-74	22%	23%	24%	Congestive heart failure	14%	28%	36%
75-79	14%	10%	14%	Valvular disease	10%	18%	24%
80-84	5%	3%	5%	Pulmonary circulation disorder	2%	5%	4%
85+	2%	1%	0%	Peripheral vascular disorder	13%	22%	30%
<i>Gender</i>				Paralysis	2%	1%	3%
Male	49%	54%	64%	Other neurological	4%	4%	4%
Female	51%	46%	36%	Chronic pulmonary disease	26%	18%	35%
<i>Race</i>				Diabetes w/o chronic comp.	14%	15%	22%
Asian	0%	0%	1%	Diabetes w/ chronic comp.	5%	6%	11%
Black	11%	8%	6%	Hypothyroidism	6%	9%	3%
Hispanic	4%	2%	3%	Renal failure	2%	3%	5%
White	83%	87%	89%	Liver disease	1%	1%	0%
Mult./Oth/Unknown	1%	2%	2%	Chronic Peptic ulcer disease	0%	1%	1%
<i>Education</i>				HIV and AIDS	0%	0%	0%
Elem (K-8)	20%	23%	21%	Lymphoma	0%	1%	1%
HS (non-grad); GED	23%	21%	22%	Metastatic cancer	1%	0%	0%
HS grad	29%	33%	30%	Solid tumor without metastasis	5%	4%	3%
Some col; AA deg.	18%	19%	15%	Rheumatoid arthritis	3%	2%	1%
BA degree	6%	1%	8%	Coagulation deficiency	2%	4%	9%
Grad. Degree	3%	2%	4%	Obesity	4%	7%	8%
Unknown	1%	0%	1%	Weight loss	2%	2%	2%
<i>Family Income</i>				Fluid and electrolyte disorders	11%	18%	28%
\$0 to \$9,999	23%	23%	16%	Blood loss anemia	1%	2%	5%
\$10,000 to \$19,999	28%	21%	24%	Deficiency anemias	10%	12%	29%
\$20,000 to \$35,000	19%	25%	22%	Alcohol abuse	2%	2%	4%
\$35,000 or over	14%	14%	23%	Drug abuse	1%	1%	2%
Unknown	16%	16%	15%	Psychoses	2%	1%	1%
				Depression	6%	11%	8%
				Hypertension	38%	44%	44%
<i>Count (N = 3,065)</i>	2,657	199	209	<i>Count (N = 3,065)</i>	2,657	199	209

Note: Sample includes only those individuals who were included in the regression because they smoked 9.5 years prior to the initiation of their treatment. All data are weighted by the NHIS probability weights. Age is as of diagnosis (CAD) or procedure (PCI / CABG). Comorbidities based on Elixhauser et al. (1998)