

This PDF is a selection from an out-of-print volume from the National Bureau of Economic Research

Volume Title: Annals of Economic and Social Measurement, Volume 6, number 5

Volume Author/Editor: NBER

Volume Publisher: NBER

Volume URL: <http://www.nber.org/books/aesm77-5>

Publication Date: December 1977

Chapter Title: Adapted Methods for Solving and Optimizing Quasi-Triangular Econometric Models

Chapter Author: Pierre Nepomiestehy, Alain Ravelli

Chapter URL: <http://www.nber.org/chapters/c10543>

Chapter pages in book: (p. 555 - 562)

## ADAPTED METHODS FOR SOLVING AND OPTIMIZING QUASI-TRIANGULAR ECONOMETRIC MODELS

BY PIERRE NEPOMIASTCHY AND ALAIN RAVELLI

*This paper sets out a Newton-like method for solving the model and a method for computing the gradient (adjoint variable technique) which are both adapted to econometric models with a quasi-triangular structure (after a possible renumbering of the equations). Comparisons are made with classical methods on the 130 equations Star model.*

### I. INTRODUCTION

Let us consider a non-linear and dynamic macroeconomic model with  $n$  endogeneous variables. To solve this model, we propose the following method. A set of  $s$  variables, called *loop variables*, and a set of  $s$  equations, called *loop equations* are selected such as: for given values of the loop variables, the remaining model with  $n - s$  equations and  $n - s$  variables is triangular and can be solved directly; then, an algorithm is chosen to iterate on the values of the loop variables in order to satisfy the loop equations.

The efficiency of this method obviously depends on the number of loop variables. We propose in [1] a method to renumber the equations and the variables in order to minimize this number.

If, after a possible renumbering,  $s$  is small compared to  $n$ , then the jacobian matrix of the model is lower-quasi-triangular (i.e. most of its non zero elements are under the main diagonal) and the model is called *quasi-triangular*. In the paper, we shall consider only this kind of model but it seems to be the case for many macroeconomic models. Here are a few examples:

Model	Number of equations ( $n$ )	Number of loop variables ( $s$ )
Andomini [2]	4	1
Pimpon [3]	14	2
Fair [4]	83	7
Star [5]	130	3
DMS [6]	about 1000	less than 100

The quasi-triangular structure was obtained by hand for Andomini and Pimpon, by the algorithm described in [1] for Fair, and was provided by the author of Star; for DMS, no renumbering was made.

In section III, we propose an adapted Newton method easy to use and we give numerical results using comparisons with Gauss-Seidel.

We have shown in [7] on the small Pimpon model that the adjoint variable method is more effective for computing the gradient than the generally used finite difference method proposed by Fair [8]. We propose in section IV an adaptation of this adjoint variable method to the quasi-triangular structure of the model and we give numerical results of comparisons with the finite difference method made using the Star model.

These comparisons provide most encouraging results to support the efficiency of the two proposed methods.

## II. STRUCTURE OF THE MODEL

For sake of clarity of the presentation, we shall simplify the structure of the model in the sense that we are not going to distinguish between equations which do not depend on loop variables and which correspond to pre-determined variables, equations which depend on loop variables (heart of the model) and the triangular set of output equations. The methods proposed in sections III and IV can easily be adapted to take into account these differences between equations (see [1]).

Let us consider a discrete time dynamic model with periods  $t$  going from 1 to  $T$ . Let  $n$  be the number of endogenous variables,  $x_t^i$  be the value of variable  $i$  at period  $t$  and  $x_t$  be the vector  $\{x_t^1, \dots, x_t^n\}$ . Let  $r$  be the number of control variables,  $u_t^i$  be the vector  $\{u_t^1, \dots, u_t^r\}$ . Let  $p$  and  $q$  be the maximum lag appearing in the model on (respectively) endogenous and control variables. Then, the model is described by the following set of equations:

$$(1a) \quad x_t^i = f_i^t(x_t, x_{t-1}, \dots, x_{t-p}, u_t, u_{t-1}, \dots, u_{t-q}), \\ i = 1, \dots, n, \quad t = 1, \dots, T$$

$$(1b) \quad x_t^i \text{ given for } i = 1, \dots, n \quad \text{and} \quad t = -p + 1, -p + 2, \dots, -1, 0$$

We shall assume that the given functions  $f_i^t$  are continuously differentiable and that, at least for any control taken in a reasonable range, the system (1) has a unique solution.

For solving the model at period  $t$ , the vector:

$$(2) \quad e_t = \{x_{t-1}, x_{t-2}, \dots, x_{t-p}, u_t, u_{t-1}, \dots, u_{t-q}\}$$

is known and the problem is to find the solution  $x_t$  of the problem:

$$(3) \quad x_t^i = f_i^t(x_t, e_t), \quad i = 1, \dots, n$$

After a possible renumbering of equations and variables, the loop equations are the  $s$  last equations of the model and  $y_t = \{x_t^{n-s+1}, \dots, x_t^n\}$  is the vector of the loop variables. Then, by definition of the loop variables, the system (3) is triangular for any given  $e_t$  and (3) can be

written:

$$(4a) \quad x_i^t = f_i^t(x_1^{t-1}, \dots, x_{i-1}^{t-1}, y_i, e_i), \quad i = 1, \dots, n - s$$

$$(4b) \quad x_i^t = f_i^t(x_i, e_i), \quad i = n - s + 1, \dots, n$$

### III. SOLUTION OF THE MODEL

For any given  $y_i$ , the system (4a) is triangular and can be solved by simple evaluations of functions  $f_i^t$ , which gives us values of  $x_i^t$  for any  $i \leq n - s$ . Since  $y_i = \{x_i^{n-s+1}, \dots, x_i^n\}$ , it can be seen that, taking (4a) into account,  $x_i$  can be considered as a function  $x_i(y_i)$  of  $y_i$ . Of course, with an arbitrary value of  $y_i$ ,  $x_i(y_i)$  does not in general satisfy equations (4b), there is an error which depends on and only on  $y_i$ . We shall denote  $\varphi_i(y_i)$  this error. The problem is to find  $y_i$  in such a way that this error is equal to zero:

$$(5) \quad \varphi_i^t(y_i) = 0, \quad i = 1, \dots, s$$

*In theory*, it is possible to eliminate variables  $x_i^t$ ,  $i < n - s$ , from equations (4) and obtain the *analytical* expressions of functions  $\varphi_i^t(y_i)$ . For large models, this is obviously impossible for practical reasons but it is clear that, using (4), one is able to compute the numerical value of  $\varphi_i(y_i)$  for any given numerical value of  $y_i$ . In this case, the partial derivatives of  $\varphi_i^t$  cannot be analytically computed and the simplest method for solving (5) is the well-known Gauss-Seidel algorithm.

It should be noted that this Gauss-Seidel algorithm, applied to the system (5) of dimension  $s$  ( $s = 3$  for Star) and not to the total system (3) of dimension  $n$  ( $n = 130$  for Star) is adapted to the quasi-triangular structure of the model. But, still, it has the usual disadvantages of the Gauss-Seidel methods, namely:

- a) the convergence is slow;\*
- b) the convergence is strongly dependent on the ordering and normalization of the equations and on the weights chosen for the feedback, weights which can be determined for each model only by a large number of random tests.

We propose solving (5) by using the Newton method which has a convergence rate of 2\*\* and no parameters to determine in the feedback rule. It should be noted that if the Newton method had been applied directly to the total model (3), then each iteration would have required the solution of a linear system of dimension  $n$ . Using a general package

\*Its convergence rate is 1: the error at iteration  $k$  is proportional to the error at iteration  $k - 1$ .

\*\*The error at iteration  $k$  is proportional to the square of the error at iteration  $k - 1$ .

to solve this linear system, we have obtained very poor results. This linear system is sparse and can be solved by adapted methods like those proposed by Drud [9] and we shall wait until Drud's routine is available to see if the "global Newton method" with the sparse technique for solving the linear system is more effective than our method or not.

Applying the Newton method to (5), at each iteration we have only to solve a linear system of dimension  $s$  (with  $s = 3$  for Star). Consequently, the only problem is the computation of the partial derivatives  $\partial \varphi_i^t / \partial y_i^t$ . If an analytical derivation package (as Formac) is available, then the analytical expressions of the derivatives  $\partial f_i^t / \partial x_i^t$  can be deduced from (4). By definition of  $\varphi_i^t(y_i)$ , we have:

$$(6) \quad \varphi_i^t(y_i) = y_i^t - f_i^{n-s+t}[x_i^t(y_i), e_i], \quad i = 1, \dots, s$$

with  $y_i^t = x_i^{n-s+t}$ ; from (4a) we have:

$$(7) \quad \frac{\partial x_i^t}{\partial y_i^t} = \sum_{k=1}^{i-1} \frac{\partial f_i^t}{\partial x_i^k} \frac{\partial x_i^k}{\partial y_i^t} + \frac{\partial f_i^t}{\partial y_i^t}, \quad i = 1, \dots, n-s, \quad j = 1, \dots, s$$

which for any  $j$ , is a triangular system in  $\partial x_i^t / \partial y_i^t$ ,  $i = 1, \dots, n-s$ .

Then, from derivation of (6), we have:

$$(8) \quad \frac{\partial \varphi_i^t}{\partial y_i^t} = \delta_{ij} - \sum_{k=1}^{n-s} \frac{\partial f_i^{n-s+t}}{\partial x_i^k} \frac{\partial x_i^k}{\partial y_i^t} - \frac{\partial f_i^{n-s+t}}{\partial y_i^t}, \quad i, j = 1, \dots, s$$

If no analytical derivation package is available, then finite difference approximations of the  $\partial \varphi_i^t / \partial y_i^t$  are computed. The computation of  $\varphi_i^t(y_i)$  for  $s+1$  values of vector  $y_i$ , that is  $s+1$  solutions of the triangular system (4), gives an approximation of these derivatives; for more details, see [1].

With programs written in Fortran H extended on an IBM 370-168 computer and with a unit of one millisecond of CPU time, Table 1 gives, for different values of the required accuracy, the computing time of one simulation over 10 periods of the 130 equations model Star.

TABLE 1

$\epsilon$	GS	N1	N2
$10^{-2}$	14.7	9.6	12.0
$10^{-4}$	51.9	14.5	18.5
$10^{-6}$	129.0	20.2	20.6
$10^{-8}$	265.9	21.9	26.3

GS: Gauss-Seidel method.

N1: Newton method with analytical derivation of the model.

N2: Newton method with numerical derivation of the model.

$\epsilon$ : accuracy required.

Remember that the model is solved when  $\varphi_t(y_t) = 0$ , with  $\varphi_t(y_t)$  given by formula (6). We have chosen as the test to stop the algorithm:

$$(9) \quad \sum_{t=1}^s \left| \frac{f_t^{n-s+t} - y_t^t}{y_t^t} \right| \leq \epsilon$$

All the methods tested were adapted to the quasi-triangular structure of the model. A run of the "global Newton method" discussed above with a standard package for the linear system produced disastrous results: 13 *seconds* for  $\epsilon = 10^{-8}$  (50 times worse than GS).

The Gauss-Seidel method chosen for comparison was the one described above, namely Gauss-Seidel applied to (5), *after optimization* of the  $s = 3$  weights of the feed-back rule. It is obvious that Gauss-Seidel applied to the total system (3), even after the optimization of the  $n = 130$  weights of the feed-back rule, would have given worse results.

In methods N1 and N2, the jacobian ( $(\partial\varphi_t^i/\partial y^j)$ ) was computed for each iteration of the Newton methods. Computing it only sometimes gives a small improvement (for more details, see [1]).

From table 1, it appears obvious that when the Newton method is adapted to the quasi-triangular structure of the model, then this method is much more effective than the Gauss-Seidel, especially when a high accuracy is required, which is the case when the simulation algorithm is only a part of an optimization problem solved by sophisticated algorithms like Davidon-Fletcher-Powell.

#### IV. OPTIMIZATION OF THE MODEL

We have to minimize the following loss function:

$$(10) \quad j(x, u) = \sum_{t=1}^T j_t(x_t, x_{t-1}, \dots, x_{t-p}, u_t, u_{t-1}, \dots, u_{t-q})$$

where  $x$  and  $u$  are linked by the model (1). We shall assume that the functions  $j_t$  are continuously differentiable.

In this paper, we are only concerned by the search for an efficient algorithm to compute the gradient  $J'(u)$  of  $J(u) = j[x(u), u]$ , where  $x(u)$  is the solution of (1) associated with the control  $u$ .

The finite difference method [8] is a very simple method for gradient computation, but it requires  $rT$  model simulations for one gradient computation, where  $r$  is the control dimension and  $T$  the number of periods. We propose using the adjoint-variable method. This method is well known in optimal control theory (see, for example [10]) but, as far as we know, was applied to the optimization of a macroeconomic model for the first time by us [7]. In [7], we have shown how it can be applied to the 14-equations Pimpon model: here we show how it can be adapted to the

quasi-triangular structure of the model and we give results for the 130-equations Star model.

To describe the adjoint-variable method, let us introduce several notations:

$$(11) \quad \delta_t = \begin{cases} 1 & \text{if } t < T \\ 0 & \text{if } t = T \end{cases}$$

$$(12) \quad y_t^i = \sum_{k=0}^p \delta_{t+k} \frac{\partial j_{t+k}}{\partial x_t^i}, \quad i = 1, \dots, n, \quad t = 1, \dots, T$$

$$(13) \quad w_t^i = \sum_{k=0}^q \delta_{t+k} \frac{\partial j_{t+k}}{\partial u_t^i}, \quad i = 1, \dots, r, \quad t = 1, \dots, T$$

$$(14) \quad F_t^A = \left( \left( \frac{\partial f_{t+k}^i}{\partial x_t^i} \right) \right); \quad G_t^A = \left( \left( \frac{\partial f_{t+k}^i}{\partial u_t^i} \right) \right)$$

Let  $\psi = \{\psi_1, \dots, \psi_T\}$ , with  $\psi_t \in R^n$ , be the solution of:

$$(15) \quad \psi_t = \sum_{k=0}^p \delta_{t+k} (F_t^A)^* \psi_{t+k} + y_t, \quad t = 1, \dots, T$$

where  $(F_t^A)^*$  is the transposed matrix of  $F_t^A$ . The equation (15) is called the adjoint system of the optimization problem and its solution  $\psi$  is called the adjoint-variable. It can be proven (see [1]) that the gradient can be deduced from the value of  $\psi$  by the formula:

$$(16) \quad J_t'(u) = w_t + \sum_{k=0}^q \delta_{t+k} (G_t^A)^* \psi_{t+k}, \quad t = 1, \dots, T$$

For the gradient computation, one must solve the system (15) for  $t = T$ , then for  $t = T - 1$ , and so on till  $t = 1$ . Indeed, when  $\psi_t$  is computed, the values of  $\psi_{t'}$ , for any  $t' > t$  are known, hence the vector:

$$(17) \quad c_t = y_t + \sum_{k=1}^p \delta_{t+k} (F_t^A)^* \psi_{t+k}$$

is known (from (11) we see that, for  $t = T$ ,  $c_t = y_t$ ) and computing  $\psi_t$  is reduced to the solution of the linear system:

$$(18) \quad \psi_t = (F_t^A)^* \psi_t + c_t$$

From the notation (14) and the structure of the model given by (4), it can be seen that (18) can be written:

$$(19a) \quad \psi_t^i = \sum_{j=i+1}^n \frac{\partial f_t^j}{\partial x_t^i} \psi_t^j + c_t^i, \quad i = 1, \dots, n - s$$

$$(19b) \quad \psi_i^t = \sum_{j=1}^n \frac{\partial f_j^t}{\partial x_i^t} \psi_j^t + c_i^t, \quad i = n - s + 1, \dots, n$$

Using the same argument as for solving the model, let  $\lambda = \{\psi_1^{n-s+1}, \dots, \psi_s^n\}$  be the vector of the adjoint loop variables. For any given  $\lambda$ , using (19a), it is possible to compute the corresponding values of  $\psi_i^{n-s}$ , then  $\psi_i^{n-s-1}$  and so on till  $\psi_i^1$  and check whether these values satisfy (19b). Consequently, we are able to compute the errors:

$$(20) \quad E_i(\lambda) = \lambda_i - \sum_{j=1}^n \frac{\partial f_j^t}{\partial x_i^t} \psi_j^t(\lambda) - c_i^{n-s+1}, \quad i = 1, \dots, s$$

and the only problem is to find  $\lambda$  such as  $E(\lambda) = 0$ . From (19), it is clear that  $\lambda \rightarrow E(\lambda)$  is a linear mapping which can be denoted by  $E(\lambda) = A\lambda - b$ . The computation of  $E(\lambda)$  for  $\lambda = 0$  gives the vector  $b$ . It can be proven\* that the matrix  $A$  is the transposed of the jacobian matrix of the system (5), hence  $A$  is known. Consequently, the system  $A\lambda = b$ , which is only of dimension  $s$ , has to be solved, then, the adjoint variable  $\psi_i$  is computed with the help of (19a) and, finally, the gradient is obtained by the formula (16).

The adjoint variable method requires, for one gradient computation,  $2T$  evaluations of equations (19) and  $T$  solutions of a linear system of dimension  $s$ ; on the other hand, the finite difference method requires  $rT^2/2$  solutions of the non-linear system (3) of dimension  $n^{**}$ . Consequently, it is clear that the adjoint variable method is much more effective as shown on Table 2 which gives, for the model Star, the computing time\*\*\* of a gradient computation using the two methods:

TABLE 2

$T$	$M1$	$M2$	$M1/M2$
10	0.65 sec.	0.016 sec.	40.6
20	2.39 sec.	0.031 sec.	77.0
30	4.90 sec.	0.046 sec.	106.5

$T$  : number of simulation periods.

$M1$  : gradient computing time using the finite difference method.

$M2$  : gradient computing time using the adjoint variable method.

$M1/M2$ : ratio of computing times.

\*this result was kindly suggested to us by one of the referees of this paper. For proof, see [1].

\*\*for Star, we have  $r = 10$ ,  $n = 130$ ,  $s = 3$  and  $T = 10$ .

\*\*\*with programs in Fortran H extended on an IBM 370-168 computer.

In this comparison, the matrices  $F_1^A$  and  $G_1^A$  were obtained by analytical derivation of the equations of the model. A finite difference approximation of these matrices multiply the computing time of method  $M2$  by a factor 10 (independently of  $T$ ) and the adjoint variable method still remains much powerful than the finite difference method.

IRIA-LABORIA

Rocquencourt, 78150 Le Chesnay  
France

#### REFERENCES

- [1] D. Gabay; P. Néponiastchy; M. Rochdi; A. Ravelli: "Etude, résolution et optimisation de modèles macroéconomiques", Rapport IRIA, 1978.
- [2] K. Wall; *NBER Cambridge Memo*, June 26, 1975.
- [3] A. Coufiere: "Un modèle du système monétaire français," *Statistiques et Etudes Financières*, INSEE (Paris), 1975/17.
- [4] R. C. Fair: "A Model of Macroeconomic Activity," Vol. II. The Empirical Model, Ballinger Publishing Company, 1976.
- [5] J. Mistral: "STAR, Modèle de prévision à court/moyen terme," *Séminaire sur les modèles macroéconomiques* (CNRS-NSF), Gif/s/Yvette (France), Nov. 22-26, 1976.
- [6] D. Fouquet: "Les multiplicateurs de DMS," *Publ. INSEE*, Jan. 24, 1977.
- [7] P. Nepomiastchy: "Méthodes d'optimisation adaptées aux modèles macroéconomiques," *International Symposium on New Trends in System Analysis*, Versailles (France), Dec. 13-17, 1976.
- [8] R. C. Fair: "On the Solution of Optimal Problems as Maximization Problems," *Annals of Economic and Social Measurement*, Vol. 3, No. 1, Jan. 1974, pp. 135-154.
- [9] A. Drud: "Method for Control of Complex Dynamic Systems," *IMSOR*, No. 27, Lyngby (Denemark), 1976.
- [10] A. Bensoussan; E. Hurst; B. Naslund: "Management Applications of Modern Control Theory," North-Holland, 1974.