

NBER WORKING PAPER SERIES

INFORMAL CARE AND CAREGIVER'S HEALTH

Young Kyung Do
Edward C. Norton
Sally Stearns
Courtney H. Van Houtven

Working Paper 19142
<http://www.nber.org/papers/w19142>

NATIONAL BUREAU OF ECONOMIC RESEARCH
1050 Massachusetts Avenue
Cambridge, MA 02138
June 2013

The data for this study were made available by the Korea Labor Institute and Korea Employment Information Service. We thank David Grabowski and other participants at the International Health Economics Association's 8th World Congress on Health Economics and seminar participants at the University of Rochester and Johns Hopkins University for helpful comments. C.H. Van Houtven was supported by the Veterans Affairs Health Services Research and Development Merit Review Program (MRP 05-311). The views expressed herein are those of the authors and do not necessarily reflect the views of the National Bureau of Economic Research.

NBER working papers are circulated for discussion and comment purposes. They have not been peer-reviewed or been subject to the review by the NBER Board of Directors that accompanies official NBER publications.

© 2013 by Young Kyung Do, Edward C. Norton, Sally Stearns, and Courtney H. Van Houtven. All rights reserved. Short sections of text, not to exceed two paragraphs, may be quoted without explicit permission provided that full credit, including © notice, is given to the source.

Informal Care and Caregiver's Health

Young Kyung Do, Edward C. Norton, Sally Stearns, and Courtney H. Van Houtven

NBER Working Paper No. 19142

June 2013

JEL No. I1

ABSTRACT

This study aims to measure the causal effect of informal caregiving on the health and health care use of women who are caregivers, using instrumental variables. We use data from South Korea, where daughters and daughters-in-law are the prevalent source of caregivers for frail elderly parents and parents-in-law. A key insight of our instrumental variable approach is that having a parent-in-law with functional limitations increases the probability of providing informal care to that parent-in-law, but a parent-in-law's functional limitation does not directly affect the daughter-in-law's health. We compare results for the daughter-in-law and daughter samples to check the assumption of the excludability of the instruments for the daughter sample. Our results show that providing informal care has significant adverse effects along multiple dimensions of health for daughter-in-law and daughter caregivers in South Korea.

Young Kyung Do

Program in Health Services and Systems Research

Duke-NUS Graduate Medical School Singapore

8 College Road, Singapore 169857

young.do@duke-nus.edu.sg

*****Sally Stearns

*****Department of Health Policy and Management

*****The University of North Carolina at Chapel Hill

*****Chapel Hill, NC 27599-7411

*****sstearns@email.unc.edu

Edward C. Norton

Department of Health Management and Policy

Department of Economics

University of Michigan

School of Public Health

1415 Washington Heights, M3108 SPHII

Ann Arbor, MI 48109-2029

and NBER

ecnorton@umich.edu

*****Courtney H. Van Houtven

*****Department of Medicine

Division of General Intern

*****Duke University Medical Center

Durham, NC 27705, USA

courtney.vanhoutven@duke.edu

1. INTRODUCTION

While much economic research on informal care focuses on the health and health care use of an elderly parent, informal care may also adversely affect the health of the caregiver. Several mechanisms may lead to this effect (Schulz *et al.*, 1995). Caregiving can worsen health through increased emotional stress and physical strain. Caregiving also inevitably involves observing a loved one's decline and anticipatory bereavement, which itself may affect the caregiver's health (Bobinac *et al.*, 2010; Amirkhanyan and Wolf, 2006). Through these negative health effects, caregiving may increase a caregiver's health care use.

The provision of informal care may, however, be endogenous to caregivers' health, making the modeling of the endogeneity problem important to uncover the true effect (Coe and Van Houtven, 2009). Health status may affect the individual and family decisions on who provides informal care ("selection in") as well as whether the caregiver continues or ceases caregiving ("selection out") (Coe and Van Houtven, 2009). The selection criteria may not be monotonic in health. While a healthier family member is more likely to take up the caregiving role because of the burdensome nature of caregiving, it is also possible that a less healthy or less productive family member may choose to provide informal care instead of participating in paid employment. Selection out may arise when less healthy caregivers are more likely to stop providing informal care. This endogeneity issue makes it difficult to untangle the causal effect of informal caregiving on caregivers' health in observational data.

This study aims to measure the causal effect of caregiving on caregivers' health and health care use, using instrumental variables (IVs). We use nationally representative data from South Korea, where daughters and daughters-in-law are the prevalent source of caregivers for frail elderly parents and parents-in-law. A key insight is that having a parent-in-law with one or

more Activity of Daily Living (ADL) limitations increases the probability of providing informal care to that parent-in-law, but a parent-in-law's ADL limitation does not directly affect the daughter-in-law's health. We focus, therefore, on daughter-in-law caregivers and use the health of their parents-in-law as our instruments. We begin with a simple Wald estimator and perform careful robustness and falsification tests of the instruments to show that our instruments are theoretically and empirically valid in their application to informal caregiving in South Korea.

This paper makes several contributions to the empirical literature on caregiver health. First, we estimate the causal effect of caregiving on the health of the caregiver, controlling for the endogeneity between informal care and health using empirically strong instrumental variables. Second, we estimate how caregiving affects a caregiver's own health care use and expenditures to provide policy-relevant information about how the health effects of caregiving may translate to public health care costs in an aging society. Third, we use data from South Korea, which has a unique cultural context about filial duty and caregiving as well as an emerging strong market of publicly provided long-term care. The unique data source allows us to estimate caregiver health effects separately for daughters and daughters-in-law, the latter being an important source of informal care in South Korea. The comparison between daughters and daughters-in-law is interesting empirically.

We find that informal care has adverse effects on health and health care for daughter-in-law and daughter caregivers in South Korea. Caregiving has significant spillover effects to caregiver health care expenditures. For example, compared to their non-caregiving counterparts, caregiving daughters-in-law annually spend on average 47 percent more out-of-pocket on their own outpatient costs among those who use any outpatient care.

2. BACKGROUND

A large body of research has found negative health effects of caregiving (for reviews and meta-analyses see Schulz *et al.*, 1990; Schulz *et al.*, 1995; Walker *et al.*, 1995; Bookwala *et al.*, 2000; Yee and Schulz, 2000; Dilworth-Anderson *et al.*, 2002; Pinquart and Sörensen, 2003; Vitaliano *et al.*, 2003; Pinquart and Sörensen, 2006; Pinquart and Sörensen, 2007). Despite considerable differences in the study design and outcome measures, the literature finds that, compared with non-caregivers, caregivers experience overall poorer psychological and physical health (Schulz *et al.*, 1995; Pinquart and Sörensen, 2003; Vitaliano *et al.*, 2003).

The main empirical challenge of the literature on caregiver health effects has been that caregiving (or care intensity among caregivers) is potentially endogenous. In their critical review of the earlier literature, Barer and Johnson (1990) point out that much of the literature uses self-selected samples. Schulz (1990) also suggests that health status may determine who will provide informal care in the family. While well acknowledged, this methodological challenge has not been adequately addressed in the caregiving literature, with one recent exception (Coe and Van Houtven, 2009). Coe and Van Houtven (2009) address the endogeneity of both selection in and selection out of caregiving using different sets of instrumental variables. They use the death of a parent to instrument for selection out of caregiving, and sibling and family characteristics to instrument for selection into caregiving. In their longitudinal study, they find no evidence of endogeneity for selecting into caregiving, but do find evidence of endogeneity for selecting out of caregiving.

We extend the literature by taking into account the endogeneity of informal caregiving in the instrumental variable framework, using data from an Asian cultural setting — South Korea. South Korea provides a particularly interesting setting for our empirical work. Although South

Korea has become one of the fastest aging societies in the world, its strong tradition of filial piety means caregiving for frail elderly parents is the responsibility of adult children. Informal caregiving for disabled parents is embedded as an inseparable component of old-age support in traditional Korean culture (Sung, 1990), as in other East Asian cultures (Hsu and Shyu, 2003; Yamamoto and Wallhagen, 1997). Filial piety, an essential element of Confucianism, served as the major principle for the everyday lives of ordinary people, helping to keep the family as well as society in harmony (Chee, 2000). In an agrarian society until the 1960s, the Korean extended family served as a production unit as well as a communal living unit. Traditional Korean extended families continued with the eldest son's marriage and intergenerational co-residence with his parents. An eldest son inherited a larger share of bequests than his siblings would, in the form of house, farming land, and other real estate assets. The disproportionately larger bequest to the eldest son implied that the eldest son assumed the greatest role in familial responsibilities, which included co-residing with, supporting, and caring for elderly parents. The daughter-in-law married to the eldest son played a central role in familial responsibilities, including caring for ill and disabled parents. In a sense, the extended Korean family started with a designated future caregiver, the daughter-in-law, for aging parents (Choi, 1993).

Although the traditional form of parental caregiving is care by the eldest son's wife, recent decades have seen important changes in attitudes towards parental support. Eldest sons' wives now are less likely to assume their traditional role of parental caregiving than they were in the historically patrilineal society; at the same time, daughters increasingly play a greater role in caring for their own parents. Traditionally, daughters were viewed as leaving their family to join their husband's family after marriage, and therefore did not receive equal bequests from their own parents. A trend toward gender equality coupled with weakening traditional norms has

changed this pattern. Therefore, when a woman has both parents and parents-in-law who need care, it is now less clear for whom she will provide care than in the past. Still, older women, especially those married to an eldest son, are more likely to uphold the traditional practice of daughters-in-law sharing the responsibility of caregiving in the husband's family. A younger woman who is the eldest daughter in her pre-marriage family and lives close by her parents tends to play an increasingly greater role in parental caregiving than in the past, suggesting potential heterogeneity among women who serve as caregivers to their parents or their parents-in-law. When faced with the competing demands for caregiver time (i.e., a woman has both a parent and a parent-in-law with limitations), the decision for whom a woman will provide care involves a more complex set of factors, including her birth order, her husband's birth order, the availability of substitutes (e.g., whether both parents-in-law are living together), and geographic proximity.

It is also less clear who should provide parental care in an extended family with several adult children. Economic theory suggests that the primary informal caregiver selected is more likely to be the one with the lowest opportunity cost of time, which in turn may be correlated with poorer health and other socioeconomic disadvantages. There is, however, a growing consensus among Koreans that "more able" children, who have higher incomes and spacious houses, should take care of their parents. Despite these changes, parental caregiving by daughters-in-law and daughters remains the major form of elderly long-term care in South Korea.

In this cultural and institutional context, functional limitations of elderly parents and in-laws can be a strong predictor of caregiving by daughters and daughters-in-law, and therefore a good source of instruments. However, parental care needs may have direct influences on adult children's psychological health (Amirkhanyan and Wolf, 2006). Having a functionally dependent parent may adversely affect the health of adult children, which is termed as "the family effect"

independent of "the caregiving effect" (Bobinac *et al.*, 2010). For example, an extra-residential daughter with a demented mother may still suffer psychological consequences even though she is not providing informal care to her mother. In the same vein, Amirkhanyan and Wolf (2003) distinguish non-caregiver stress from caregiver stress. Moreover, parents' functional limitations may also be correlated with adult children's (but not children-in-law's) health outcomes through genetic and behavioral similarities. For example, a 55-year-old daughter of an 85-old-year mother bedridden with stroke may share an increased susceptibility to many chronic diseases. There is less concern with instrument validity in models of physical health effects for the sample of daughters-in-law (where the instruments are functional limitations of parents-in-law, who obviously have no genetic link to the caregiver). The relatively high prevalence of caregiving by daughters-in-law in South Korea allows for studying daughter-in-law caregivers separately from daughter caregivers.

The foregoing discussion provides a conceptual framework for a woman's caregiving decisions. Although the functional limitations of elderly parents and parents-in-law are the primary determinant of this decision, the availability of alternative sources of care, as well as location and the opportunity cost of time, are also likely to influence the decision process.

3. METHODS

We estimate several models to predict a woman's adverse health outcome or health care use (y) as a function of whether she provides informal care to her parents (or parents-in-law) and other personal characteristics (X).

$$y = f(\text{Informal care}, X)$$

The test of the main hypothesis is whether the coefficient on the informal care variable is statistically significant. We hypothesize that the coefficient on informal care will be positive, indicating that women who are informal caregivers have worse health and greater health care use. The goal is to obtain causal estimates of this relationship.

There are two main econometric issues. First, the dependent variables (health and health care use) are multifaceted, and therefore can be measured in several different ways. We estimate two models of health and we estimate both extensive and intensive models of health care use and costs.

Second, informal care may be endogenous in a model that predicts health. Informal care is not provided randomly. The decision to provide informal care will depend in part on unobservable dimensions of health. Whether selection occurs in or out of caregiving, informal caregivers are likely selected, at least partially, based on health status. We need instrumental variables to correct for the potential endogeneity of informal care. These instruments must be correlated with the decision to provide informal care but unrelated directly to the potential caregiver's health. We use indicators of any ADL limitations of the mother-in-law and of the father-in-law as two instruments for the sample of daughters-in-law. Having a parent-in-law with ADL limitations is highly predictive of providing informal care for that parent-in-law.

Counter examples are instructive. Infectious diseases can spread within families, but we are not measuring infectious disease. Some health problems are inherited, but in-laws are not typically closely genetically related. Assortative mating could lead to a correlation in the health and health behaviors of daughters-in-law and parents-in-law; however, this link cannot be as strong as the link between daughters and parents. While some women may develop strong emotional bonds with their in-laws, in general the daughter-in-law will be less emotionally

vested in the health decline of the parent-in-law. In short, we argue that a parent-in-law's health problems primarily affect a daughter-in-law's health through caregiving, rather than directly. The exclusion argument is harder to make for daughters caring for their parents due to heritability and closeness, but for completeness and comparison we also estimate models with daughters as caregivers.

Before running the full IV models, we first obtain Wald estimators for each of our health outcome measures to show the plausibility of the proposed instruments and the rough estimate of the magnitude of the effect. The simple Wald estimator is calculated as follows (Angrist and Evans, 1998). Consider the following regression model.

$$y = \alpha + \beta IC + \varepsilon$$

where y is a health outcome measure, and IC is the endogenous binary variable of interest, whether the respondent provided any informal care. Let α be the constant term and β be the coefficient of interest. The random error is ε . Now we consider a binary instrument (z) for the sample of daughters-in-law — whether either parent-in-law has any ADL limitations ($z=1$ if any ADL limitations, 0 otherwise). The IV-Wald estimate of β is the ratio of two differences. The numerator is the difference between the expected value of y for those whose parents-in-law have ADL limitations ($\overline{y_1}$) and for those whose parents-in-law do not have ADL limitations ($\overline{y_0}$). The denominator is the difference between the expected value of informal care for those whose parents-in-law have ADL limitations ($\overline{IC_1}$) and for those whose parents-in-law do not have ADL limitations ($\overline{IC_0}$). The estimate $\beta_{IV-Wald}$ is the average treatment effect of informal caregiving on a particular health outcome in question (y) for the subgroup of individuals whose probability of providing informal care (IC) has been affected by having a parent-in-law with any ADL limitations (z).

$$\beta_{IV-Wald} = \frac{(\overline{y_1} - \overline{y_0})}{(\overline{IC_1} - \overline{IC_0})}$$

In this sense, $\beta_{IV-Wald}$ can be interpreted as a local average treatment effect (LATE) (Imbens and Angrist, 1994; Angrist *et al.*, 1996; Angrist and Evans, 1998). Robust standard errors for these Wald estimators can be easily obtained by running regression models with the only endogenous explanatory variable of informal care for parent(s)-in-law and one IV of whether either parent-in-law has ADL limitations. In a similar way, we also calculate Wald estimators for the samples of daughters, using a different binary instrument of whether either mother or father has any ADL limitations.

As a next step, we conduct falsification tests of the Wald estimator by using the wrongly matched instrument for each sample; namely using the instrument of whether either mother or father has any ADL limitations to predict the probability of providing informal care for parents-in-law for the sample of daughters-in-law, and using the instrument of whether either the mother-in-law or father-in-law has any ADL limitations to predict the probability of providing informal care for parents for the sample of daughters. The Wald estimators from these falsification tests should not produce sensible results.

We then estimate the full IV models controlling for other observable characteristics. These other characteristics include daughter(-in-law)'s age, education, wealth, home ownership, health insurance, and whether both parents(-in-law) are living together. In this step, we use two instruments for each sample, allowing for overidentification: whether the mother-in-law has any ADL limitations (z_1) and whether the father-in-law has any ADL limitations (z_2) for the sample of daughters-in-law, and whether the mother has any ADL limitations (z_1) and whether the father has any ADL limitations (z_2) for the sample of daughters. We use specification tests to assess the instrument validity and the exogeneity of the potentially endogenous variable of informal care.

Another potential method of dealing with endogeneity is to use longitudinal data and exploit within-individual variations while also possibly controlling for baseline health status (Coe and Van Houtven, 2009). Our data from South Korea have only three waves and thus have limited variation between waves in key variables. We estimated random- and fixed-effects models as well as models including lagged health variables, but found most estimation methods show signs of insufficient variation and limited sample size, particularly for the models with a binary dependent variable. Given the data limitations, our preferred method is the IV model using repeated observations.

As a sensitivity test, we repeat our analysis using logged informal care hours instead of the indicator variable of informal caregiving, and examine whether accounting for the extent of caregiving changes our main results. Because the results from the alternative specification remain largely similar, we present our results using the specification of the indicator variable of informal caregiving. Other results are available upon request.

4. DATA

This study uses data from the first three waves (2006, 2008, and 2010) of the Korean Longitudinal Study of Aging (KLoSA). The KLoSA is a nationally representative study of non-institutionalized South Korean adults aged 45 or older in 15 large administrative areas. The KLoSA is designed to be comparable to aging panel studies from other regions, including the US Health and Retirement Study and the European Survey of Health and Retirement in Europe (Boo and Chang, 2006). The KLoSA contains detailed information on the respondents as well as their children, siblings, and parents. In its first wave, conducted in 2006, 10,254 individuals (5,788 women) in 6,171 households were interviewed face-to-face using the computer-assisted personal

interviewing method. In the second (2008) and third (2010) waves, 84.7% and 77.2% of the original sample were respectively followed up without replacement.

We restrict our study sample to women with any living parents-in-law and women with any living parents, leaving 2,531 observations in the daughter-in-law sample and 4,110 observations in the daughter sample. Because information on parents-in-law can only be identified among currently married women in our data, the daughter-in-law sample is smaller in number than the daughter sample. There are 1,348 observations with both “any living parents-in-law” and “any living parents.” After removing five observations with missing values for the study variables, our main statistical analysis uses the final daughter-in-law ($N=2,528$) and daughter ($N=4,108$) samples.

4.1. Outcome Variables

Various outcome measures have been used to capture the effects of informal caregiving on psychological and physical health as well as on health care use and medication (Schulz, 1990; Haley *et al.*, 1987; Pang, 2000; Yong and McCallion, 2003). In this study, we focus on physical health outcomes, and health care use and costs, rather than psychological health outcomes, because of the concern that psychological outcomes are likely prone to the family effect (Bobinac *et al.*, 2010).

First, we define a dichotomous variable indicating *pain affecting daily activities*. Caregiving often involves physical efforts and may produce physical pain. Moreover, individuals with psychological distress often present with physical symptoms, which is known as somatization (Pang, 2000). While this argument for somatization may raise the concern that pain can result from the family effect, we believe this concern is lessened among daughter-in-law

caregivers. Second, we construct a binary variable of *fair to poor self-rated health* to capture overall subjective health. Although additional health variables are available from the data, most of these variables are excluded because their psychological or disease-specific nature fails to meet two criteria: being able to capture physical health and being sufficiently generic.

Our health care use and costs variables are derived from information on outpatient care use and regular prescription drug use in the past 12 months. We define a binary indicator variable of *any outpatient care use* = 1 if the respondent reported having visited a doctor's office and hospital outpatient office, including the emergency room, at least once in the past 12 months. For those with any outpatient care use, we create a continuous variable of the total *out-of-pocket spending for outpatient care* paid by the respondent during the same period. We also create two variables for regular prescription drug use in the same way as for outpatient care use: *any regular prescription drug use* and *out-of-pocket spending for regular prescription drug use* conditional on any use during the past 12 months. We focus on the use of outpatient care and prescription drugs, rather than inpatient care use and other types of health care use that may be limited in capturing the patient's overall health status. Although our measures of health care use and costs may reflect influences other than health status itself (e.g., insurance status and health seeking behavior) and possibly pose concerns over the accuracy of reporting, we choose to use these measures because they are relatively objective and they allow for summarizing multiple sources of the negative health effects of caregiving. In presenting the potential effect of caregiving on health care financing, the use of these measures becomes highly policy relevant. Because the distribution of out-of-pocket spending shows the typical right-skewed pattern, we tested for the most appropriate Box-Cox transformation and use the natural logarithm of these two cost

variables.

4.2. Explanatory Variables

The key explanatory variable is a binary indicator variable of any informal care provided to parents-in-law (daughter-in-law sample) or a binary variable of any informal care provided to parents (daughter sample). In the KLoSA, ADL care was asked regardless of co-residential status, but IADL care was only asked about care provided to non-co-residing parents(-in-law).

Our statistical models control for a set of demographic and socioeconomic factors, including age, education, home ownership, health insurance status, whether both parents(-in-law) are living together, type of residential area, and survey year. In South Korea, every Korean citizen is insured by either National Health Insurance or Medical Aid. Medical Aid, which is a public assistance program for the poor and other specified groups, has two types. Medical Aid Type 1 is for the poorer among the poor, and charges lower out-of-pocket costs than Type 2. In addition to these statutory medical security programs, individuals can purchase supplemental voluntary private health insurance for protection from the financial burden of out-of-pocket medical expenditures.

5. RESULTS

The main instrument is largely balanced across other covariates, providing evidence that the instrument is unlikely to be correlated with anything other than the probability of being a caregiver. Comparisons of the daughters-in-law by having at least one parent-in-law with ADL limitations (Table 1A) show that the women are similar on all measured characteristics except age, home ownership and, of course, caregiver status. Women with at least one parent-in-law

with ADL limitations are slightly older on average (55.3 years versus 53.3 years) and are more likely to own a home; 60.0% of these women provided informal care to their parents-in-law compared with only 0.2% (4 out of 2,433) of the women who had parents-in-law without ADL limitations. These four caregivers are non-residential caregivers who provided IADL care only. Thus, parent-in-law ADL limitations satisfy one criteria for a good instrument, namely that it be highly correlated with the potentially endogenous variable. The Wald estimators based on ADL limitations for at least one parent-in-law show statistically significant adverse health effects ($p < 0.1$) in the outcome measures except for any outpatient care use and the log of out-of-pocket prescription drug costs, conditional on any regular prescription drug use (see Table 1B). The basic results also hold for the estimates that control for other covariates, as will be shown later.

Group comparisons and Wald estimators for the daughter sample show similar patterns (Table 2), although daughters having parent(s) with ADL limitations are also less educated, more likely to be on the Medical Aid program, and are slightly older. Among daughters having parent(s) with any ADL limitations, 26.3% provided informal care to their parent(s), while informal caregiving was reported by only 0.3% (=10/3850) of women whose parent(s) had no ADL limitations (last row of Table 2A). The Wald estimator for the difference in outcomes measures based on parental ADL limitations show significant adverse effects ($p < 0.05$) for all outcome measures except the log of out-of-pocket prescription drug costs, conditional on any regular prescription drug use (Table 2B).

The falsification tests for the Wald estimators in Table 3 (based on the effect of parental ADL limitation on the probability of providing informal care for parents-in-law in the daughter-in-law sample and that of parent-in-law ADL limitation on the probability of providing informal care for parents in the daughter sample) show no statistically significant adverse health outcomes

($p < 0.10$), even suggesting beneficial effects. The estimated differences in outcomes are extremely large for many measures and have large standard errors. The erratic results from using a “nonsensical” instrument for each sample, combined with the Wald estimators shown in Tables 1B and 2B, suggest that the differential probabilities of providing parental care induced by conceptually appealing instruments (i.e., the presence of functionally dependent parents or parents-in-law for the corresponding sample of daughters or daughters-in-law) can be used to identify the estimated health effects of caregiving. Therefore, the main limitation of the Wald estimators in Tables 1B and 2B is the fact that they are based only on the allocation into groups according to ADL status of parents (for the daughter sample) and parents-in-law (for the daughter-in-law sample) and do not adjust for any of the observed characteristics of the women.

The relatively high proportions of women (daughters or daughters-in-law) who respond to the corresponding instruments of ADL status (for parents or parents-in-law) by providing care, and the negligible proportion of women providing care for parents or parents-in-law without ADL limitations suggest that the instruments should have a strong predictive power in the first-stage regression (Angrist *et al.*, 1996). Indeed, the indicators of ADL limitations among parents or parents-in-law are essentially the most important predictors of informal caregiving by daughters or daughters-in-law (Table 4), which is not surprising since those IVs are the reason for providing parental care. The F -test at the bottom of Table 4 indicates the instruments are strong (F -statistics 86 and 37). In contrast, most other factors have small coefficients and fail to reach statistical significance at the 10% level. All else being equal, having both parents(-in-law) living together reduces the probability of a daughter(-in-law) providing care by 0.8 (1.4) percentage points, and these effects are statistically significant despite the small magnitude. The

surprisingly small magnitude is likely reflected in the small proportion (<20%) of daughters(-in-law) with both parents(-in-law) living together in our study sample (Tables 1 and 2).

Table 5 provides the IV estimates of the effect of caregiving on health outcomes and the results from specification tests (for overidentification and exogeneity) for the separate samples of daughters-in-law and daughters. The tests for overidentification (column 8) show support for valid identification in all models. The tests for exogeneity are more equivocal, with statistical evidence of endogeneity in five of the 12 models estimated.

For the daughter-in-law sample, the models show an adverse impact of caregiving on caregiver health for the following outcome measures ($p < 0.10$, the magnitude of effect in parenthesis): having pain affecting daily activities (13 percentage points), having fair or poor self-reported health (16 percentage points), and out-of-pocket outpatient care costs if any use (47 percent), although the specification test does not reject exogeneity of the IV-Probit estimates.

For the daughter sample, the models show an adverse impact of caregiving on caregiver health for the following outcome measures ($p < 0.10$, the magnitude of effect in parenthesis): having pain affecting daily activities (41 percentage points), having fair or poor self-reported health (56 percentage points), having any outpatient care use (30 percentage points), out-of-pocket outpatient care costs if any use (75 percent) and having any regular prescription drug use (34 percentage points).

No beneficial and statistically significant impacts on health outcomes from caregiving were estimated for any model, as expected. Therefore, although controlling for additional covariates in a full IV framework reduced the number of health measures with adverse impacts from caregiving, adverse impacts from caregiving appears to have occurred for a number of measures. We find that most of the coefficient estimates on the alternative specification of

logged informal care hours do not change qualitatively from the coefficient estimates on the indicator variable of caregiving (Table 6), suggesting the robustness of our results.

6. DISCUSSION

Our results suggest that informal care has adverse health effects along multiple dimensions for daughter-in-law caregivers in South Korea. Our study has carefully tested for the possibility that selection into caregivers by health status might affect the estimation results of caregiver health effects. In doing so, we show IV estimation can help control for endogeneity, starting with more intuitive Wald estimators.

The main methodological contribution of this paper is that our IV approach solves the problem of endogeneity and allows us to estimate causal parameters. The LATE interpretation is directly meaningful because the instruments are directly related to why most informal care is provided. Informal care is more likely provided when an elderly family member has more ADL limitations. Finding appropriate instruments has been a problem in this literature, making most of the published results inconclusive about the causal effects of informal caregiving on caregivers' health.

This study has limitations. First, because of the multiple eligibility criteria required for our IV estimation (e.g., currently married women aged 45 and over with any living parent-in-law) combined with the strict definition of caregiving in the data (i.e., not including co-residential IADL caregiving), our daughter-in-law sample has a modest size with a small number of caregivers, even after three waves of data are combined. One related concern is that our estimates may have been influenced by possible outliers when outcome variables are continuous. Further examining our data, however, excludes this possibility because extreme outliers for the

two continuous outcome variables were observed in the comparison group (i.e., non-caregiver) rather than the treatment group (i.e., caregiver). Second, we focus on obtaining valid population-level estimates of the health effects of informal care, and our models do not account for specific caregiving contexts. Our data do not allow for examining contextual factors: whether the care recipient showed problem behaviors often seen in patients with Alzheimer's dementia; how important the caregiver's choice was in the caregiving decision; what the quality of the dyadic relationship between caregiver and care recipient was; what the main type of caregiving tasks was; and whether the caregiver had role conflict. These limitations, of course, apply to all studies that use large national data sets. Although many of these factors may moderate the health effects of caregiving, several subgroup analyses attempted were limited by small sample sizes even when necessary variables were available. Third, our data do not allow for examining potentially differential effects by phase and duration of caregiving. On the one hand, caregiver health effects can be more pronounced in the earlier phases than later phases. New caregivers may experience numerous changes in their lives and suffer adverse health effects. On the other hand, it is also possible that caregivers who have provided care for an extended period may have poorer health outcomes than new caregivers. The accumulation of longitudinal data would help in accounting for this heterogeneity in future research.

Despite these limitations, we found statistically significant effects of providing informal care on the health of caregivers. In addition to effects on specific measures of health, we found that health care costs of caregivers increased. Thus, given the public financing of health care in South Korea, there are potential tradeoffs between encouraging informal care over long-term care for elders and expending more on the caregiver's health care.

Our study has two policy implications. First, the results support the perspective that informal caregiving is an emerging public health issue (Talley and Crews, 2007). Our results suggest that caregivers may suffer multiple negative health effects across many domains of physical health. Given the demographic trends in South Korea and a strong tradition of informal care, the number of female informal caregivers will likely grow sharply.

Second, our results on increased health care costs among informal caregivers suggest that informal caregiving also has direct economic costs that partially offset any savings from lower formal health care costs for elderly. Estimating the costs of informal care has traditionally focused on the opportunity costs of caregivers' time (Arno *et al.*, 1999; Van Houtven and Norton, 2008). Our results, however, suggest that cost-effectiveness analyses related to long-term care policies also need to take into account the health care costs associated with informal caregiving.

7. ACKNOWLEDGEMENTS

The data for this study were made available by the Korea Labor Institute and Korea Employment Information Service. We thank David Grabowski (Harvard Medical School) and other participants at the International Health Economics Association's 8th World Congress on Health Economics and seminar participants at the University of Rochester and Johns Hopkins University for helpful comments. C.H. Van Houtven was supported by the Veterans Affairs Health Services Research and Development Merit Review Program (MRP 05-311).

8. REFERENCES

- Amirkhanyan AA, Wolf DA. 2003. Caregiver stress and noncaregiver stress: exploring the pathways of psychiatric morbidity. *The Gerontologist* **43**: 817–827.
- Amirkhanyan AA, Wolf DA. 2006. Parent care and the stress process: findings from panel data. *Journals of Gerontology B: Psychological Science and Social Sciences* **61**: S248–S255.
- Angrist JD, Evans WN. 1998. Children and their parents' labor supply: evidence from exogenous variation in family size. *American Economic Review* **88**: 450–477.
- Angrist JD, Imbens GW, Rubin DB. 1996. Identification of causal effects using instrumental variables. *Journal of the American Statistical Association* **91**: 444–455.
- Arno PS, Levine C, Memmott MM. 1999. The economic value of informal caregiving. *Health Affairs (Project Hope)* **18**: 182–188.
- Barer BM, Johnson CL. 1990. A critique of the caregiving literature. *The Gerontologist* **30**: 26–29.
- Bobinac A, Van Exel NJA, Rutten FFH, Brouwer WBF. 2010. Caring for and caring about disentangling the caregiving effect and the family effect. *Journal of Health Economics* **29**: 549–556.
- Boo K, Chang J. 2006. Korean Longitudinal Study of Ageing: research design for international comparative studies. *Survey Research* **7**: 97–122.
- Bookwala J, Yee JL, Schulz R. 2000. Caregiving and Detrimental Mental and Physical Health Outcomes. In: Williamson GM, Shaffer DR, Parmelee PA (Eds). *Physical Illness and Depression in Older Adults; A Handbook of Theory, Research, and Practice*. Kluwer Academic/Plenum Publishers: New York. 93–131.
- Chee YK. 2000. Elder care in Korea: the future is now. *Aging International* **26**: 25–37.

- Choi H. 1993. Cultural and noncultural factors as determinants of caregiver burden for the impaired elderly in South Korea. *The Gerontologist* **33**: 8–15.
- Coe NB, Van Houtven CH. 2009. Caring for mom and neglecting yourself? The health effects of caring for an elderly parent. *Health Economics* **18**: 991–1010.
- Dilworth-Anderson P, Williams IC, Gibson BE. 2002. Issues of race, ethnicity, and culture in caregiving research: a 20-year review (1980–2000). *The Gerontologist* **42**: 237–272.
- Haley WE, Levine EG, Brown SL, Bartolucci AA. 1987. Stress, appraisal, coping, and social support as predictors of adaptational outcome among dementia caregivers. *Psychology and Aging* **2**: 323–330.
- Hsu, HC, Shyu YI. 2003. Implicit exchanges in family caregiving for frail elders in Taiwan. *Qualitative Health Research* **13**: 1078–1093.
- Imbens GW, Angrist JD. 1994. Identification and estimation of local average treatment effects. *Econometrica* **62**: 467–475.
- Pang KYC. 2000. Symptom expression and somatization among elderly Korean immigrants. *Journal of Clinical Geropsychology* **6**: 199–212.
- Pinquart M, Sörensen S. 2003. Differences between caregivers and noncaregivers in psychological health and physical health: a meta-analysis. *Psychology and Aging* **18**: 250–267.
- Pinquart M, Sörensen S. 2006. Gender differences in caregiver stressors, social resources, and health: an updated meta-analysis. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences* **61**: 33–45.

- Pinquart M, Sörensen S. 2007. Correlates of physical health of informal caregivers: a meta-analysis. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences* **62**: 126–137.
- Schulz R. 1990. Theoretical perspectives on caregiving: concepts, variables, and methods. In: Biegel DE, Blum A. (Eds). *Aging and Caregiving: Theory, Research and Policy*. Sage Publications: New York. 27–52.
- Schulz R, Visintainer P, Williamson GM. 1990. Psychiatric and physical morbidity effects of caregiving. *Journal of Gerontology* **45**: 181–191.
- Schulz R, O'Brien AT, Bookwala J, Fleissner K. 1995. Psychiatric and physical morbidity effects of dementia caregiving: prevalence, correlates, and causes. *The Gerontologist* **35**: 771–791.
- Sung KT. 1990. A new look at filial piety: ideals and practices of family-centered parent care in Korea. *The Gerontologist* **30**: 610–617.
- Talley RC, Crews JE. 2007. Framing the public health of caregiving. *American Journal of Public Health* **97**: 224–228.
- Van Houtven CH, Norton EC. 2008. Informal care and Medicare expenditures: testing for heterogeneous treatment effects. *Journal of Health Economics* **27**: 134–156.
- Vitaliano PP, Zhang J, Scanlan JM. 2003. Is caregiving hazardous to one's physical health? A meta-analysis. *Psychological Bulletin* **129**: 946–972.
- Walker AJ, Pratt CC, Eddy L. 1995. Informal caregiving to aging family members: a critical review. *Family Relations* **44**: 402–411.
- Yamamoto N, Wallhagen MI. 1997. The continuation of family caregiving in Japan. *Journal of Health and Social Behavior* **38**: 164–176.

- Yee JL, Schulz R. 2000. Gender differences in psychiatric morbidity among family caregivers: a review and analysis. *The Gerontologist* **40**: 147–164.
- Yong F, McCallion P. 2003. Hwabyung as caregiving stress among Korean-American caregivers of a relative with dementia. *Journal of Gerontological Social Work* **42**: 3–19.

Table 1. Summary statistics and Wald estimators in daughters-in-law ($N=2528$)

Variable	Parent(s)-in-law have no ADL limitations (n=2433)	Parent(s)-in-law have any ADL limitations (n=95)		
Table 1A: Daughter-in-law characteristics				
	Mean	Mean	<i>p</i> -value†	
Age (year)	53.29	55.31	<0.01	
Education (%)			0.21	
Elementary school or less	24.58	27.37		
Middle school	20.72	25.26		
High school	44.43	43.16		
College or more	10.28	4.21		
Home ownership (%)	83.26	92.63	0.02	
Statutory medical security (%)			0.52	
National Health Insurance	98.31	96.84		
<i>Medical Aid</i> Type 1	0.70	1.05		
<i>Medical Aid</i> Type 2	0.99	2.11		
Voluntary private health insurance	56.39	52.63	0.47	
Both parents-in-law living together (%)	16.19	10.53	0.14	
Residential area (%)			0.40	
Metropolitan	46.65	42.11		
Small city	34.57	33.68		
Rural	18.78	24.21		
KLoSA wave (%)			0.68	
Wave 1 (2006)	41.10	41.05		
Wave 2 (2008)	32.22	35.79		
Wave 3 (2010)	26.67	23.16		
Caregiving for a least one parent-in-law	0.0016	0.6000	-	
Table 1B: Wald estimator††				
	Mean	Mean	Wald	(S.E.)†††
Pain affecting daily activities	0.14	0.26	0.20***	(0.08)
Self-reported health <i>fair</i> to <i>poor</i>	0.36	0.49	0.22**	(0.09)
Any outpatient care use	0.62	0.60	-0.03	(0.09)
Log(Out-of-pocket costs) if any use	1.81	2.13	0.55**	(0.28)
Any regular prescription drug use	0.29	0.39	0.17*	(0.09)
Log(Out-of-pocket costs) if any use	2.96	3.14	0.25	(0.30)

Notes:

ADL, activity of daily living

† Test statistics for *t*-test (continuous variable) or chi-sq test (categorical variable).

†† Table 5A provides the number of observations used for each outcome measure.

††† Robust standard errors in parentheses

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$.

Table 2. Summary statistics and Wald estimators among daughters (N=4108)

Variable	Parent(s) have no ADL limitations (n=3850)	Parent(s) have any ADL limitations (n=258)		
Table 2A: Daughter characteristics				
	Mean	Mean	<i>p</i> -value†	
Age (year)	54.16	56.56	<0.01	
Education (%)			<0.01	
Elementary school or less	27.58	41.86		
Middle school	21.74	19.77		
High school	40.52	31.01		
College or more	10.16	7.36		
Home ownership (%)	81.17	78.29	0.26	
Statutory medical security (%)			0.03	
National Health Insurance	96.23	93.41		
<i>Medical Aid</i> Type 1	1.64	3.88		
<i>Medical Aid</i> Type 2	2.13	2.71		
Voluntary private health insurance	53.64	49.61	0.21	
Both parents living together (%)	6.49	3.49	0.06	
Residential area (%)			0.21	
Metropolitan	47.51	53.10		
Small city	34.39	30.23		
Rural	18.1	16.67		
KLoSA wave (%)			0.11	
Wave 1 (2006)	40.34	44.96		
Wave 2 (2008)	32.05	33.33		
Wave 3 (2010)	27.61	21.71		
Caregiving for a least one parent	0.0026	0.2636	-	
Table 2B: Wald estimator††				
	Mean	Mean	Wald	(S.E.)†††
Pain affecting daily activities	0.17	0.37	0.75***	(0.16)
Self-reported health <i>fair</i> to <i>poor</i>	0.39	0.61	0.85***	(0.16)
Any outpatient care use	0.63	0.71	0.33***	(0.13)
Log(Out-of-pocket costs) if any use	1.85	2.08	0.92**	(0.39)
Any regular prescription drug use	0.31	0.47	0.63***	(0.15)
Log(Out-of-pocket costs) if any use	2.88	2.98	0.43	(0.55)

Notes:

ADL, activity of daily living

† Test statistics for *t*-test (continuous variable) or chi-sq test (categorical variable).

†† Table 5B provides the number of observations used for each outcome measure.

††† Robust standard errors in parentheses

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$.

Table 3. Falsification test for Wald estimation in daughters-in-law and daughters

Sample	Daughters-in-law		Daughters	
Potentially endogenous variable	Caregiving for parent-in-law		Caregiving for parent	
Instrumental variable	Parent has ADL limitations		Parent-in-law has ADL limitations	
Outcome	Coeff.	(S.E.)†	Coeff.	(S.E.)†
Prob(Pain affecting daily activities)	4.56	(5.88)	-11.23*	(6.70)
Prob(Self-reported health <i>fair to poor</i>)	8.19	(11.15)	-15.93*	(8.32)
Prob(Any outpatient care use)	5.54	(8.09)	-3.87	(6.30)
Log(Out-of-pocket costs) if any use	10.38	(12.58)	-40.84	(29.56)
Prob(Any regular prescription drug use)	11.16	(15.11)	-14.55*	(8.32)
Log(Out-of-pocket costs) if any use	8.76	(11.94)	-6.98	(18.45)

Notes:

ADL, activity of daily living

† Robust standard errors in parentheses

* $p < 0.1$.

Table 4. First-stage regression of caregiving for parent-in-law and for parent

Variable	Caregiving for parent-in-law by daughter-in-law		Caregiving for parent by daughter	
	Coefficients	(S.E.)†	Coefficients	(S.E.)†
Instrumental variables				
Father-in-law has ADL limitations	0.3940***	(0.1067)	-	
Mother-in-law has ADL limitations	0.6178***	(0.0554)	-	
Father has ADL limitations	-		0.2206***	(0.0644)
Mother has ADL limitations	-		0.2564***	(0.0327)
Age (year)	0.0007	(0.0004)	-0.0003	(0.0003)
Education				
Elementary school or less	-0.0069	(0.0059)	0.0016	(0.0067)
Middle school	-0.0056	(0.0054)	0.0086	(0.0063)
High school	-0.0032	(0.0043)	0.0060	(0.0056)
College or more (ref.)	-		-	
Home ownership	-0.0023	(0.0064)	0.0001	(0.0057)
Statutory medical security				
National Health Insurance (ref.)	-		-	
Medical Aid Type 1	-0.0055	(0.0076)	0.0336	(0.0301)
Medical Aid Type 2	0.0385	(0.0238)	0.0014	(0.0137)
Voluntary private health insurance	-0.0081**	(0.0041)	0.0065	(0.0040)
Both parents(-in-law) living together	-0.0141**	(0.0056)	-0.0078*	(0.0041)
Residential area				
Metropolitan	-		-	
Small city	-0.0059	(0.0046)	-0.0145***	(0.0041)
Rural	-0.0073	(0.0064)	-0.0103*	(0.0058)
Survey wave				
Wave 1 (ref.)	-		-	
Wave 2	-0.0027	(0.0040)	-0.0033	(0.0043)
Wave 3	-0.0008	(0.0059)	-0.0025	(0.0046)
Constant	-0.0188	(0.0247)	0.0184	(0.0182)
Observations	2528		4108	
R-squared	0.5841		0.2151	
IV strength (<i>F</i> -statistic)	86.29***		36.73***	

Notes:

ADL, activity of daily living

† Robust standard errors in parentheses

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$.

Table 5: IV Estimates and specification tests for the effect of caregiving

Outcome	Obs.	OLS or Probit		IV-2SLS or IV-Probit		Marginal effects of IV-Probit		IV specification tests	
		(1)	(2) Coeff.	(3) (S.E.) [†]	(4) Coeff.	(5) (S.E.) [†]	(6) %points	(7) (S.E.) [†]	(8) Overid
Table 5A: Daughter-in-law sample									
Prob(Pain affecting daily activities)	2528	0.41**	(0.18)	0.62***	(0.23)	13.1***	(4.9)	NS	NS
Prob(Self-reported health <i>fair to poor</i>)	2528	0.23	(0.17)	0.46**	(0.23)	15.7**	(7.9)	NS	NS
Prob(Any outpatient care use)	2525	-0.09	(0.18)	-0.18	(0.23)	-6.6	(8.6)	NS	NS
Log(Out-of-pocket costs) if any use	1548	0.19	(0.21)	0.47*	(0.27)	-		NS	**
Prob(Any regular prescription drug use)	2528	0.21	(0.17)	0.27	(0.23)	8.2	(7.2)	NS	NS
Log(Out-of-pocket costs) if any use	721	0.29	(0.29)	0.30	(0.31)	-		NS	NS
Table 5B: Daughter sample									
Prob(Pain affecting daily activities)	4108	0.31*	(0.17)	1.73***	(0.40)	41.3***	(9.4)	NS	***
Prob(Self-reported health <i>fair to poor</i>)	4108	0.22	(0.15)	1.62***	(0.37)	56.1***	(12.5)	NS	***
Prob(Any outpatient care use)	4105	0.10	(0.17)	0.82**	(0.38)	29.8**	(13.9)	NS	**
Log(Out-of-pocket costs) if any use	2567	0.46**	(0.21)	0.75*	(0.39)	-		NS	NS
Prob(Any regular prescription drug use)	4108	0.22	(0.18)	1.07***	(0.39)	34.0***	(12.2)	NS	***
Log(Out-of-pocket costs) if any use	1268	0.41	(0.31)	0.57	(0.54)	-		NS	NS

Notes: NS denotes 'not significant' at the 10% level. *F*-statistics for first-stage regressions for the four conditional models were over 50 in the daughter-in-law sample, and 22 and 15 in the daughter sample. All models control for other explanatory variables presented in Table 4, namely age, education, home ownership, health insurance status, and whether both parents(-in-law) are living together, type of residential area, and survey wave.

[†] Robust standard errors in parentheses.

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$.

Table 6: IV Estimates and Specification Tests for the Effect of Caregiving Hours

Outcome	Obs.	OLS or Probit		IV-2SLS or IV-Probit		IV specification tests	
	(1)	(2) Coeff.	(3) (S.E.)†	(4) Coeff.	(5) (S.E.)†	(6) Overid	(7) Exo.
<i>Daughter-in-law sample</i>							
Prob(Pain affecting daily activities)	2527	0.18**	(0.08)	0.33**	(0.13)	NS	NS
Prob(Self-reported health <i>fair</i> to <i>poor</i>)	2527	0.08	(0.08)	0.25**	(0.13)	NS	*
Prob(Any outpatient care use)	2524	0.03	(0.09)	-0.10	(0.12)	NS	NS
Log(Out-of-pocket costs) if any use	1548	0.10	(0.10)	0.24*	(0.14)	NS	NS
Prob(Any regular prescription drug use)	2527	0.14*	(0.08)	0.16	(0.12)	NS	NS
Log(Out-of-pocket costs) if any use	720	0.13	(0.10)	0.15	(0.15)	NS	NS

Notes: NS denotes 'not significant' at the 10% level. Logarithm of weekly care hours was used. F -statistics for first-stage regressions were 26 in the four IV-Probit models and 23 and 12 in the two conditional IV-2SLS models. The same set of control variables were used as in Table 5. One observation was dropped because of missing value for caregiving hours.

† Robust standard errors in parentheses.

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$.