

NBER WORKING PAPER SERIES

THE IMPACT OF PHARMACEUTICAL INNOVATION ON PREMATURE CANCER
MORTALITY IN CANADA, 2000-2011

Frank R. Lichtenberg

Working Paper 21239

<http://www.nber.org/papers/w21239>

NATIONAL BUREAU OF ECONOMIC RESEARCH

1050 Massachusetts Avenue

Cambridge, MA 02138

June 2015

This research was supported by Canada's ResearchBased Pharmaceutical Companies (Rx&D), the national association that represents 50 researchbased pharmaceutical companies in Canada. The sponsor placed no restrictions or limitations on data, methods, or conclusions and had no right of review or control over the outcome of the research. The views expressed herein are those of the author and do not necessarily reflect the views of the National Bureau of Economic Research.

NBER working papers are circulated for discussion and comment purposes. They have not been peer-reviewed or been subject to the review by the NBER Board of Directors that accompanies official NBER publications.

© 2015 by Frank R. Lichtenberg. All rights reserved. Short sections of text, not to exceed two paragraphs, may be quoted without explicit permission provided that full credit, including © notice, is given to the source.

The Impact of Pharmaceutical Innovation on Premature Cancer Mortality in Canada, 2000-2011
Frank R. Lichtenberg
NBER Working Paper No. 21239
June 2015
JEL No. C23,C33,I10,J11,J17,L65,O33

ABSTRACT

The premature cancer mortality rate has been declining in Canada, but there has been considerable variation in the rate of decline across cancer sites. I analyze the effect that pharmaceutical innovation had on premature cancer mortality in Canada during the period 2000-2011, by investigating whether the cancer sites that experienced more pharmaceutical innovation had larger declines in the premature mortality rate, controlling for changes in the incidence rate.

The estimates imply that pharmaceutical innovation during the period 1985-1996 reduced the number of years of potential life lost to cancer before age 75 in 2011 by 105,366. The cost per life-year before age 75 gained from previous pharmaceutical innovation is estimated to have been 2730 USD. The evidence suggests that, even if these drugs had been sold at branded rather than generic prices, the cost per life-year gained would have been below 11,000 USD, a figure well below even the lowest estimates of the value of a life-year gained.

Frank R. Lichtenberg
Columbia University
504 Uris Hall
3022 Broadway
New York, NY 10027
and NBER
frl1@columbia.edu

I. Introduction

Previous authors have argued that “reducing premature mortality is a crucial public health objective” (Renard, Tafforeau, and Deboosere (2014)). A widely used measure of premature mortality is years of potential life lost (YPLL) before a given age (e.g. age 75), i.e. the number of years *not* lived by an individual who died before that age (Association of Public Health Epidemiologists in Ontario (2015)). Statistics of YPLL are published by the World Health Organization, the OECD, and government agencies of Canada, the U.S., and other countries. Burnet et al (2005) argue that YPLL “should be considered when allocating research funds.” In the U.S., “cancer [was] responsible for more [YPLL] than all other causes of death combined” in 2008 (National Cancer Institute (2015c)). In Canada, premature (before age 75) mortality from cancer is about twice as great as premature mortality from circulatory diseases.

But as shown in Figure 1, the premature cancer mortality rate has been declining; it declined about 20% between 1996 and 2006. The cancer incidence rate remained approximately constant during that period.

While the premature mortality rate from all cancers combined has been declining in Canada, Figure 2 indicates that there has been considerable variation in the rate of decline across cancer sites. During the period 2000-2011, the premature mortality rate from breast cancer declined 20%, and from cancer of lymphoid, haematopoietic and related tissue declined 27%, but the premature mortality rate from lip, oral cavity, and pharynx cancer increased 6%, and from cancer of female genital organs increased 8%. I will show that this variation in the rate of decline of premature mortality cannot be explained by variation in the rate of decline of incidence.

In this paper, I will analyze the effect that pharmaceutical innovation has had on premature cancer mortality in Canada during the period 2000-2011.^{1,2} The analysis will be

¹ Lichtenberg (2014b) analyzed the impact of pharmaceutical innovation and other types of medical innovation on cancer mortality in the U.S. during the period 2000–2009. But as the Commonwealth Fund (2011) demonstrated, the U.S. health care system differs dramatically from the health care systems of other OECD countries, including Canada. For example, in 2008 per capita spending on health was 85% higher in the U.S. than it was in Canada. Also, the outcome measure and the measure of pharmaceutical innovation used in the present study will differ from those used in Lichtenberg (2014b).

² I use the sample period 2000-2011 to avoid potential discontinuities in the mortality data, because Canada used the ICD9 cause-of-death classification during 1979-1999, and the ICD10 cause-of-death classification during 2000-2011.

performed using a difference-in-differences research design based on longitudinal disease-level data. In essence, I will investigate whether the cancer sites that experienced more pharmaceutical innovation had larger declines in the premature mortality rate, controlling for changes in the incidence rate. Figure 3 illustrates that the rate of pharmaceutical innovation, as measured by the number of drugs registered during the period 1988-2013, varied considerably across cancer sites. Only 5 drugs for cancer of the eye, brain and central nervous system were registered, while 19 drugs for cancer of digestive organs were registered.

The analysis will be based on aggregate data—longitudinal data on 15 cancer sites³—rather than patient-level data. Stukel et al (2007) argue that comparisons of outcomes between patients treated and untreated in observational studies may be biased due to differences in patient prognosis between groups, often because of unobserved treatment selection biases. I believe that difference-in-differences estimates based on aggregate panel data are much less likely to be subject to unobserved treatment selection biases than estimates based on cross-sectional patient-level data.⁴ Moreover, the outcome measures that we analyze (premature mortality rates) are not subject to lead-time bias.⁵

In Section II, I describe an econometric model of premature cancer mortality. The data sources used to construct the data to estimate this model are described in Section III. Empirical results are presented in Section IV. Key implications of the estimates are discussed in Section V. Section VI provides a summary and conclusions.

II. Premature cancer mortality model

In his model of endogenous technological change, Romer (1990) hypothesized an aggregate production function such that an economy's output depends on the "stock of ideas"

³ The 15 cancer sites are the 15 malignant neoplasm ICD-10 blocks defined by the World Health Organization.

⁴ Jalan and Ravallion (2001) argued that "aggregation to village level may well reduce measurement error or household-specific selection bias" (p. 10).

⁵ Survival time for cancer patients is usually measured from the day the cancer is diagnosed until the day they die. Patients are often diagnosed after they have signs and symptoms of cancer. If a screening test leads to a diagnosis before a patient has any symptoms, the patient's survival time is increased because the date of diagnosis is earlier. This increase in survival time makes it seem as though screened patients are living longer when that may not be happening. This is called lead-time bias. It could be that the only reason the survival time appears to be longer is that the date of diagnosis is earlier for the screened patients. But the screened patients may die at the same time they would have without the screening test. See National Cancer Institute (2015a).

that have previously been developed, as well as on the economy's endowments of labor and capital. The premature mortality model that I will estimate may be considered a health production function, in which premature mortality is an inverse indicator of health output or outcomes, and the cumulative number of drugs approved is analogous to the stock of ideas. The first model will be of the following form:

$$\ln(YPLL75_{st}) = \beta_k \text{CUM_NCE}_{s,t-k} + \gamma \ln(\text{INC_RATE75}_{st}) + \alpha_s + \delta_t + \varepsilon_{st} \quad (1)$$

where

$YPLL75_{st}$ = years of potential life lost before age 75 from cancer at site s per 100,000 population age 0-74 in year t ($t = 2000, \dots, 2011$)

$\text{CUM_NCE}_{i,t-k} = \sum_d \text{IND}_{ds} \text{REGISTERED}_{d,t-k}$ = the number of new chemical entities (drugs) to treat cancer at site s that had been registered in Canada by the end of year $t-k$

IND_{ds} = 1 if drug d is used to treat (indicated for) cancer at site s
 = 0 if drug d is not used to treat (indicated for) cancer at site s

$\text{REGISTERED}_{d,t-k}$ = 1 if drug d was registered in Canada by the end of year $t-k$
 = 0 if drug d was not registered in Canada by the end of year $t-k$

INC_RATE75_{st} = the average annual incidence rate of cancer at site s per 100,000 population age 0-74 in years $t-5, t-4, \dots, t-1$ ⁶

α_i = a fixed effect for cancer at site s

δ_t = a fixed effect for year t

Inclusion of year and cancer-site fixed effects controls for the overall decline in premature cancer mortality and for stable between-disease differences in premature mortality. A negative and significant estimate of β_k in eq. (1) would signify that diseases for which there was more pharmaceutical innovation had larger declines in premature mortality. The functional form of eq. (1) has the property of diminishing marginal productivity: the absolute reduction in premature mortality declines with each successive increase in the number of drugs.

As illustrated by Figure 4, the data exhibit heteroskedasticity—diseases with larger mean premature mortality rates had smaller (positive and negative) annual percentage fluctuations in

⁶ The most recent available incidence data are for the year 2010.

YPLL75. Eq. (1) will therefore be estimated by weighted least-squares, weighting by the mean premature mortality rate during 2000-2011 ($(\sum_t \text{YPLL75}_{it}) / 12$). The standard errors of eq. (1) will be clustered within cancer sites.

Although one would expect an increase in true cancer incidence to increase premature cancer mortality, cancer incidence rates are subject to measurement error, so one should not necessarily expect the coefficient on measured cancer incidence (γ) to be positive. Let I and I^* represent measured and true cancer incidence, respectively. Then $I = (I / I^*) \times I^*$, and $\log(I) = \log(I / I^*) + \log(I^*)$. Measured cancer incidence can increase for two reasons: an increase in true cancer incidence, or an increase in the ratio of measured incidence to true incidence. The latter could occur as a result of increasing quantity or quality of cancer screening. More and better cancer screening could lead to earlier diagnosis, which might reduce premature mortality. Therefore the effect on premature mortality of increases in I^* and increases in (I / I^*) may offset one another: the former is likely to increase premature mortality, but the latter may reduce it. For this reason, although controlling (in an unrestrictive manner) for measured incidence in the premature mortality model seems appropriate, we should not be surprised if we don't find a significant effect of measured incidence on premature mortality.

Estimation of eq. (1) enables determination of how much of the decline in Canadian premature mortality during the sample period (2000-2011) can be attributed to the introduction of new drugs. The expression $(\delta_{2011} - \delta_{2000})$ indicates the 2000-2011 decline in premature mortality, controlling for (holding constant) the number of drugs and cancer incidence, i.e., in the absence of pharmaceutical innovation. Suppose eq. (1) is estimated, excluding $\text{CUM_NCE}_{i,t-k}$, and that the year fixed effects from that equation are denoted by δ'_t . Then $(\delta'_{2011} - \delta'_{2000})$ indicates the 2000-2011 decline in Canadian premature mortality, not holding constant the number of drugs, i.e., in the presence of pharmaceutical innovation, and $(\delta'_{2011} - \delta'_{2000}) - (\delta_{2011} - \delta_{2000})$ is an estimate of the 2000-2011 decline in premature mortality attributable to pharmaceutical innovation.

The measure of pharmaceutical innovation in eq. (1)—the number of chemical substances previously commercialized to treat a disease—is not the theoretically ideal measure. Premature mortality is presumably more strongly related to the drugs *actually* used to treat a disease than it is to the drugs that *could be* used to treat the disease. A preferable measure is the mean *vintage* of drugs used to treat a disease, defined as $\text{VINTAGE}_{it} = \sum_d Q_{dit} \text{LAUNCH_YEAR}_d / \sum_d Q_{dit}$,

where Q_{dit} = the quantity of drug d used to treat disease i in year t , and $LAUNCH_YEAR_d$ = the world launch year of drug d .⁷ Unfortunately, measurement of $VINTAGE_{it}$ is infeasible: even though data on the total quantity of each drug in each year ($Q_{d,t} = \sum_i Q_{dit}$) are available, many drugs are used to treat multiple diseases,⁸ and there is no way to determine the quantity of drug d used to treat cancer at site s in year t .⁹ However, it is shown in Appendix 1 of Lichtenberg (2014a) that there is a highly significant positive correlation across *drug classes* between changes in the (quantity-weighted) vintage of drugs and changes in the number of chemical substances previously commercialized within the drug class.

Pharmaceutical innovation is not the only type of medical innovation that is likely to contribute to premature mortality. Other medical innovation, such as innovation in diagnostic imaging, surgical procedures, and medical devices, is also likely to affect premature mortality. Therefore, measures of these other types of medical innovation should be included in the eq. (1). Unfortunately, longitudinal disease-level measures of non-pharmaceutical medical innovation are not available for Canada. But failure to control for non-pharmaceutical medical innovation is unlikely to bias estimates of the effect of pharmaceutical innovation on premature mortality, for two reasons. First, pharmaceuticals are more research-intensive than other types of medical care: in 2007, prescription drugs accounted for 10% of U.S. health expenditure (Center for Medicare and Medicaid Services (2013: Table 2)), but more than half of U.S. funding for biomedical research came from pharmaceutical and biotechnology firms (Dorsey et al (2010)). Much of the rest came from the federal government (i.e. the NIH), and new drugs often build on upstream government research (Sampat and Lichtenberg (2011)). The National Cancer Institute (2015b) says that it “has played an active role in the development of drugs for cancer treatment for 50 years... [and] that approximately one half of the chemotherapeutic drugs currently used by

⁷ According to the Merriam Webster dictionary, one definition of vintage is “a period of origin or manufacture (e.g. a piano of 1845 vintage)”. <http://www.merriam-webster.com/dictionary/vintage>. Robert Solow (1960) introduced the concept of vintage into economic analysis. Solow’s basic idea was that technical progress is “built into” machines and other goods and that this must be taken into account when making empirical measurements of their roles in production. This was one of the contributions to the theory of economic growth that the Royal Swedish Academy of Sciences cited when it awarded Solow the 1987 Alfred Nobel Memorial Prize in Economic Sciences (Nobelprize.org (2015)).

⁸ For example, dactinomycin is used to treat C45-C49 connective and soft tissue neoplasms, C51-C58 female genital organ neoplasms, C60-C63 male genital organ neoplasms, and C64-C68 urinary organ neoplasms.

⁹ Outpatient prescription drug claims usually don’t show the indication of the drug prescribed. Claims for drugs administered by doctors and nurses (e.g. chemotherapy) often show the indication of the drug. In the US, 70% of spending on anticancer drugs is for drugs covered under the medical benefit and infused or injected. However, t data on claims for drugs administered by doctors and nurses are not available for Canada.

oncologists for cancer treatment were discovered and/or developed” at the National Cancer Institute.

Second, previous research based on U.S. data indicates that non-pharmaceutical medical innovation is not positively correlated across diseases with pharmaceutical innovation. In Appendix 2 of Lichtenberg (2014a), it is shown that, in the U.S. during the period 1997-2007, the rate of pharmaceutical innovation was not positively correlated across diseases with the rate of medical procedure innovation and may have been *negatively* correlated with the rate of diagnostic imaging innovation. Also, Lichtenberg (2014b) found that estimates of the effect of pharmaceutical innovation on U.S. cancer mortality rates were insensitive to the inclusion or exclusion of measures of non-pharmaceutical medical innovation. This suggests that failure to control for other medical innovation is unlikely to result in overestimation of the effect of pharmaceutical innovation on longevity growth.

In eq. (1), premature mortality from cancer at site s in year t depends on the number of new chemical entities (drugs) to treat cancer at site s registered in Canada by the end of year $t-k$, i.e. there is a lag of k years. Eq. (1) will be estimated for different values of k : $k = 0, 5, 10, 15, 20, 25$.¹⁰ One would expect there to be a substantial lag because new drugs diffuse gradually—they won’t be used widely until years after registration. Two kinds of evidence—“within molecule” and “between molecule”—support the gradual diffusion hypothesis. The first kind consists of estimates based on the π_y parameters from the following equation:

$$\ln(\text{SU}_{my}) = \rho_m + \pi_y + \varepsilon_{my} \quad (2)$$

where

SU_{my} = the number of standard units¹¹ of molecule m sold in Canada y years after registration ($y = 0, 1, \dots, 11$)

ρ_m = a fixed effect for molecule m

π_y = a fixed effect for age y

¹⁰ A separate model is estimated for each value of k , rather than including multiple values ($\text{CUM_NCE}_{i,t-1}$, $\text{CUM_NCE}_{i,t-2}$, $\text{CUM_NCE}_{i,t-3}, \dots$) in a single model because CUM_NCE is highly serially correlated (by construction), which would result in extremely high multicollinearity if multiple values were included.)

¹¹ The number of standard ‘dose’ units sold is determined by taking the number of counting units sold divided by the standard unit factor which is the smallest common dose of a product form as defined by IMS HEALTH. For example, for oral solid forms the standard unit factor is one tablet or capsule whereas for syrup forms the standard unit factor is one teaspoon (5 ml) and injectable forms it is one ampoule or vial. Other measures of quantity, such as the number of patients using the drug, prescriptions for the drug, or defined daily doses of the drug, are not available.

The expression $\exp(\pi_y - \pi_0)$ is a “relative utilization index”: it is the mean ratio of the number of units of a molecule sold y years after registration to the number of units of the same molecule sold in the year that it was registered. Using annual data on the number of standard units of molecules sold in Canada during the period 1999-2010, I estimated eq. (2). Estimates of the “relative utilization index,” based on data on 25 molecules used to treat cancer that were registered after 1998, are shown in Figure 5. These estimates indicate that the number of units sold 10 years after registration is about ten times as great as the number of units sold one year after registration. Moreover, Figure 5 provides a conservative estimate of the slope of the age-utilization profile, because there was zero utilization of many of these molecules in the first few years after they were registered.¹²

Figure 6 provides “between-molecule” evidence of gradual diffusion; it shows data on the mean number of standard units of cancer drugs sold (in thousands) in Canada in 2010, by period of registration in Canada. Mean utilization in 2010 of drugs registered after 2000 is only 15% as high as mean utilization of drugs registered during 1991-2000, and 17% as high as mean utilization of drugs registered during 1981-1990.

The relatively low utilization of new drugs may be due to several factors. One is that the prices of old drugs (most of which are no longer patent-protected) are considerably lower than the prices of new, patent-protected drugs. A second factor may be that it takes time for physicians to become knowledgeable about new treatment options. A third potential factor is that new drugs may be targeted at smaller patient populations. Data from the U.S. Food and Drug Administration (2015) indicate that drugs approved by the FDA since 2000 were twice as likely to include pharmacogenomic information in their labeling as drugs approved before 2000. A fourth potential factor is that older drugs are more likely to have supplemental indications, i.e. indications approved after the drug was initially launched, than new drugs.¹³

The measure of pharmaceutical innovation, $CUM_NCE_{s,t-k} = \sum_d IND_{ds} REGISTERED_{d,t-k}$, is based on whether drug d had an indication for cancer at site s *at the end of 2011*. One would

¹² Since the dependent variable of eq. (2) is logarithmic, observations for which $SU_{my} = 0$ had to be excluded.

¹³ The measure of pharmaceutical innovation, $CUM_NCE_{s,t-k} = \sum_d IND_{ds} REGISTERED_{d,t-k}$, is based on whether drug d had an indication for cancer at site s *at the end of 2011*. One would prefer to base the measure on whether drug d had an indication for cancer at site s *at the end of year $t-k$* . Data in the U.S. FDA’s Drugs@FDA data files indicate that about one in four new molecular entities has supplemental indications, i.e. indications approved after the drug was initially launched.

prefer to base the measure on whether drug d had an indication for cancer at site s *at the end of year $t-k$* . FDA data indicate that about one in four new molecular entities has supplemental indications, i.e. indications approved after the drug was initially registered.¹⁴

In eq. (1), the measure of premature mortality is the number of years of potential life lost before age 75. This is the age threshold used in Statistics Canada's key socioeconomic database (CANSIM). Other authorities use different age thresholds; the CDC (2013) provides estimates of YPLL before ages 65, 70, 75, 80, and 85. To assess the robustness of my results, I will estimate models similar to eq. (1), using age thresholds 65 and 55 as well as 75.

Chemical substances are divided into different groups according to the organ or system on which they act and their therapeutic, pharmacological and chemical properties. In the Anatomical Therapeutic Chemical (ATC) classification system developed by the World Health Organization Collaborating Centre for Drug Statistics Methodology, drugs are classified in groups at five different levels. The highest (1st) level is the “anatomical main group” level; there are 14 anatomical main groups. The 2nd, 3rd, 4th, and 5th levels are “therapeutic subgroup,” “pharmacological subgroup,” “chemical subgroup,” and “chemical substance,” respectively.¹⁵ Premature mortality from a disease may depend on the number of chemical (or pharmacological) *subgroups* that have previously been developed to treat the disease rather than, or in addition to, the number of chemical *substances* (drugs) that have previously been developed to treat the disease. This will be investigated by estimating versions of eq. (1) in which $CUM_SUBGROUP_{s,t-k}$ is included in addition to or instead of $CUM_NCE_{s,t-k}$, where

¹⁴ Source: Drugs@FDA Data Files

¹⁵ For example, the five levels associated with the chemical subgroup “nitrogen mustard analogues” are:

L	ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS
L01	ANTINEOPLASTIC AGENTS
L01A	ALKYLATING AGENTS
L01AA	Nitrogen mustard analogues
L01AA01	cyclophosphamide
L01AA02	chlorambucil
L01AA03	melphalan
L01AA05	chlormethine
L01AA06	ifosfamide
L01AA07	trofosfamide
L01AA08	prednimustine
L01AA09	bendamustine

$$\text{CUM_SUBGROUP}_{s,t-k} = \sum_g \text{IND_SUBGROUP}_{gs} \text{REGISTERED_SUBGROUP}_{g,t-k}$$

IND_SUBGROUP_{gs} = 1 if any drugs in chemical subgroup g are used to treat (indicated for) cancer at site s

= 0 if no drugs in chemical subgroup g are used to treat (indicated for) cancer at site s

$\text{REGISTERED_SUBGROUP}_{g,t-k}$ = 1 if any drugs in chemical subgroup g had been registered in Canada by the end of year $t-k$

= 0 if no drugs in chemical subgroup g had been registered in Canada by the end of year $t-k$

III. Data

NCE registrations in Canada (REGISTERED). Data on new chemical entities registered in Canada were constructed from the [Health Canada Drug Product Database](#), which contains product-specific information on drugs approved for use in Canada. The database is managed by Health Canada and includes human pharmaceutical and biological drugs, veterinary drugs and disinfectant products. It contains approximately 15,000 products which companies have notified Health Canada as being marketed.

Drug indications (IND). Data on drug indications were obtained from [Thériaque](#), a database of official, regulatory, and bibliographic information on all drugs available in France, intended for health professionals. This database is produced by the Centre National Hospitalier d'Information sur le Médicament. In this database, drugs are coded according to WHO ATC codes, and diseases are coded according to WHO ICD-10 codes.¹⁶ The drug indications listed in [Thériaque](#) are labeled indications, as defined by the [Collège de la Haute autorité de Santé](#).

Appendix Table 1 shows drugs (sorted by registration year) used to treat various types of cancer in Canada.

Premature mortality data (YPLL75, YPLL65, YPLL55). Data on years of potential life lost before ages 75, 65, and 55, by cancer site and year (2000-2011), were constructed from the

¹⁶ Many drug databases contain information about drug indications, but this information is usually in text form only.

[WHO Mortality Database](#), a compilation of mortality data by age, sex and cause of death, as reported annually by Member States from their civil registration systems.¹⁷

Cancer incidence data. Data on the number of new cancer cases, by cancer site, age, and year, were obtained from [CANSIM Table 103-0550](#).

Population data. Data on population, by age and year, were obtained from [CANSIM Table 051-0001](#).

IV. Empirical results

Estimates of the β_k parameters from eq. (1) and similar equations are shown in Table 1 and plotted (on an inverted scale) in Figure 7. Each estimate was obtained from a separate model. All equations include a cancer incidence measure (e.g. $\ln(\text{INC_RATE}_{75_{st}})$), cancer-site fixed effects and year fixed effects. To conserve space and simplify the presentation, estimates of the cancer incidence coefficient (γ) are not included in Table 1.¹⁸ None of the estimates of this coefficient were statistically significant, and controlling for cancer incidence had very little effect on the estimates of β_k . As discussed earlier, this may be due to offsetting effects of increases in I^* and increases in (I / I^*) on premature mortality.

In part A of Table 1 and Figure 7, the age threshold for calculating premature mortality is 75 years, i.e. the dependent variable is years of potential life lost before age 75. In model 1, the lag (k) from drug registrations to premature mortality equals zero, i.e. we are examining the effect of the cumulative number of drugs registered by the end of year t on premature mortality in year t . The estimate of β_0 is not statistically significant. In model 2, the lag is 5 years; the estimate of β_5 is also statistically insignificant. In models 3-6, the lags are 10, 15, 20, and 25 years, respectively. All of these coefficients are negative and highly statistically significant (p -value $< .0001$), indicating that premature mortality before age 75 is significantly inversely related to the cumulative number of drugs registered at least 10 years earlier. The estimate of β_{15} is the most statistically significant, and the magnitude of the point estimate of β_{15} is 66% larger than the magnitude of the point estimate of β_{10} . Since, as discussed earlier, mean utilization of

¹⁷ Mortality data are reported in 5-year age groups in the WHO Mortality Database. I assume that deaths in a 5-year age group occur at the midpoint of the age group. For example, I assume that deaths at age 35-39 years occurred at age 37.5. The Association of Public Health Epidemiologists in Ontario (2015) uses this method.

¹⁸ Complete estimates of model 4 are presented in Appendix Table 2.

drugs that have been marketed for less than 10 years is only one-sixth as great as mean utilization of drugs that have been marketed for at least a decade, it is not surprising that premature mortality is strongly inversely related only to the cumulative number of drugs that had been registered at least ten years earlier.

In parts B and C of Table 1 and Figure 7, the age thresholds for calculating premature mortality are 65 years and 55 years, respectively. The estimates based on these age thresholds are very similar to the estimates based on the age threshold of 75 years: premature mortality before age 65 and 55 is strongly inversely related only to the cumulative number of drugs that had been registered at least ten years earlier.

Figure 8 shows a bubble plot of the long-run (2000-2011) log change in YPLL before age 75 [$\ln(\text{YPLL}_{75,s,2011}) - \ln(\text{YPLL}_{75,s,2000})$] against the long-run change in the cumulative number of drugs registered 15 years earlier [$\text{CUM_NCE}_{s,1996} - \text{CUM_NCE}_{s,1985}$], i.e. the number of drugs registered during the period 1985-1996. The bubble size is proportional to the mean premature mortality rate during 2000-2011 ($(\sum_t \text{YPLL}_{75,it}) / 12$). This figure confirms the finding from model 4 in Table 1 of a highly significant inverse relationship. The point estimate of β_{15} from the long-difference model ($\beta_{15} = -0.0247$, $t\text{-value} = -4.42$; $p\text{-value} = 0.0010$) is similar to the point estimate of β_{15} from model 4 in Table 1.

Figure 8 reveals that the largest number of drugs that were launched during 1985-1996 were for cancers of lymphoid, haematopoietic and related tissue (ICD-10 block C81-C96), which include leukemia. In principle, it is possible that excluding this ICD-10 block could have a substantial effect on the estimates, although it does not appear from Figure 8 to be an outlier. As shown in Appendix Table 3, when ICD-10 block C81-C96 is excluded from the sample, estimates of eq. (1) are very similar to the estimates when it is included.

As discussed above, the hypothesis that premature mortality from a disease depends on the number of chemical (or pharmacological) *subgroups* that have previously been developed to treat the disease rather than, or in addition to, the number of chemical *substances* (drugs) that have previously been developed to treat the disease can be tested by estimating versions of eq. (1) in which $\text{CUM_SUBGROUP}_{s,t-k}$ is included instead of, or in addition to, $\text{CUM_NCE}_{s,t-k}$. Table 2 provides estimates of models suitable for testing this hypothesis. Estimates from three different models are presented there. In all three models, the dependent variable is $\ln(\text{YPLL}_{75,st})$ and $k = 15$. The first model shown is the same as model 4 in Table 1, in which the only

pharmaceutical variable is $CUM_NCE_{s,t-15}$. In the second model (model 19), $CUM_NCE_{s,t-15}$ is replaced by $CUM_SUBGROUP_{s,t-15}$. The coefficient on this variable is negative and significant, but it is less significant than the coefficient on $CUM_NCE_{s,t-15}$ in model 4. The third model (model 20) includes both $CUM_NCE_{s,t-15}$ and $CUM_SUBGROUP_{s,t-15}$. Controlling for the cumulative number of drugs, the cumulative number of chemical subgroups is not statistically significant. These estimates suggest that drugs (chemical substances) within the same class (chemical subgroup) are not “therapeutically equivalent,”¹⁹ i.e. they do not have essentially the same effect in the treatment of a disease or condition.

V. Discussion

During the period 2000-2011, the premature (before age 75) cancer mortality rate (the number of years of potential life lost due to cancer before age 75 per 100,000 population age 0-74) declined by about 9%. The estimates of model 4 imply that, in the absence of pharmaceutical innovation during the period 1985-1996, the premature cancer mortality rate would have *increased* about 12% during the period 2000-2011.^{20, 21} As shown in Figure 9, the premature mortality rate would have been 1788, rather than its actual value of 1459. In 2011, the population age 0-74 was about 32.1 million (or 321 hundred thousand), so the estimates of model 4 imply that *pharmaceutical innovation during the period 1985-1996 reduced the number of years of potential life lost to cancer before age 75 in 2011 by 105,366* ($= 321 * (1788 - 1459)$).

This reduction in premature mortality is an estimate of the *benefit* to Canadians below age 75 in 2011 of pharmaceutical innovation during the period 1985-1996. Now I will calculate an estimate of the (social) *cost* of this innovation. As shown in Appendix Table 1, 40 drugs that are used to treat cancer were registered during the period 1985-1996. Data from IMS Health indicate that in 2010,²² expenditure on products containing these molecules was 409 million

¹⁹ According to one medical dictionary, drugs that have “essentially the same effect in the treatment of a disease or condition” are therapeutically equivalent. Drugs that are therapeutically equivalent may or may not be chemically equivalent, bioequivalent, or generically equivalent. <http://medical-dictionary.thefreedictionary.com/therapeutic+equivalent>

²⁰ In Appendix Table 2, the point estimate of δ_{2000} is -0.1157 (p-value = 0.0003). If $CUM_NCE_{s,t-15}$ is excluded from the equation, the point estimate of δ_{2000} is 0.0873 (p-value = 0.0008).

²¹ An important possible reason why the premature cancer mortality rate would have increased in the absence of previous pharmaceutical innovation is a substantial decline in the “competing risk” of death from cardiovascular disease. See Honoré and Lleras-Muney (2006).

²² 2010 is the most recent year for which these data are available.

USD, which is about 1.9% of total Canadian drug expenditure (21.6 billion USD). About 70% of cancer patients were diagnosed before the age of 75, so it seems reasonable to assume that 288 million USD ($= 70\% * 409$ million USD) was spent on these drugs for cancer patients below the age of 75.²³ This implies that the cost per life-year before age 75 gained from previous pharmaceutical innovation was 2730 USD ($= 288$ million USD / 105,366 life-years).

Presumably most of the drugs registered during the period 1985-1996 were off-patent by 2010, so these cost estimates reflect prices of generic drugs. Law (2013) argues that Canadian generic drug prices have traditionally been set using a percentage of the equivalent brand-name price as a ceiling, and that typically, these percentages ranged between 60% and 70% of the brand price.²⁴ This suggests that if these drugs had been sold at branded rather than generic prices, the cost per life-year gained would have been between 3900 ($= 2730 / 60\%$) and 4550 ($= 2730 / 70\%$) USD. However, the ratio of generic price to branded price may be significantly lower for cancer drugs (which are often infused or injected) than it is for other drugs (which are primarily administered orally). For example, when imatinib, which is used to treat a set of leukemias, went generic in Canada in 2013, the generic drug price was approximately 25% of the branded price. If the generic/branded price ratio were 25%, and if these drugs had been sold at branded rather than generic prices, the cost per life-year gained would have been 10,920 ($= 2730 / 25\%$) USD.

Hirth et al (2000) performed a search of the value-of-life literature and identified 41 estimates of the value of life from 37 articles.²⁵ From estimates of the value of life, they calculated estimates of the value (in 1997 dollars) of a quality-adjusted life-year (QALY).²⁶ Four types of methods were used to produce those estimates: revealed preference/job risk, contingent valuation, revealed preference/non-occupational safety, and human capital. Median implied values (in 1997 and 2011 dollars²⁷) of a QALY estimated in those studies are shown in the following table.

²³ Since some of these drugs are used to treat diseases other than cancer, this is probably an overestimate.

²⁴ Law (2013) also argues that recent changes have moved these price ceilings lower in almost every province, to a nationwide low of 25% in Ontario.

²⁵ Twenty-eight of the reviewed articles used U.S. data; the remaining articles used data from the U.K. (4), Canada (3), France (1), and Denmark (1). National origin did not significantly affect the values.

²⁶ Lichtenberg (2009) demonstrated that, although the health of cancer patients is less than perfect, the number of QALYs gained from pharmaceutical innovation could be either greater than or less than the number of life-years gained.

²⁷ The U.S. Consumer Price Index increased by 40% between 1997 and 2011.

Study method	Number of studies	Median value of a QALY	
		1997 dollars	2011 dollars
revealed preference/job risk	19	\$428,286	\$600,243
contingent valuation	8	\$161,305	\$226,069
revealed preference/non-occupational safety	8	\$93,402	\$130,903
human capital	6	\$24,777	\$34,725

My estimate of the cost per life-year before age 75 gained from previous pharmaceutical innovation is well below even the lowest estimates of the value of a life-year saved.

VI. Summary and conclusions

The premature cancer mortality rate has been declining in Canada, and there has been considerable variation in the rate of decline across cancer sites. I analyzed the effect that pharmaceutical innovation has had on premature cancer mortality in Canada during the period 2000-2011, by investigating whether the cancer sites that experienced more pharmaceutical innovation had larger declines in the premature mortality rate, controlling for changes in the incidence rate.

The study is subject to several limitations. First, the measures of pharmaceutical innovation that were used were based only on labeled indications, but the National Cancer Institute (2015d) says that “off-label use of drugs is very common in cancer treatment.” Second, it was not possible to measure or control for non-pharmaceutical medical innovation. Third, the outcome measures used were life-years gained, not quality-adjusted life-years gained.

The estimates indicated that premature mortality before age 75 is significantly inversely related to the cumulative number of drugs registered at least 10 years earlier. Since mean utilization of drugs that have been marketed for less than 10 years is only one-sixth as great as mean utilization of drugs that have been marketed for at least a decade, it is not surprising that premature mortality is strongly inversely related only to the cumulative number of drugs that had been registered at least ten years earlier. Premature mortality before age 65 and 55 is also strongly inversely related to the cumulative number of drugs that had been registered at least ten

years earlier. None of the estimates of the effect of incidence on mortality were statistically significant.

Controlling for the cumulative number of drugs, the cumulative number of chemical subgroups does not have a statistically significant effect on premature mortality. This suggests that drugs (chemical substances) within the same class (chemical subgroup) are not therapeutically equivalent.

During the period 2000-2011, the premature (before age 75) cancer mortality rate declined by about 9%. The estimates imply that, in the absence of pharmaceutical innovation during the period 1985-1996, the premature cancer mortality rate would have increased about 12% during the period 2000-2011. A substantial decline in the “competing risk” of death from cardiovascular disease could account for this. The estimates imply that pharmaceutical innovation during the period 1985-1996 reduced the number of years of potential life lost to cancer before age 75 in 2011 by 105,366.

The cost per life-year before age 75 gained from previous pharmaceutical innovation is estimated to have been 2730 USD. Most of the previously-registered drugs were off-patent by 2011, but evidence suggests that, even if these drugs had been sold at branded rather than generic prices, the cost per life-year gained would have been below 11,000 USD, a figure well below even the lowest estimates of the value of a life-year gained.

References

- Association of Public Health Epidemiologists in Ontario (2015). Calculating Potential Years of Life Lost (PYLL), <http://www.apheo.ca/index.php?pid=190>
- Brustugun OT, Møller B, Helland Å (2014). Years of life lost as a measure of cancer burden on a national level. *British Journal of Cancer* 111, 1014–1020, <http://www.nature.com/bjc/journal/v111/n5/full/bjc2014364a.html>
- Burnet NG, Jefferies SJ, Benson RJ, Hunt DP, Treasure FP (2005). Years of life lost (YLL) from cancer is an important measure of population burden--and should be considered when allocating research funds. *British Journal of Cancer* 92 (2): 241–5, January, <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2361853/>
- Center for Medicare and Medicaid Services (2013), National Health Expenditure Data, <http://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/Downloads/tables.pdf>.
- Commonwealth Fund (2011). *The U.S. Health System in Perspective: A Comparison of Twelve Industrialized Nations*.
- Dorsey ER, et al (2010). Financial Anatomy of Biomedical Research, 2003 – 2008. *Journal of the American Medical Association* 303(2), 137–143.
- Gardner JW, Sanborn JS (1990). Years of potential life lost (YPLL)--what does it measure? *Epidemiology* 1(4): 322-9, July.
- Hirth RA, Chernew ME, Miller E, Fendrick AM, Weissert WG (2000), Willingness to Pay for a Quality-adjusted Life Year: In Search of a Standard, *Med Decis Making* 20(3): 332-342, Jul-Sep, <http://users.phhp.ufl.edu/jharman/healthecon/Hirth%20WTP%20QALY.pdf>
- Honoré BE, Lleras-Muney A (2006). Bounds in Competing Risks Models and the War on Cancer. *Econometrica* 74 (6): 1675-1698, November.
- Jalan J, Ravallion M (2001) Does Piped Water Reduce Diarrhea for Children in Rural India?, World Bank Development Research Group, Policy Research Working Paper 2664, http://www1.worldbank.org/prem/poverty/ie/dime_papers/332.pdf
- Law MR (2013). Money Left on the Table: Generic Drug Prices in Canada. *Healthcare Policy* 8(3): 17–25, February, <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3999558/#!po=44.7368/>
- Lichtenberg FR (2014a). [The impact of pharmaceutical innovation on longevity and medical expenditure in France, 2000–2009](#), *Economics and Human Biology* 13: 107-127, March.

Lichtenberg FR (2014b). [Has Medical Innovation Reduced Cancer Mortality?](#) *CEsifo Economic Studies* (2014) 60 (1): 135-177.

National Cancer Institute (2015a). Cancer Screening Overview, <http://www.cancer.gov/cancertopics/pdq/screening/overview/patient/page5>

National Cancer Institute (2015b). Drug Discovery at the National Cancer Institute, <http://www.cancer.gov/cancertopics/factsheet/NCI/drugdiscovery>

National Cancer Institute (2015c). Person-Years of Life Lost. *Cancer Trends Progress Report, 2009/2010 Update*, http://progressreport.cancer.gov/doc_detail.asp?pid=1&did=2007&chid=76&coid=730&mid=#estimate

National Cancer Institute (2015d). Off-Label Drug Use in Cancer Treatment, <http://www.cancer.gov/about-cancer/treatment/drugs/off-label>

Nobelprize.org (2015). The Prize in Economics 1987 - Press Release". http://www.nobelprize.org/nobel_prizes/economic-sciences/laureates/1987/press.html

Renard F, Tafforeau J, Deboosere P (2014). Premature mortality in Belgium in 1993-2009: leading causes, regional disparities and 15 years change. *Arch Public Health*. 2014 Oct 1;72(1):34.

Romer P (1990). Endogenous Technological Change. *Journal of Political Economy* 98 (5, Part 2), S71-S102.

Sampat BN, Lichtenberg FR (2011). What are the Respective Roles of the Public and Private Sectors in Pharmaceutical Innovation? *Health Affairs* 30(2), 332-9.

Solow RM (1960). Investment and technological progress, in: Arrow, K., Karlin, S., Suppes, P. (eds.), *Mathematical Methods in Social Sciences 1959*, Stanford University Press, 89-104.

Stukel TA, Fisher ES, Wennberg DE, Alter DA, Gottlieb DJ, Vermeulen MJ. (2007) Analysis of Observational Studies in the Presence of Treatment Selection Bias: Effects of Invasive Cardiac Management on AMI Survival Using Propensity Score and Instrumental Variable Methods. *JAMA* 297(3):278-285, <http://jama.jamanetwork.com/article.aspx?articleid=205172>

U.S. Food and Drug Administration (2015), Table of Pharmacogenomic Biomarkers in Drug Labeling, <http://www.fda.gov/drugs/scienceresearch/researchareas/pharmacogenetics/ucm083378.htm>

World Health Organization (2015), ICD-10 Version 2015, Chapter II, Neoplasms, <http://apps.who.int/classifications/icd10/browse/2015/en#/II>

Figure 1
Trends in premature cancer mortality and cancer incidence, Canada, 1997-2006





Figure 2

Log change from 2000 to 2011 in the premature mortality rate, by type of cancer, Canada



The premature mortality rate is the number of potential years of life lost before age 75 per 100,000 population age 0-74.

Figure 3
Cumulative number of drugs launched in Canada
for 5 types of cancer, 5-year intervals, 1988-2013




Figure 4
Plot of the residuals from the unweighted regression
 $[\ln(YPLL75_{st}) = \alpha_s + \delta_t + \varepsilon_{st}]$ on the
mean premature mortality rate $(\sum_t YPLL75_{it}) / 12$





Figure 5
Estimates of the relative utilization index (year 0 = 1.0)



The relative utilization index is the mean ratio of the number of units of a molecule sold y years after registration to the number of units of the same molecule sold in the year that it was registered. These estimates are based on data on 25 molecules used to treat cancer that were registered after 1998. The figure provides a conservative estimate of the slope of the age-utilization profile, because there was zero utilization of many of these molecules in the first few years after they were registered.

Figure 6
Mean number of standard units of cancer drugs sold (in thousands) in 2010,
by period of launch in Canada




Figure 7
 Estimates of the β_k parameters from eq. (1) and similar equations






Figure 8

Relationship across cancer sites between the number of drugs launched during 1985-1996 and the 2000-2011 log change in the premature (before age 75) mortality rate





Figure 9
Premature (before age 75) cancer mortality rate:
actual vs. estimated in the absence of previous pharmaceutical innovation



The premature (before age 75) cancer mortality rate is the number of years of potential life lost due to cancer before age 75 per 100,000 population age 0-74.

Table 1

Estimates of the β_k parameters from eq. (1) and similar equations

Model	Parameter	Lag	Estimate	Standard Error	Z	Pr > Z
A. Dependent variable: $\ln(YPLL75_{st})$ Weight: $((\sum_t YPLL75_{it}) / 12)$						
1	β_0	0	-0.003	0.009	-0.33	0.7385
2	β_5	5	-0.006	0.008	-0.73	0.4659
3	β_{10}	10	-0.013	0.003	-4.69	<.0001
4	β_{15}	15	-0.021	0.002	-8.96	<.0001
5	β_{20}	20	-0.019	0.003	-7.58	<.0001
6	β_{25}	25	-0.023	0.003	-8.09	<.0001
B. Dependent variable: $\ln(YPLL65_{st})$ Weight: $((\sum_t YPLL65_{it}) / 12)$						
7	β_0	0	-0.006	0.009	-0.61	0.5427
8	β_5	5	-0.006	0.007	-0.88	0.3784
9	β_{10}	10	-0.012	0.003	-4.80	<.0001
10	β_{15}	15	-0.023	0.003	-6.78	<.0001
11	β_{20}	20	-0.022	0.003	-8.72	<.0001
12	β_{25}	25	-0.026	0.003	-10.74	<.0001
C. Dependent variable: $\ln(YPLL55_{st})$ Weight: $((\sum_t YPLL55_{it}) / 12)$						
13	β_0	0	-0.013	0.009	-1.43	0.1519
14	β_5	5	-0.014	0.009	-1.62	0.1045
15	β_{10}	10	-0.016	0.005	-3.19	0.0014
16	β_{15}	15	-0.024	0.004	-6.10	<.0001
17	β_{20}	20	-0.025	0.004	-6.92	<.0001
18	β_{25}	25	-0.029	0.005	-5.53	<.0001

Each estimate was obtained from a separate model. All equations include a cancer incidence measure (e.g. $\ln(INC_RATE75_{it})$), cancer-site fixed effects and year fixed effects. Standard errors are clustered within cancer sites.

Table 2

Estimates of models of years of potential life lost before age 75, including cumulative number of drugs, cumulative number of chemical subgroups, or both

Dependent variable: $\ln(YPLL75_{st})$ Weight: $(\sum_t YPLL75_{it}) / 12$

Model	Regressor	Estimate	Standard Error	Z	Pr > Z
4	CUM_NCE _{s,t-15}	-0.021	0.002	-8.96	<.0001
19	CUM_SUBGROUP _{s,t-15}	-0.023	0.009	-2.55	0.0107
20	CUM_NCE _{s,t-15}	-0.024	0.004	-6.78	<.0001
20	CUM_SUBGROUP _{s,t-15}	0.010	0.012	0.83	0.4042

All equations include a cancer incidence measure (e.g. $\ln(INC_RATE75)$), cancer-site fixed effects and year fixed effects. Standard errors are clustered within cancer sites.

Appendix Table 1

Drugs (sorted by launch year) used to treat various types of cancer in Canada

Drug	Canada launch year	C00-C14 lip, oral cavity and pharynx	C15-C26 digestive organs	C30-C39 respiratory system and intrathoracic organs	C40-C41 bone and articular cartilage	C43-C44 skin	C45-C49 connective and soft tissue	C50-C50 breast	C51-C58 female genital organs	C60-C63 male genital organs	C64-C68 urinary organs	C69-C72 eye, brain and central nervous system	C73-C75 endocrine glands and related structures	C76-C80 secondary and ill-defined	C81-C96 stated or presumed to be primary, of lymphoid, haematopoietic and related tissue	C97 independent (primary) multiple sites
J06BA01 Immunglobulins, normal human, for extravascular adm.	1951														X	
L01AA05 Chlormethine	1951														X	
H02AB09 Hydrocortisone	1954											X		X		
L01AB01 Busulfan	1954														X	
L01BB02 Mercaptopurine	1954														X	
G03CA03 Estradiol	1955									X						
L01BA01 Methotrexate	1955			X	X		X	X	X		X	X		X	X	
L02AA01 Diethylstilbestrol	1956									X						
L01AA02 Chlorambucil	1957														X	
H02AB02 Dexamethasone	1958	X	X	X	X	X	X								X	
H02AB08 Triamcinolone	1958			X										X		
L01AA01 Cyclophosphamide	1959			X			X	X	X	X	X	X	X	X	X	
L01AC01 Thiotepa	1959							X	X		X					
H02AB04 Methylprednisolone	1960	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
H02AB01 Betamethasone	1961	X	X	X	X	X	X					X		X	X	
L01DA01 Dactinomycin	1963						X		X	X	X					
L01AA03 Melphalan	1964						X	X	X			X	X		X	
D07AC01 Betamethasone	1965														X	
H02AB06 Prednisolone	1967											X		X		
C03CC01 Etacrynic acid	1968													X		
L01XB01 Procarbazine	1970			X								X		X	X	
L01DB02 Daunorubicin	1971						X								X	
L01DC01 Bleomycin	1973			X			X		X	X				X	X	
L01BB03 Tioguanine	1974														X	
L01XX02 Asparaginase	1974														X	
L01AD01 Carmustine	1975					X						X			X	
L01BC02 Fluorouracil	1975	X	X	X				X	X					X		
L01AD02 Lomustine	1976			X		X						X		X	X	
L01AX04 Dacarbazine	1976					X	X								X	
L01DB01 Doxorubicin	1976		X	X	X		X	X	X		X				X	
L01BC01 Cytarabine	1977														X	
L01DC03 Mitomycin	1977		X					X			X			X		
B01AB04 Dalteparin	1979	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
L01XX05 Hydroxycarbamide	1979														X	
L01XX23 Mitotane	1979												X			
V03AF03 Calcium folinate	1980	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
L01CB01 Etoposide	1981			X				X	X	X				X	X	
L01CA03 Vindesine	1982	X	X	X				X						X	X	

Appendix Table 1

Drugs (sorted by launch year) used to treat various types of cancer in Canada

Drug	Canada launch year	C00-C14 lip, oral cavity and pharynx	C15-C26 digestive organs	C30-C39 respiratory system and intrathoracic organs	C40-C41 bone and articular cartilage	C43-C44 skin	C45-C49 connective and soft tissue	C50-C50 breast	C51-C58 female genital organs	C60-C63 male genital organs	C64-C68 urinary organs	C69-C72 eye, brain and central nervous system	C73-C75 endocrine glands and related structures	C76-C80 secondary and ill-defined	C81-C96 stated or presumed to be primary, of lymphoid, haematopoietic and related tissue	C97 independent (primary) multiple sites
H05BA01 Calcitonin (salmon synthetic)	1983													X	X	
L01XX01 Amsacrine	1983														X	
L02BG01 Aminogluthetimide	1983												X			
L01CA02 Vincristine	1984			X	X		X	X	X		X	X	X	X	X	
L01CB02 Teniposide	1984								X		X				X	
L01DB07 Mitoxantrone	1984							X		X					X	
L02BB01 Flutamide	1984									X						
D05BA02 Methoxsalen	1985														X	
L01AD04 Streptozocin	1985		X													
L01DB03 Epirubicin	1985		X	X			X	X	X		X			X	X	
L01XA01 Cisplatin	1985		X	X					X	X	X					
L02AE02 Leuprorelin	1985							X		X						
D05AD02 Methoxsalen, topical	1986														X	
J06BA02 Immunoglobulins, normal human, for intravascular adm.	1986														X	
L01XA02 Carboplatin	1986	X		X					X					X		
L03AB05 Interferon alfa-2b	1986					X									X	
G03HA01 Cyproterone	1987									X						
L02AE01 Buserelin	1988									X						
H01CB02 Octreotide	1989		X											X		
L01AA06 Ifosfamide	1989			X	X		X	X	X	X				X	X	
L03AB04 Interferon alfa-2a	1989					X	X				X				X	
B03XA01 Erythropoietin	1990	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
J02AC01 Fluconazole	1990	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
L03AX03 BCG vaccine	1990										X					
L01DB06 Idarubicin	1991														X	
L01BB05 Fludarabine	1992														X	
L01CA01 Vinblastine	1992						X	X	X	X	X				X	
L01CD01 Paclitaxel	1992			X			X	X	X					X		
L02BB02 Nilutamide	1992									X						
M05BA03 Pamidronic acid	1992							X						X	X	
L01BB04 Cladribine	1993														X	
L01XX08 Pentostatin	1993														X	
L01XX11 Estramustine	1993									X						
L01CA04 Vinorelbine	1994			X				X						X		
L02AE03 Goserelin	1994							X		X						
L02BG02 Formestane	1994							X								
L01CD02 Docetaxel	1995	X	X	X				X		X				X		
L01XD01 Porfimer sodium	1995		X	X										X		

Appendix Table 2

Complete estimates of model 4

Dependent variable: $\ln(YPLL75_{st})$ Weight: $(\sum_t YPLL75_{it}) / 12$

Parameter	Estimate	Standard Error	Z	Pr > Z
CUM_NCE _{s,t-15}	-0.021	0.002	-8.96	<.0001
$\ln(INC_RATE75_{st})$	0.064	0.116	0.55	0.5831
Year 2000	-0.116	0.032	-3.61	0.0003
Year 2001	-0.113	0.036	-3.16	0.0016
Year 2002	-0.118	0.035	-3.38	0.0007
Year 2003	-0.121	0.029	-4.25	<.0001
Year 2004	-0.095	0.026	-3.68	0.0002
Year 2005	-0.072	0.022	-3.35	0.0008
Year 2006	-0.079	0.019	-4.12	<.0001
Year 2007	-0.042	0.016	-2.60	0.0092
Year 2008	-0.044	0.021	-2.10	0.0359
Year 2009	-0.049	0.022	-2.18	0.0290
Year 2010	-0.016	0.015	-1.08	0.2821
Year 2011	0.000	0.000	.	.
group C00-C14 lip, oral cavity and pharynx	-2.154	0.156	-13.81	<.0001
group C15-C26 digestive organs	0.285	0.116	2.45	0.0141
group C30-C39 respiratory system and intrathoracic organs	0.567	0.085	6.70	<.0001
group C40-C41 bone and articular cartilage	-3.028	0.403	-7.52	<.0001
group C43-C44 skin	-2.048	0.124	-16.47	<.0001
group C45-C49 connective and soft tissue	-1.979	0.248	-7.98	<.0001
group C50-C50 breast	-0.434	0.089	-4.90	<.0001
group C51-C58 female genital organs	-1.011	0.060	-16.87	<.0001
group C60-C63 male genital organs	-2.132	0.089	-23.94	<.0001
group C64-C68 urinary organs	-1.481	0.065	-22.75	<.0001
group C69-C72 eye, brain and central nervous system	-1.069	0.169	-6.31	<.0001
group C73-C75 endocrine glands and related structures	-3.277	0.143	-22.93	<.0001
group C76-C80 secondary and ill-defined	-0.965	0.123	-7.82	<.0001
group C81-C96 stated or presumed to be primary, of lymphoid, haematopoietic and related tissue	0.000	0.000	.	.
Intercept	5.730	0.397	14.44	<.0001

Standard errors are clustered within cancer sites.

Appendix Table 3

Estimates of the β_k parameters from eq. (1) and similar equations, including and excluding C81-C96 lymphoid, haematopoietic and related tissue

Model	Parameter	Lag	Estimate	Std. Err.	Z	Pr > Z	Estimate	Std. Err.	Z	Pr > Z	
			Include C81-C96 lymphoid, haematopoietic and related tissue				Exclude C81-C96 lymphoid, haematopoietic and related tissue				
A. Dependent variable: $\ln(YPLL75_{st})$ Weight: $((\sum_t YPLL75_{it}) / 12)$											
1	β_0	0	-0.003	0.009	-0.33	0.7385	0.004	0.008	0.55	0.585	
2	β_5	5	-0.006	0.008	-0.73	0.4659	-0.008	0.007	-1.06	0.291	
3	β_{10}	10	-0.013	0.003	-4.69	<.0001	-0.012	0.003	-3.74	0.000	
4	β_{15}	15	-0.021	0.002	-8.96	<.0001	-0.020	0.003	-5.91	<.0001	
5	β_{20}	20	-0.019	0.003	-7.58	<.0001	-0.017	0.006	-3.07	0.002	
6	β_{25}	25	-0.023	0.003	-8.09	<.0001	-0.029	0.009	-3.18	0.002	
B. Dependent variable: $\ln(YPLL65_{st})$ Weight: $((\sum_t YPLL65_{it}) / 12)$											
7	β_0	0	-0.006	0.009	-0.61	0.5427	0.006	0.010	0.56	0.579	
8	β_5	5	-0.006	0.007	-0.88	0.3784	-0.007	0.008	-0.93	0.353	
9	β_{10}	10	-0.012	0.003	-4.80	<.0001	-0.011	0.003	-3.43	0.001	
10	β_{15}	15	-0.023	0.003	-6.78	<.0001	-0.019	0.004	-5.60	<.0001	
11	β_{20}	20	-0.022	0.003	-8.72	<.0001	-0.017	0.005	-3.33	0.001	
12	β_{25}	25	-0.026	0.003	-10.74	<.0001	-0.030	0.009	-3.46	0.001	
C. Dependent variable: $\ln(YPLL55_{st})$ Weight: $((\sum_t YPLL55_{it}) / 12)$											
13	β_0	0	-0.013	0.009	-1.43	0.1519	-0.003	0.014	-0.18	0.860	
14	β_5	5	-0.014	0.009	-1.62	0.1045	-0.013	0.009	-1.39	0.166	
15	β_{10}	10	-0.016	0.005	-3.19	0.0014	-0.014	0.004	-3.81	0.000	
16	β_{15}	15	-0.024	0.004	-6.10	<.0001	-0.022	0.006	-4.05	<.0001	
17	β_{20}	20	-0.025	0.004	-6.92	<.0001	-0.031	0.007	-4.49	<.0001	
18	β_{25}	25	-0.029	0.005	-5.53	<.0001	-0.043	0.010	-4.15	<.0001	

Each estimate was obtained from a separate model. All equations include a cancer incidence measure (e.g. $\ln(INC_RATE75_{st})$), cancer-site fixed effects and year fixed effects. Standard errors are clustered within cancer sites.