This PDF is a selection from a published volume from the National Bureau of Economic Research

Volume Title: American Agriculture, Water Resources, and Climate Change

Volume Authors/Editors: Gary D. Libecap and Ariel Dinar, editors

Volume Publisher: University of Chicago Press

Volume ISBNs: 978-0-226-83061-2 (cloth); 978-0-226-83062-9 (electronic)

Volume URL: https://www.nber.org/books-andchapters/american-agriculture-water-resources-and-climatechange

Conference Date: May 12-13, 2022

Publication Date: December 2023

Chapter Title: Cover Crops, Drought, Yield, and Risk: An Analysis of US Soybean Production

Chapter Author(s): Fengxia Dong

Chapter URL: https://www.nber.org/books-andchapters/american-agriculture-water-resources-and-climatechange/cover-crops-drought-yield-and-risk-analysis-ussoybean-production

Chapter pages in book: p. 241 - 267

Cover Crops, Drought, Yield, and Risk An Analysis of US Soybean Production

Fengxia Dong

7.1 Introduction

Climate change has caused increasing frequency and severity of drought stress in the US. Water scarcity has become one of the most severe constraints to agricultural production, adversely affecting crop yields and presenting a major challenge to sustainable food production. Along with prescribed grazing, mulching, micro-irrigation, and conservation tillage, cover cropping is among the five short- and long-term strategies for dealing with drought conditions that farmers can receive financial assistance for from the Natural Resources Conservation Service (NRCS) (USDA Climate Hubs 2021). Cover crops are defined by USDA NRCS in Cover Crop Termination Guidelines Version 4 (2019) as "crops including grasses, legumes and forbs for seasonal cover and other conservation purposes. Cover crops are primarily used for erosion control, soil health improvement, weed and other pest control, habitat for beneficial organisms, improved water efficiency, nutrient cycling, and water quality improvement. A cover crop managed and terminated according to these Guidelines is not considered a 'crop' for crop insurance purposes."

Legumes and grasses are currently the two most popular cover crop types

Fengxia Dong is a Research Agricultural Economist in the Agricultural Policy and Models Branch of the Market and Trade Economics Division at USDA's Economic Research Service (ERS).

The findings and conclusions in this paper are those of the author and should not be construed to represent any official USDA or US Government determination or policy. For acknowledgments, sources of research support, and disclosure of the author's material financial relationships, if any, please see https://www.nber.org/books-and-chapters/american -agriculture-water-resources-and-climate-change/cover-crops-drought-yield-and-risk -analysis-us-soybean-production.

(SARE 2007). Hairy vetch is the most widely used winter annual legume in northern regions because of its high N content, winter hardiness, and high productivity (Lu et al. 2000), and crimson clover is considered one of the best cover crops for southern regions due to its fast matureness and large N addition to the following crops (SARE 2007). Grass cover crops include annual cereals (such as rye, wheat, barley, and oats), annual or perennial forage grasses (such as ryegrass), and warm-season grasses (such as sorghum–Sudan grass) (SARE 2007). Besides legumes and grasses, buckwheat and *Brassica* (such as mustard, rapeseed, and forage radish) can also be used as cover crops (SARE 2007).

The use of cover crops as a cropping strategy is not new. It was practiced by people in ancient Greece, Rome, and China as early as 3,000 years ago (Langdale et al. 1991). Cover crops were first used in the US in the 18th century and extensively expanded in the 19th century (Groff 2015), although by that time they were used mainly as green manures. The affordability and ease of use of synthetic fertilizer at the end of World War II, however, attracted farmers to utilize more synthetic fertilizer instead of cover crops to further improve crop yields. The use of cover crops in conventional agriculture has gradually become less common since then (Groff 2015).

Currently, the adoption rate of cover crops is low and varies by agricultural commodity type. According to Agricultural Resource Management Surveys (ARMS) in years of 2010, 2016, 2017, and 2018, the adoption rate ranged from just over 5 percent of acreage on corn-for-grain (2016) to 8.4 percent on soybeans (2018), around 13 percent on cotton (2015), and over 24 percent on corn-for-silage (2016) (Wallander et al. 2021), in stark contrast to the adoption rate of conservation tillage, for example, which is 67 percent of soybean acreage, according to USDA's 2018 ARMS Phase II, Soybean Production Practices and Costs Report.

Cover crops can protect and improve soil between periods of regular crop production (Schnepf and Cox 2006). Besides a variety of production, soil health, and environmental benefits such as increasing weed and pest suppression, reducing runoff of sediments and nutrients into waterways, and reducing soil erosion and compaction, cover crops can improve water infiltration, reduce water evaporation, and increase soil's water holding capacity (e.g., USDA NRCS 2018; Mitchell, Shrestha, and Irmak 2015; Blanco-Canqui et al. 2015; McDaniel, Tiemann, and Grandy 2014; Laloy and Bielders 2010; Dean and Weil 2009; Sainju, Singh, and Whitehead 2002; Sainju et al. 2006).

There are, however, well-recognized trade-offs and limitations in adopting cover crops as a conservation strategy (SARE 2017). In addition to the costs of soil preparation, seeds, and labor, there are challenges in implementation and management, such as the selection of cover crop species, planting and termination time—which may interfere with fall harvest or spring planting—and producing too much surface residue (CTIC 2015; Sackett

2013; Miller, Chin, and Zook 2012; Snapp et al. 2005). Moreover, there are concerns that water needed by cover crops may reduce the amount of water available to the following main crop (SARE 2017; Clark et al. 1997; Corak, Frye, and Smith 1991; Ebelhar, Frye, and Blevins 1984; Munawar et al. 1990).

Along with the above concerns, the effect on crop yield and risk is another important factor in farmers' adoption decision of cover crops, as yield and risk directly affect farmers' economic returns. The Iowa Farm and Rural Life Poll in 2015 (Arbuckle 2016) reveals that 74 percent of the farmers believe that economic factors have a moderate to very strong influence on their changes in management practices. The 2017 Cover Crop Survey conducted by SARE (2017) also shows that the fear of a lack of economic returns (54 percent of respondents), increasing production risk (48 percent of respondents), and potential yield reduction (44 percent) are among the major concerns for non-users. Another cover crop survey conducted in 2015 shows that the potential yield benefit to cash crops is an important factor in decision making, especially for non-adopters (CTIC 2015). Therefore, an analysis of the effects of management practices on yield and its risk is essential to find effective supporting programs to promote good management practice adoption.

The results of existing studies of yield effects of cover crops are mixed. A meta-analysis of the response of corn yield to cover crops by Miguez and Bollero (2005) concludes that legume cover crops increase corn yield by 37 percent. Similarly, Andraski and Bundy (2005) and Muñoz et al. (2014) find a positive effect of cover crop biomass on corn yields. Contrastingly, Reddy (2017) discovers lower soybean yield with cover crops compared with no cover crops. Nielsen et al. (2016) find that there was an average 10 percent reduction in wheat yield following a cover crop compared with following fallow, regardless of whether the cover crop was grown in a mixture or in a single-species planting; in addition, yield reductions were greater under drier conditions. In comparison, Acharya et al. (2019); Smith, Atwood, and Warren (2014); Hunter et al. (2019), and Acuña and Villamil (2014) locate no benefits of growing cover crops on subsequent crop yield. Note that all the findings are subject to certain conditions, such as soil types, other production practices (e.g., tillage), cover crop species, and precipitation.

Previous studies of yield effects of cover crops are mainly conducted in field experimental plots using agronomic models. A study based on a large number of fields with different agroecological characteristics and under varied weather is in need. In addition, increasing the frequency and severity of adverse events can expose farms to significant production uncertainty. Therefore, special attention is paid to downside risk exposure. In general, the downside risk is the risk associated with unfavorable events and located in the lower tail of the yield or return distribution (Kim et al. 2014). As pointed out by Hardaker et al. (2004), Kim et al. (2014), and OECD (2011), analyzing both the exposure to risk and levels of downside risk in agriculture is a key component in assessing welfare impacts. While cover cropping is recommended to farmers to deal with drought, its effects on farm yield risk and especially downside risk are not well documented. This study aims to fill the literature gap by analyzing the effects of cover crops on yield and its risk with varied weather, regional, and field characteristics.

In this study, we focus on US soybean production. The US is the world's second-largest soybean producer and exporter, accounting for 31 percent of world total production and 36 percent of world total exports, respectively. US farmers planted 87.2 million acres of soybeans in 2021, behind only corn. The growth and productivity of soybeans are adversely affected by various environmental stresses, among which drought stress is considered the most devastating event (Le et al. 2012; Shaheen et al. 2016). Drought stress, especially occurring at late vegetative stages, may cause significant soybean production losses of up to 40 percent (Specht, Hume, and Kumudin 1999; Le et al. 2012) by inhibiting increases in the soybean plant height and leaf area (Dong et al. 2019). Several studies find that cover crops improve soybean soil moisture (e.g., Acharya et al. 2019; Chu et al. 2017), although some do not (Barker et al. 2018).

The paper makes three contributions. First, it explores the factors that affect the adoption of cover cropping. We consider not only land characteristics and farmers' demographics and concerns but also droughts in previous years. This enables us to reveal whether farmers view cover cropping as an effective means of increasing resilience to drought. Second, the paper examines the effects of cover crops on yield variation and downside risk. We employ moments of yield distribution to evaluate the exposure to yield variation and downside risk. Disentangling the yield effects of adaptation is of paramount importance. It will reveal whether farmers who adopt cover crops are indeed getting benefits in terms of an increase in crop yield, a benefit crucial to broader adoption. Third, the paper utilizes a data set covering the majority of soybean fields in the US with significantly different soil types and weather conditions. Two interplays, one between soil types and the adaptation strategy (namely, cover cropping), and the other between weather and the cover cropping practice, are included in yield and its risk analysis. The interplay along with farmer demographics, farm characteristics, and input use allow us to examine the effectiveness of the managerial options for risk mitigation under varied soil and weather conditions, especially in the threat of drought.

The rest of the paper is organized as follows. The next section discusses the theoretical framework and empirical models, followed by a description of data and variables. Then estimation results are discussed, followed by conclusions at the end.

7.2 Theoretical Framework and Empirical Models

Consider a farmer who uses a vector of inputs **x** and drought adaptation strategies (e.g., cover cropping, mulching, or drought-resistant seeds) C to produce a single output Q through a technology described by a wellbehaved (i.e., continuous and twice differentiable) production function $Q(\cdot)$. The farmer can choose to adopt (C = 1) or not (C = 0) a drought adaptation strategy. Use *e* to indicate random and uncontrollable factors reflecting production risk (e.g., drought effect) whose distribution is F(e). The production technology can thus be represented by $Q = Q(\mathbf{x}, C, e)$. Use *p* to indicate the output price and *w* a vector of input prices. The net return is represented by $\pi = pQ(\mathbf{x}, C, e) - w\mathbf{x}$.

Let $U(\pi)$ be a von Neumann-Morgenstern utility function that represents farmers' preferences regarding income. To simplify the analysis, I assume that the only risk that farmers are facing is production risk and both output and input prices are given or nonrandom. Being risk averse, farmers are assumed to maximize expected utility $EU(\pi)$, where $EU(\pi) = \int U(\pi) dF(e)$ with E as the expectation operator. For the decision on the drought adaptation strategy C, for example, if $EU(\pi \mid C = 1) - EU(\pi \mid C = 0) > 0$, then the farmer would choose to adopt the drought adaptation strategy; otherwise, the farmer would choose not to do so. In addition, the greater the difference between the expected utilities, the higher the probability of adoption. Very often there is a requirement of investment and/or possible uncertainty in profit due either to a lack of the exact performance of the adaptation strategy or to the higher probability of erring in the use of the adaptation strategy (Koundouri, Nauges, and Tzouvelekas 2006). In those cases, the farmer may choose to delay the adoption to achieve more information (Koundouri, Nauges, and Tzouvelekas 2006). Consequently, the farmer will choose the adaptation strategy iff $EU(\pi \mid C = 1) - EU(\pi \mid C = 0) > V$, where $V \ge 0$ is the value of new information essential for the farmer to make adoption decision which depends on the investment, the uncertainty related to the use of the strategy, and the farmers' characteristics (Koundouri, Nauges, and Tzouvelekas 2006). Therefore, drought adaptation strategies that require less investment and have less uncertainty in profit will have a higher level of adoption. For example, if technical assistance and extension service are provided to farmers for adopting an adaptation strategy, then the uncertainty in profit will be lower. Consequently, it is more possible that farmers will choose to adopt the strategy. In addition, farmers' characteristics such as their education level or their concerns about the environment may also play a role in the adoption decision. The more concerned the farmer is about an environmental issue, the higher probability of adopting a practice that can address the concern.

With differentiability of $U(\pi)$, $EU(\pi)$ can be approximated by taking the

expectation of an *k*th-order Taylor series expansion of $U(\pi)$ at the mean net return $E\pi$ where $E\pi = \int \pi d\mathcal{F}(\epsilon) = u_1$ and is written as

(1)
$$EU(\pi) \approx E\left[\sum_{j=0}^{k} \left(\frac{1}{j!} \frac{\partial^{j}U}{\partial \pi^{j}}(u_{1}) \times (\pi - u_{1})^{j}\right)\right]$$
$$= U(u_{1}) + \sum_{j=1}^{k} \left[\frac{1}{j!} \frac{\partial^{j}U}{\partial \pi^{j}}(u_{1}) \times E(\pi - u_{1})^{j}\right]$$
$$= U(u_{1}) + \sum_{j=1}^{k} \left[\frac{1}{j!} \frac{\partial^{j}U}{\partial \pi^{j}}(u_{1}) \times E(\pi - u_{1})^{j}\right]$$

Equation (1) shows that the expected utility depends on the mean net return u_1 and the *j*th (j = 2, 3, ..., k) central moment of net return, $u_j = E[(\pi - u_1)^j]$. When $j=2, u_j$ is the second moment or the variance, and when j = 3, the third moment or the skewness of the net return. The skewness measures the asymmetry of the distribution around its mean, with a negative skewness implying a distribution skewed to the left; and a positive one implying a distribution skewed to the right. A lower skewness generates a greater exposure to downside risk.

By normalizing prices so that p = 1, a farm's net return can be expressed by $\pi = Q(\mathbf{x}, C, \mathbf{e}) - (\mathbf{wx} / p)$. The equation explicitly shows that the production function $Q(\mathbf{x}, C, \mathbf{e})$ provides all the relevant information for analyzing risk exposure on farms adopting drought adaptation strategies. To empirically investigate the impacts of cover crops on crop yield and yield risks, we start with the moment functions of crop yield.

7.2.1 Moment Representation of Production Function

Our empirical model is based on Antle's (1983) moment-based approach, which provides a flexible and convenient basis for evaluating exposure to production risk. As discussed in Antle (1983), a stochastic production function can be represented by a general parameterization of the moment functions. Using β_1 to indicate a vector of technology parameters and as discussed above, the production technology can be represented by $Q = Q(x, C, e, \beta_1)$. The production function is stochastic given the random error term *e*. Let the stochastic output *Q* have a cumulative distribution *F(e)*, then the first and the *i*th central moments of output *Q* can be represented, respectively, as

(2) $m_1(\mathbf{x}, C, \boldsymbol{\beta}_1) = E[Q(\mathbf{x}, C, \boldsymbol{e}, \boldsymbol{\beta}_1)] = \int Q(\mathbf{x}, C, \boldsymbol{e}, \boldsymbol{\beta}_1) dF(\boldsymbol{e}),$

(3)
$$m_i(\mathbf{x}, C, \boldsymbol{\beta}_i) = E[\{Q(\mathbf{x}, C, e, \boldsymbol{\beta}_1) - m_1(\mathbf{x}, C, \boldsymbol{\beta}_1)\}^i]$$

 $= \int (Q(\mathbf{x}, C, \mathbf{e}, \boldsymbol{\beta}_1) - m_1(\mathbf{x}, S, \boldsymbol{\beta}_1))^i dF(\mathbf{e}) \text{ for } i \ge 2.$

Here, *E* is the expectation operator; m_1 is the mean and m_i is the *i*th moment of output (for $i \ge 2$); and β_i is a vector of parameters. The models in (1) and

(2) have the advantage of being flexible as there are no restrictions within or cross moments.

By rewriting equations (2) and (3), we get the following equations

(4)
$$\varepsilon_1 = Q(\mathbf{x}, C, \boldsymbol{\beta}_1) - m_1(\mathbf{x}, C, \boldsymbol{\beta}_1),$$

(5)
$$(\boldsymbol{\varepsilon}_1)^i = m_i(\boldsymbol{x}, \boldsymbol{C}, \boldsymbol{\beta}_i) + \boldsymbol{\varepsilon}_i, i \ge 2.$$

Here, $E(\varepsilon_i)$ and $E(\varepsilon_i \varepsilon_{i'}) = 0$ $(j = 1, 2, ..., n \text{ and } j \neq j')$.

As discussed by Kendall and Stuart (1977) and shown in many empirical analyses (e.g., Day 1965; Di Falco and Chavas 2009; Di Falco and Veronesi 2014; Tack, Harri, and Coble 2012; and Anderson, Dillon, and Hardaker 1980), the first three moments including location (mean), dispersion (variance), and skewness (the third moment) of a given distribution can adequately approximate the distribution. We, therefore, choose the first three moments to represent the distribution of yield in our analysis. While the variance (m_2) is a traditional measure of risk, the skewness of the output measure (m_3) captures the tail asymmetry of a yield distribution around its mean. A negative (positive) skewness implies a distribution skewed to the left (right). A lower skewness presents a greater exposure to the downside risk of unexpected low yield, i.e., crop failure.

From equations (4) and (5), we have the mean, variance, and skewness of yield as the following,

(6)
$$Q = m_1(\mathbf{x}, C, \boldsymbol{\beta}_1) + \boldsymbol{\varepsilon}_1,$$

(7)
$$(\boldsymbol{\varepsilon}_1)^2 = m_2(\boldsymbol{x}, \boldsymbol{C}, \boldsymbol{\beta}_i) + \boldsymbol{\varepsilon}_2,$$

(8)
$$(\boldsymbol{\varepsilon}_1)^3 = m_3(\boldsymbol{x}, \boldsymbol{C}, \boldsymbol{\beta}_i) + \boldsymbol{\varepsilon}_3.$$

Here again, $E(\varepsilon_j) = 0$ and $E(\varepsilon_j, \varepsilon_{j'}) = 0$ (j = 1, 2, 3 and $j \neq j'$). Empirically, if β_1^* is a consistent estimator of β_1 from a sample of observed outputs, then $\varepsilon_1^* = Q(\mathbf{x}, C, \beta_1) - m_1(\mathbf{x}, C, \beta_1^*)$ is a consistent estimator of $\varepsilon_1 = Q(\mathbf{x}, C, \beta_1) - m_1(\mathbf{x}, C, \beta_1)$. It also suggests that $(\varepsilon_1^*)^i = m_i(\mathbf{x}, S, \beta_i^*) + \varepsilon_i$, $i \ge 2$ is a consistent estimator of $(\varepsilon_1)^i$. The models in (5), (6), and (7) have no restrictions within or cross moments and thus are flexible.

The adoption of cover cropping *C* is variance increasing, variance neutral, or variance decreasing if $(\partial m_2 / \partial_C) > 0$, = 0, or < 0, respectively. For a risk averse farmer, $(\partial m_2 / \partial_C) > 0$ meaning that the adoption of cover crops creates a greater risk in output is undesirable. Similarly, the adoption of cover cropping *C* is skewness increasing, skewness neutral, or skewness decreasing if $(\partial m_3 / \partial_C) > 0$, = 0, or < 0, respectively. And, for a risk-averse farmer, $(\partial m_3 / \partial_C) < 0$ meaning that the adoption of cover crops increasing the exposure to a lower output is undesirable.

Some factors that are known by farmers but unknown to economists may affect both yield and the cover cropping decision. Consequently, when empirically estimating the yield equation as shown in (6), what arises is a concern that the adoption of cover crops may be endogenous. The endogeneity may result in inconsistent and biased estimates. To address the potential endogeneity issue, we use a two-stage method, which is one of the most potent and versatile tools available to treat endogeneity (Antonakis et al. 2014).

We will first estimate the use of cover crops with an instrumental variable approach. We model the adoption of cover crops in a logit model as follows:

Cover cropping: Cover cropping: $y_{cc}^* = \alpha' x + \gamma Z + \varepsilon$

(9)
$$y_{cc} = \begin{cases} 1 \text{ if } y_{cc}^* > 0\\ 0 \text{ otherwise} \end{cases}$$

Here y_{cc}^* is a latent continuous variable associated with the adoption of cover cropping; y_{cc} is the corresponding observed binary outcome with a value of 1 if cover cropping is adopted and 0 otherwise; α' is a transposed vector for parameters to be estimated; and Z is a vector of instrumental variables.

7.3 Data and Statistics

We apply our analysis to US soybean production. We construct the data from the USDA's 2018 ARMS Phase II, Soybean Production Practices and Costs Report, and Phase III Soybean Costs and Returns Report. The Phase II survey covers a cross-section of soybean fields in 19 states and collects information on production and management practices, input uses, and field characteristics. The Phase III report provides information on farm operators and financial characteristics. Farm-level survey data provide us a good opportunity to look more closely at farm activities and the motives behind them (Dong, Hennessy, and Jensen 2010; Dong, Hennessy, Jensen, and Volpe 2016).

As conventional and organic production are significantly different in production practices, we only use data from conventional soybean growers. We delete all observations with missing values, leaving a total of 1,177 observations. ARMS has a complex survey design and is a probability-based survey with unequal probability sampling (National Research Council 2007). To account for the survey design, we use the sampling weights (expansion factors) provided by USDA NASS to expand the sample to generate population estimates in the statistical analysis. With the survey weights applied to the sample observations, the weighted sample represents approximately 835,530 soybean fields in the United States.

7.3.1 Variables in Moment Equations

The dependent variables in equations (6) to (8) are the first three moments of the distribution of soybean yield per acre, respectively. Variable *phos*-

phorus measuring the use of phosphorus per acre is included along with its quadratic term. They are expected to increase the yield. Other independent variables include those on production practices, field characteristics, regional location, and weather/climate. To capture regional differences, indicator variables were constructed based on Farm Resource Regions (USDA ERS 2000), which are defined based on farm, soil, and climate characteristics rather than state boundaries. The regional dummy variables (Eastern Uplands, Heartland, Mississippi Portal, Northern Crescent, Northern Great Plains, Prairie Gateway, and Southern Seaboard) are equal to 1 if the field locates in the corresponding region and 0 otherwise. Two variables (PlantLate and ReplantPct) are used to capture the impacts of adverse factors negatively affecting soybean yields at the start of the planting season. Variable PlantLate is equal to 1 if the planting date fell in the last 15 percent percentile of the state, and 0 otherwise. Variable Replant Pct is the proportion of fields that was replanted. Both might be resulted from adverse weather and result in shorter growing seasons for soybeans, and thus are expected to have a negative effect on yield. Dummy variable manure has a value of 1 if manure was applied to the field and 0 otherwise. Several field characteristics may affect yields such as soil texture and slope of the field (Butcher et al. 2018; Arora et al. 2011; Shane and Barker 1986; Kaspar et al. 2004; Jiang and Thelen 2004; Kravchenko, Bullock, and Boast 2000; Linkemer, Board, and Musgrave 1998; Nelson and Meinhardt 2011) are also included in the yield moment functions. Soil texture is categorized into five types: loam, clay, sandy, mixed, and silty. Slope is categorized into two levels: nearly level and moderate/steep grade (even or variable). While the soybean growing season is from May to September, weather in both July and August is important for soybean yields (Westcott and Jewison 2013). The county-level US Drought Monitor (USDM) indicator jointly produced by the National Drought Mitigation Center (NDMC), the National Oceanic and Atmospheric Administration (NOAA), and the US Department of Agriculture is used. The USDM indicator has five categories: D0-D4, of which D0 indicates abnormally dry but not in drought, while D1-D4 indicates moderate drought to exceptional drought. The USDM indicator is based on inputs including the Palmer Drought Severity Index (PDSI), the Standardized Precipitation Index (SPI), satellite-based assessments of vegetation health, and various indicators of soil moisture as well as hydrologic data (NDMC 2021). If on a weekly average over 10 percent of a county area is categorized as D1 or above in either July or August of 2018, then an indicator variable *drought18* is equal to 1 and 0 otherwise. Moreover, modified growing degree days (mGDD) and overheating growing days (ODD) for soybeans during July and August of 2018 are also included to explicitly capture temperature or heat effects on the growth and development of soybean plants. GDD is one of the most important

| Variable and unit | Survey Mean | Std. Err |
|--------------------------------|-------------|----------|
| Yield (bushels/acre) | 52.354 | 0.097 |
| Heartland (0/1) | 0.498 | 0.003 |
| Northern Crescent (0/1) | 0.141 | 0.003 |
| Northern Great Plains (0/1) | 0.066 | 0.001 |
| Prairie Gateway (0/1) | 0.054 | 0.001 |
| Eastern Uplands (0/1) | 0.053 | 0.002 |
| Southern Seaboard (0/1) | 0.105 | 0.002 |
| Mississippi Portal (0/1) | 0.083 | 0.001 |
| PlantLate (0/1) | 0.129 | 0.002 |
| Replant Pct | 0.041 | 0.002 |
| Manure (0/1) | 0.043 | 0.001 |
| Phosphorus (lbs/acre) | 25.550 | 0.229 |
| drought18 (0/1) | 0.175 | 0.002 |
| Soil texture: loam (0/1) | 0.369 | 0.003 |
| Soil texture: clay (0/1) | 0.170 | 0.003 |
| Soil texture: sandy (0/1) | 0.069 | 0.002 |
| Soil texture: mixed (0/1) | 0.369 | 0.005 |
| Soil texture: silty (0/1) | 0.019 | 0.001 |
| Growing degree days (Celsius) | 3304.675 | 4.396 |
| Overheating degree days (days) | 30.778 | 0.133 |
| Cover crops | 0.100 | 0.003 |

Table 7.1 Survey means and standard errors of variables in moment functions

factors influencing the rate of development in soybean (Major et al. 1975; Pedersen and Licht 2014; Kessler, Archontoulis and Lich 2020). Daily mGDD is calculated as

$$mGDD = \max\left(\frac{\min(daily \max temp, higher development threshold)}{+\max(daily \min temp, lower development thresholds)} - 50, 0\right).$$

GDD depends upon the minimum and maximum temperatures which affect the plant's growth. The higher and lower development thresholds are 86°F and 50°F, respectively, for soybeans. Daily mGDD in July and August is accumulated to get mGDD for the two months. ODD is the count of days in July and August with a temperature over 89.6°F and measures the heat stress for crops. Cover cropping is included in the function as a dummy variable. It is set equal to 1 if the cover cropping was adopted and 0 otherwise. In addition, interactions between cover cropping and drought status *drought18* are also included to capture the effects of the adaptation strategy on yield moments conditional on drought events. Survey population and sample summary statistics for variables used in the moment functions are reported in table 7.1.

7.3.2 Variables in Logit Model

The instrumental variables used in the cover crop equation include farmers' concerns about soil and water-related issues. Seven dummy variables are constructed, indicating concerns on water-driven erosion, wind-driven erosion, soil compaction, poor drainage, low organic matter, water quality, and other concerns, respectively, and taking a value of 1 if a farmer had such a concern and 0 otherwise. We believe that farmers having concerns about soil erosion or soil quality may have more intention to adopt soil conservation practices. The variable of land ownership is included as another instrumental variable. The variable takes a value of 1 if the operator owned the land and 0 otherwise. We expect that land ownership may increase the likelihood of fields adopting cover crops as landowners may care more about soil erosion and soil quality on their own land and thus have more motivation to adopt conservation practices.

Farm size and some other field characteristics such as field size and whether any part of the field was classified as "highly erodible," and whether the field contained a wetland are also included in the cover crop equation given their possible effect on farmers' cover crop decision making (Ding, Schoengold, and Tadesse 2009; Vitale et al. 2011; Wandel and Smithers 2000). Highly erodible land is any land that can erode at an excessive rate due to its soil properties. Farmers are required to farm such land in accordance with a conservation plan or system approved by NRCS (USDA Risk Management Agency 2015). We expect that field classified as "highly erodible" is more likely to adopt cover crops for their vulnerability to soil erosion. Both farm and field sizes are measured in acres. We expect that larger farms and larger fields may be less likely to adopt cover crops given the time and labor requirements of cover crop implementation and management. Moreover, farmers' age, education, years of experience in farming, and off-farm work are also included. We expect that older farmers or farmers with off-farm work may be less likely to adopt cover crops given the time and labor investment needed for cover crop implementation and management.

Several studies have found evidence that extreme weather affects farmers' adoption of practices. We use the variable *drought5yr* to indicate the number of years in which a D1 degree or above drought happened in at least 10 percent of the county areas during July and August in the last five years. Survey population and sample summary statistics for variables used in the logit model are reported in table 7.2. To avoid forbidden regression, all exogenous variables in yield moment equations are also included in the estimation of the cover crop equation. However, since many of them do not have realistic meanings (e.g., phosphorus use should not affect cover crop adoption), we only report several of their estimates, including regional dummy variables, soil texture, and slope of the field in the next section.

| Variable and Unit | Survey Mean | Std.Err |
|-----------------------------------------------|-------------|---------|
| Cover crops (0/1) | 0.100 | 0.003 |
| Heartland (0/1) | 0.498 | 0.003 |
| Northern Great Plains (0/1) | 0.141 | 0.003 |
| Prairie Gateway (0/1) | 0.066 | 0.001 |
| Eastern Uplands (0/1) | 0.054 | 0.001 |
| Southern Seaboard (0/1) | 0.053 | 0.002 |
| Mississippi Portal (0/1) | 0.105 | 0.002 |
| Northern Crescent (0/1) | 0.083 | 0.001 |
| Fields having moderate or steeper slope (0/1) | 0.556 | 0.004 |
| Soil texture: loam (0/1) | 0.369 | 0.003 |
| Soil texture: clay (0/1) | 0.170 | 0.003 |
| Soil texture: sandy (0/1) | 0.069 | 0.002 |
| Soil texture: mixed (0/1) | 0.369 | 0.005 |
| Soil texture: silty (0/1) | 0.019 | 0.001 |
| Concern about water-driven erosion (0/1) | 0.262 | 0.003 |
| Concern about wind-driven erosion (0/1) | 0.080 | 0.002 |
| Concern about soil compaction (0/1) | 0.261 | 0.004 |
| Concern about poor drainage (0/1) | 0.232 | 0.004 |
| Concern about low organic matter (0/1) | 0.109 | 0.003 |
| Concern about water quality (0/1) | 0.066 | 0.002 |
| Other concerns (0/1) | 0.025 | 0.001 |
| Field classified as "highly erodible" (0/1) | 0.180 | 0.003 |
| Field contains wetland (0/1) | 0.041 | 0.002 |
| Age (years) | 57.730 | 0.100 |
| Years of experience (years) | 33.326 | 0.121 |
| College education (0/1) | 0.251 | 0.002 |
| Off-farm work (0/1) | 0.185 | 0.004 |
| Land ownership (0/1) | 0.491 | 0.004 |
| Farm size (acres) | 1348.206 | 11.558 |
| drought5yr (years) | 0.764 | 0.007 |
| Field size (acres) | 50.015 | 0.328 |

Table 7.2 Survey means and standard errors of variables in the logit model

7.4 Estimation Results

Utilizing a two-stage method, we estimate the logit model specified in equation (9) in the first stage to address the endogeneity issue. We explore the determinants of cover crop adoption with a focus on the effects of climate change, farmers' demographic information, and field characteristics. In the second stage, we estimate the moment equations of (6), (7), and (8).

7.4.1 Results of Logit Model for Cover Crop Adoption

The logit model estimation results are reported in table 7.3. The results show that there existed regional differences in the adoption of cover cropping. Compared to fields in Heartland, those in Northern Crescent, Northern Great Plains, Prairie Gateway, and Mississippi Portal were less likely

| Parameter | Estimation | Bootstrapped Std. Err. |
|------------------------------------|---------------|------------------------|
| Northern Crescent | -1.049*** | 0.157 |
| Northern Great Plains | -3.100*** | 0.212 |
| Prairie Gateway | -0.605*** | 0.205 |
| Eastern Uplands | 2.363*** | 0.129 |
| Southern Seaboard | 2.788*** | 0.172 |
| Mississippi Portal | -1.187*** | 0.202 |
| slope | 0.520*** | 0.057 |
| Soil texture: clay | 0.711*** | 0.114 |
| Soil texture: sandy | -0.249* | 0.144 |
| Soil texture: mixed | 0.413*** | 0.087 |
| Soil texture: silty | 0.330 | 0.277 |
| Concern about water-driven erosion | -0.413*** | 0.121 |
| Concern about wind-driven erosion | 0.532*** | 0.102 |
| Concern about soil compaction | 0.223* | 0.129 |
| Concern about poor drainage | -0.218** | 0.105 |
| Concern about low organic matter | -0.793*** | 0.121 |
| Concern about water quality | 0.901*** | 0.112 |
| Other concerns | 0.976*** | 0.142 |
| Field "highly erodible" | 0.545*** | 0.120 |
| Field contains wetland | 0.355 | 0.531 |
| age | -0.024*** | 0.006 |
| Years of experience | -0.005 | 0.005 |
| College education | 0.551*** | 0.078 |
| Off-farm work | -0.258** | 0.109 |
| Land ownership | 0.230* | 0.080 |
| Farm size | -9.280E-05*** | 3.160E-05 |
| drought5yr | -0.089 | 0.068 |
| Field size | -0.003*** | 0.001 |
| Constant | -3.119*** | 0.667 |

Table 7.3 Estimates of parameters of the logit model

Note: Statistical significance at 1%, 5%, and 10% are denoted as ***, **, and *, respectively.

to adopt cover crops while those in Eastern Uplands and Southern Seaboard were more likely to do so. The higher adoption rate of cover crops in Heartland, Eastern Uplands, and Southern Seaboard comparting to that in other regions can be attributed to several factors. In addition to federal programs such as the Environmental Quality Incentives Program (EQIP) and Conservation Stewardship Program (CSP), many states in Heartland, Eastern Uplands, and Southern Seaboard have implemented state incentive programs, which have been found positively correlated with the adoption of cover crops (e.g., Fleming 2017; Lichtenberg, Wang, and Newburn 2018; Wallander et al. 2021). The top seven state-funded cover-crop programs in terms of acreage in the US are all in the three regions (Wallander et al. 2021). In addition, tax credits, reduction on crop insurance premiums, and programs that rent out or loan equipment related to cover cropping in the three regions (Wallander et al. 2021) may also contribute to their higher adoption rates. Moreover, access to technical assistance and extension service may also cause variations in cover crop adoption across regions. Regional differences in the adoption of cover crops or other conservation practices have been found in other studies. Unger and Vigil (1998) find that the decision on adopting cover crops may vary from semiarid regions to humid/sub-humid regions depending on whether water is scarce or not. Given that western regions had a lower adoption rate than Heartland, Eastern Uplands, and Southern Seaboard, concerns about cover crops depleting soil water for the following main crops probably dominated in those drier western regions. In addition, as suggested in some other studies (e.g., Davey and Furtan 2008, Ding, Schoengold, and Tadesse 2009, and Claassen et al. 2018), a higher rate of adoption in Eastern Uplands and Southern Seaboard may reflect the concern about soil erosion given more rainfall in the regions.

The likelihood of cover crop adoption was also affected by farmers' concerns. Farmers who had concerns over wind-driven erosion, soil compaction, water quality, or other concerns were more likely to adopt cover crops than those who did not have such concerns. The result aligns itself with the benefits that cover crops are supposed to provide. In contrast, farmers who had concerns over water-driven erosion, poor drainage, or low organic matter were less likely to adopt than those without such concerns. Steele, Coale, and Hill (2012) did not observe consistent differences in total organic matter and labile organic matter between the winter cover crop and control soils in an experiment with 13 years of cover crop use. Our finding on the effect of concerns over low organic matters is consistent with Steele, Coale, and Hill's (2012) finding. While plenty of studies have found a positive effect of cover crops on soil organic matters and erosion by water (e.g., Shanks, Moore, and Sanders 1998; Ding et al. 2006; Dube, Chiduza, and Muchaonyerwa 2012), many studies have concluded that cover crops' effect on organic matter may vary with cover crop species, soil type, and other practices, such as tillage and rotation (e.g., Wulanningtyas et al. 2021; Abdollahi and Munkholm 2014; Dube, Chiduza, and Muchaonyerwa 2012; and Motta et al. 2007). Our results imply that cover crops might either have not practically worked well on improving water-driven erosion, poor drainage and organic matter conditional on commonly used management and practices in soybean production, or complexities of management and implementation of cover crops as well as their interactions with other practices have discouraged its adoption to address those concerns.

Field characteristics also affected the adoption of cover crops. As expected, if a field was classified as "highly erodible," it was more likely to adopt cover crops; fields with slopes were also more likely to adopt cover crops since they are more vulnerable to soil erosion. Whether a field contained wetland did not statistically significantly affect the adoption. Soil texture also affected farmers' decisions on cover crop adoption. Compared to loam soil, fields

with clay or mixed soil were more likely to adopt cover crops and fields with sandy soil were less likely to do so. There was, however, no statistically significant difference between loam soil and silty soil in cover crop adoption. Larger farms and larger fields were less likely to adopt cover crops. This is expected since the larger the farm/field, the more labor and time are needed for the implementation and management of cover crops.

Farmers' demographic characteristics played an important role in practice adoption decisions. Older farmers or farmers working at least 50 percent off farms were less likely to adopt cover crops. This might be due to the labor and time requirements of cover crop implementation and management, as discussed previously. In addition, more educated farmers were more likely to adopt cover crops as they might better understand the importance of cover crops in the environment and agricultural sustainability. Consistent with expectation, farmers who had land ownership were more likely to adopt cover crops as they care more about their own land.

Drought in the last five years did not affect the likelihood of cover crop adoption. This might be due to the same reason as discussed above. Cover crops' drought mitigating effect may interact with other factors such as cover crop species, planting and termination time, tillage, rotation, soil type, etc. Proper combinations of the use of cover crops and other practices conditional on soil and weather conditions are required and many farmers might not have observed desired results of the use of cover crops as a drought adaptation strategy.

7.4.2 Results of Yield Moment Functions

Results of the first moment yield equation are presented in table 7.4. As expected, more mGDD increased the soybean yield while more ODD decreased the soybean yield. If the year had a drought in July or August, the mean yield decreased. Regional differences were shown in soybean yield. Fields in Heartland, which includes Illinois, Indiana, Iowa, and parts of Missouri, Nebraska, South Dakota, Minnesota, Ohio, and Kentucky, had the highest yield among all regions. Unsurprisingly, if soybeans were planted late or a bigger proportion of fields were replanted, the field had a lower vield, resulting from shortened vegetative and reproductive intervals. Inputs of fertilizer did help increase the yield. The more phosphorus was applied, the higher the yield. With the small parameter for the quadratic term, the effect of phosphorus on yield was close to linear. The application of manure also helped improve yield by adding more nutrients to the soil. Without cover crops, soybean yield in loam soil was higher than in clay, sandy, or mixed soil, but not significantly different from that in silty soil. This is consistent with the findings of previous studies such as Radočaj et al. (2020) and He, Luo, and Sun (2014).

Regional differences also showed in yield variance and skewness. Compared to Heartland, soybean yield in Southern Seaboard had lower varia-

| | First moment (M | ean) equation | Second mome equa | nt (variance) tion | Third moment (ske | wness) equation |
|--------------------------------|-------------------------|--------------------------|----------------------|--------------------------|--------------------|--------------------------|
| Parameter | Estimation | Bootstrapped Std. Err | Estimation | Bootstrapped Std. Err | Estimation | Bootstrapped Std. Err |
| Cover crops | -3.890 | 2.720 | 48.337*** | 9.092 | 2868.821*** | 341.341 |
| Drought × cover crops | -20.160^{***} | 4.452 | 223.235*** | 21.604 | 16562.100^{***} | 1050.960 |
| Cover crops×clay soil | 6.869*** | 2.517 | 4.441 | 32.237 | -12716.120^{***} | 1430.358 |
| Cover crops×sandy soil | 17.406^{***} | 2.509 | -79.215^{***} | 17.073 | -6299.486*** | 559.366 |
| Cover crops×mixed soil | 18.586*** | 1.655 | -48.293*** | 18.088 | -8944.720*** | 667.661 |
| Cover crops×silty soil | 14.110^{***} | 1.926 | -154.608^{***} | 23.123 | -4624.771 *** | 793.497 |
| Northern Crescent | -2.994*** | 0.314 | -3.153 | 2.946 | -263.873^{**} | 110.773 |
| Northern Great Plains | -9.363^{***} | 0.472 | 19.329*** | 4.805 | 377.511*** | 146.329 |
| Prairie Gateway | -4.436*** | 0.603 | 103.903*** | 7.312 | -1081.599*** | 259.667 |
| Eastern Uplands | -17.638^{***} | 1.147 | 39.311*** | 6.498 | 1982.181*** | 286.687 |
| Southern Seaboard | -25.983*** | 0.819 | -20.231^{***} | 4.326 | 932.423*** | 131.663 |
| Mississippi Portal | -4.093*** | 0.668 | 153.902*** | 12.949 | -2364.662*** | 500.023 |
| PlantLate | -4.934*** | 0.256 | -20.423^{***} | -20.423*** | -461.056^{***} | 77.673 |
| ReplantPct | -6.639*** | 1.069 | 42.540*** | 11.625 | 4547.295*** | 499.821 |
| Manure | 1.956^{***} | 0.440 | 3.777 | 4.742 | 547.647*** | 184.636 |
| Phosphorus | 0.020^{***} | 0.004 | -0.569*** | 0.062 | -3.022 | 1.899 |
| Phosphorus ² | 1.735E-04*** | 2.680E-05 | 0.003^{***} | 0.001 | 0.003 | 0.012 |
| Drought 18 | -3.696*** | 0.401 | 30.540*** | 4.323 | -856.462*** | 151.164 |
| Soil texture: clay | -1.507^{***} | 0.389 | 7.641** | 3.275 | -788.829*** | 113.448 |
| Soil texture: sandy | -10.515^{***} | 0.477 | 71.824*** | 5.500 | 2274.254*** | 211.552 |
| Soil texture: mixed | -3.742*** | 0.243 | -17.207^{***} | 2.521 | 446.028*** | 84.746 |
| Soil texture: silty | 0.485 | 0.429 | 1.400 | 10.029 | -1525.396^{***} | 389.240 |
| mGDD | 0.013^{***} | 0.001 | 0.041*** | 0.007 | -0.314 | 0.242 |
| ODD | -0.184^{***} | 0.016 | -0.650^{***} | 0.189 | -5.393 | 6.670 |
| Constant | 23.424*** | 1.390 | 7.472 | 18.158 | 1100.763* | 648.716 |
| R^2 | 0.334 | | 0.095 | | 0.112 | |
| Note: Statistical significance | e at 1%, 5%, and 10% ar | e denoted as ***, **, | and *, respectively. | | | |

Estimates of parameters of yield moment equations

Table 7.4

tions, and it had higher variations in the Northern Great Plains, Prairie Gateway, Eastern Uplands, and Mississippi Portal. In addition, Northern Crescent, Prairie Gateway, and Mississippi Portal had higher downside risk; in contrast, Northern Great Plains, Eastern Upland, and Southern Seaboard had lower downside risk. The higher percentage of field replanted, the more variation and the lower downside risk in yield, probably resulting from the replacement of damaged plants, for example by frosts. If soybeans were planted late, their yield varied less but had a higher downside risk, probably due to a shorter growing season or higher probability of frost before harvest. At the mean application level, phosphorus inputs reduced the variation in vield, but manure application did not. In addition, more mGDD increased the variance while more ODD decreased the variance. Both did not change the downside risk. As expected, a drought that occurred in July or August increased the variance of yield and in the meantime increased the risk of crop failure. Yield moments also showed a heterogeneous effect on soil texture. Compared to loam soil, clay and sandy soils had higher variance; clay and silty soils had lower skewness or higher downside risks.

Given the interaction terms with soil texture and weather, marginal effects of cover crops were calculated and presented in table 7.5. Standard errors are calculated using the delta method. As shown in table 7.5, if there was no drought in July and August that year, then cover crops statistically significantly increased the yield of soybeans planted in sandy, silty, or mixed soil; but there was no statistically significant effect on soybean yield in loam and clay soils. This is consistent with many studies and experiments finding that cover crops help increase cash crop yield. In the meantime, cover crops increased yield variance in loam and clay soils, but decreased yield variance in sandy and silty soils and had no significant effect in mixed soil. In addition, planting cover crops reduced downside risk in loam soil while increased downside risk in all other types of soils. From the above, we can see that the effects of cover crops depend on soil types when there was no drought. And there was always a trade-off between the mean, the variation, and the downside risk of yield, i.e., there was no simultaneous positive effect of cover crops on the three moments of yield, which affected farmers' expected utility.

With droughts in July and August, cover crops reduced soybean yield in all soil types, although the effects in sandy and mixed soil were not statistically significant. This implies that cover crops consumed water in the soil for their own growth and reduced water available for the following cash crops. When drought occurred, the water supply worsened to a point where crop yield decreased. In addition, cover crops increased yield variance but reduced the risk of crop failure in all types of soils when drought occurred. The mixture of positive and negative effects of cover crops on yield moments is somewhat consistent with the finding in the first stage that previous droughts did not affect farmers' adoption of cover crops, implying a divided acceptance of cover crops as a drought adaptation strategy among farmers.

| | | | yield equ | ation | second moment equation | (variance) | third moment (equatio | skewness) n |
|-------------|-----------|---------|-----------------|----------|---------------------------|------------|---------------------------|----------------|
| | Soil type | Drought | Estimation | Std. Err | Estimation | Std. Err | Estimation | Std. Err |
| | loam | No | -3.890 | 2.720 | 48.337*** | 9.092 | 2868.821*** | 341.341 |
| | | Yes | -24.049*** | 4.579 | 271.572*** | 26.449 | 19430.921*** | 1244.251 |
| | Clay | No | 2.980 | 3.108 | 52.778* | 31.177 | -9847.299*** | 1426.819 |
| | | Yes | -17.180^{***} | 3.826 | 276.013*** | 31.774 | 6714.801*** | 1166.486 |
| cover crops | sandy | No | 13.516*** | 2.349 | -30.878** | 13.859 | -3430.665^{***} | 391.965 |
| | | Yes | -6.643 | 4.586 | 192.357*** | 23.436 | 13131.435*** | 1080.542 |
| | mixed | No | 14.696*** | 2.375 | 0.045 | 13.002 | -6075.899*** | 461.852 |
| | | Yes | -5.463 | 4.216 | 223.280*** | 17.548 | 10486.201*** | 701.112 |
| | silty | No | 10.220^{***} | 1.839 | -106.271*** | 21.328 | -1755.950*** | 709.924 |
| | | Yes | -9.940*** | 4.234 | 116.964*** | 26.404 | 14806.150*** | 981.327 |

| conditions |
|------------|
| different |
| ps under |
| COVER CF0 |
| effects of |
| Marginal o |
| |
| Table 7.5 |

Note: Statistical significance at 1%, 5%, and 10% are denoted as ***, **, and *, respectively.

7.5 Conclusions and Discussions

We explored factors affecting farmers' adoption of cover crops by a logit model and examined the effects of cover crops on soybean yield and its risk by three moment functions. By incorporating two interplays between cover crops and soil type, and cover crops and drought, we were able to explore the varying effects of cover crops in drought and different soil types. While we found that the adoption of cover crops varied in regions and soil types and was affected by field properties and farmers' demographic characteristics and concerns, we did not find a significant effect of previous droughts on the adoption. The results from the moment functions of soybean yield confirmed what the results in the first stage suggest. When there was a drought, cover crops reduced yield and increased yield variance. However, cover crops also reduced the downside risk of crop failure in the meantime. The mixed effects of cover crops on yield and its risk associated with an occurrence of drought support the statistically insignificant effect of the previous drought on cover crop adoption, implying that farmers were divided in the acceptance of cover crops as a means to build resilience to drought. The mixed effect of cover crops also warrants a further study to calculate the certainty equivalent of net economic return of soybeans with cover crops, which requires information on cover crop seed, planting, and termination cost as well as additional or saved fertilizer and pesticide costs. The certainty equivalent of net economic returns of soybeans with cover crops may provide more information on economic impediments to farmer adoption.

The low adoption rate of cover crops may also be related to complex interactions between management and cultural practices including species selection, planting and termination date, rotation, and termination method. Achieving desired benefits requires significant training, learning, and adjustments in many aspects of the farming system (Wallander et al. 2021). As shown in the National Cover Crop Survey 2020 (Conservation Technology Information Center 2020), roughly 70 percent of respondents said that they typically used their own experience of trial and error for cover cropping. About 67 percent and 60 percent of the respondents considered the two approaches, i.e., local farm tours to see how cover crops worked and one-on-one technical assistance to select, plant, or manage cover crops, very helpful or moderately helpful, respectively, in encouraging them to try cover cropping. Therefore, programs that provide necessary training and show-case cover crop management to farmers could address an important lack of information.

Greater soil and environmental benefits can be achieved when cover crops are utilized in conjunction with other practices (Wallander et al. 2021), such as conservative tillage, irrigation, crop rotation, nutrient management, and adoption of drought tolerant seeds—which are currently available for maize. A broader range of research that finds proper combinations of cover crops and other practices conditional on soil types and weather are crucial for establishing practice guidance for farmers. Such guidance can help farmers achieve desired results by using cover crops as a drought adaptation strategy as well as a tool for improving soil and environmental benefits along with a suite of other conservation practices.

The recent increase in cover crop adoption has been accompanied by financial incentives. Given financial support, cover crop acres enrolled in the Environmental Quality Incentives Program (EQIP) increased from 312.6 thousand acres in 2009 to 2,443.1 thousand acres in 2020 (USDA Climate Hubs 2021). In 2018, about one-third of the acreage planted with a cover crop received a financial assistance payment for cover crop adoption from either federal, state, or other programs, ranging from \$12 per acre to \$92 per acre (Wallander et al. 2021). The USDA NRCS recently announced a program to promote the use of soil health practices, especially cover crops. The initiative sets a goal of doubling the number of corn and soybean acres using cover crops to 30 million acres by 2030 (USDA 2022). Given the mixed effects of cover crops on soybean yield and yield risks found in this study, financial incentives can help improve the certainty equivalent of net returns and thus encourage more risk-averse farmers to adopt cover crops. In addition, farmers have recently been paid to plant cover crops by large seeds, chemical, and food companies to generate carbon credits to offset their environmental footprints (Reuters 2022). The payments, however, are generally not as much as those from EQIP and CSP. In addition, the current carbon credit market lacks transparency and liquidity (Ag Decision Maker 2021). It is facing several challenges including setting up protocols to ensure the additionality and permanence of net greenhouse gas (GHG) reductions (Blaustein-Rejto 2021).

We conclude the paper by recognizing a key limitation of this study: it does not consider the long-term effects of sustainable practice adoption by using cross-sectional data. Cover crops have multiple benefits to soil and the environment. Cover crops can be used not only as a drought adaptation strategy but also to reduce soil erosion, enhance weed control, improve soil health, increase carbon storage, improve water quality through reduced nutrient and sediment runoff, and increase biological diversity. While solely comparing the cost of seed, seeding, and management to the impact on the yield of the following main crop may show a loss in the first few years, cover crops may possibly improve the efficiency and resiliency of the entire farm over time, resulting in a net benefit from the broad, holistic standpoint (Myers, Weber, and Tellatin, 2019). Myers, Weber, and Tellatin (2019), for example, show that the adoption of cover crops may have negative net returns in the first year, negligible net returns in three years, but about \$18 net returns in five years. In addition, if cover crops are used to address more than one yield-limiting factor in a field such as for grazing, improving soil health, and weed impression, then the net return can be larger and faster. This can be applied to many other sustainable practices. For example, the payoffs from investments in improving soil fertility and reducing soil erosion are cumulative and may take several years. And the subsequent improvement in soil fertility and reduction in soil erosion can reduce future expenses for crop nutrients, irrigation, and energy (Lee 2005; Tilman et al. 2001). If such long-term positive net returns can be demonstrated by more farmers who are supported financial assistance from federal and state programs to offset a portion of upfront investments—which have been proved very useful in increasing the adoption (Bowman and Lynch 2019)—then the adoption of cover crops, as well as other drought adaptation strategies, may surge.

References

- Abdollahi, L., and L. Munkholm 2014. "Tillage System and Cover Crop Effects on Soil Quality: I. Chemical, Mechanical, and Biological Properties." Soil Science Society of America Journal 78 (1): 262–70.
- Acharya, B., S. Dodla, L. Gaston, M. Darapuneni, J. Wang, S. Sepat, and H. Bohara. 2019. "Winter Cover Crops Effect on Soil Moisture and Soybean Growth and Yield under Different Tillage Systems." *Soil and Tillage Research* 195: 104430.
- Acuña, J. C., and M. Villamil. 2014. "Short-Term Effects of Cover Crops and Compaction on Soil Properties And Soybean Production in Illinois. Agronomy Journal 106 (3): 860–70.
- Ag Decision Maker. 2021. How to Grow and Sell Carbon Credits in US Agriculture. Accessed June 12, 2022. https://www.extension.iastate.edu/agdm/crops/pdf/a1-76.pdf.
- Anderson, J. R., J. D. Dillion, and B. Hardaker. 1980. *Agricultural Decision Analysis*. Ames, Iowa: The Iowa State University Press.
- Andraski, T. W., and L. G. Bundy. 2005. "Cover Crop Effects on Corn Yield Response to Nitrogen on an Irrigated Sandy Soil." Agronomy Journal 97 (4): 1239–1244.
- Antle, J. 1983. "Testing the Stochastic Structure of Production: A Flexible Moment-Based Approach." Journal of Business & Economic Statistics 1 (3): 192–201.
- Antonakis, J., S. Bendahan, P. Jacquart, and R. Lalive. 2014. "Causality and Endogeneity: Problems and Solutions." In *The Oxford Handbook of Leadership and Organizations*, edited by D. V. Day, 93–117. New York: Oxford University Press.
- Arbuckle, J. 2016. Iowa Farm and Rural Life Poll: 2015 Summary Report. Accessed November 12, 2021. https://dr.lib.iastate.edu/server/api/core/bitstreams/b8e999e7 -f588-4dfb-9ed6-9ff9378lf37a/content.
- Arora, V. K., C. Singh, A. Sidhu, and S. Thind. 2011. "Irrigation, Tillage and Mulching Effects on Soybean Yield and Water Productivity in Relation to Soil Texture." Agricultural Water Management 98 (4): 563–68.
- Barker, J. B., D. M. Heeren, K. Koehler-Cole, C. A. Shapiro, H. Blanco-Canqui, R. W. Elmore, C. A. Proctor, S. Irmak, C. A. Francis, T. M. Shaver, and A. T. Mohammed. 2018. "Cover Crops Have Negligible Impact on Soil Water in Nebraska Maize-Soybean Rotation." *Agronomy Journal* 110: 1–1.
- Blanco-Canqui, H., T. Shaver, J. Lindquist, C. Shapiro, R. Elmore, C. Francis, and G. Hergert. 2015. "Cover Crops and Ecosystem Services: Insights from Studies in Temperate Soils." *Agronomy Journal* 107 (6): 2449–2474.

- Blaustein-Rejto, D. 2021. "Dishing the Dirt on Ag Carbon Credits." Accessed June 13, 2022. https://agfundernews.com/carbon-credits-in-ag-dishing-the-dirt.
- Bowman, M., and L. Lynch. 2019. "Government Programs that Support Farmer Adoption of Soil Health Practices: A Focus on Maryland's Agricultural Water Quality Cost-Share Program." *Choices*. Quarter 2.
- Butcher, K., A. F. Wick, T. DeSutter, A. Chatterjee, and J. Harmon. 2018. "Corn and Soybean Yield Response to Salinity Influenced by Soil Texture." *Agronomy Journal* 110: 1243–1253.
- Chu, M., S. Jagadamma, F. Walker, N. Eash, M. Buschermohle, and L. Duncan. 2017. "Effect of multispecies cover crop mixture on soil properties and crop yield." *Agricultural & Environmental Letters* 2 (1): 170030.
- Claassen, R., M. Bowman, J. McFadden, D. Smith, and S. Wallander. 2018. "Tillage Intensity and Conservation Cropping in the United States." Economic Information Bulletin No. 197. Economics Research Services, USDA.
- Clark, A., A. Decker, J. Meisinger, and M. McIntosh. 1997. "Kill Date of Vetch, Rye, and a Vetch–Rye Mixture: I. Cover Crop and Corn Nitrogen." Agronomy Journal 89 (3): 427–34.
- Conservation Technology Information Center (CTIC). 2015. Report of the 2014–2015 Cover Crop Survey. Joint publication of the CTIC and the North Central Region Sustainable Agriculture Research and Education Program. Accessed January 15, 2022. West Lafayette, Indiana.
- ——.2020. National Cover Crop Survey. https://www.ctic.org/files/CoverCrop Survey 2020 24mb.pdf.
- Corak, S. J., W. W. Frye, and M. S. Smith. 1991. "Legume Mulch and Nitrogen Fertilizer Effects on Soil Water and Corn Production." Soil Science Society of America Journal 55 (5): 1395–1400.
- Davey, K. A., and W. H. Furtan. 2008. "Factors That Affect the Adoption Decision of Conservation Tillage in the Prairie Region of Canada." *Canadian Journal of Agricultural Economics* 56 (3): 257–75.
- Dean, J. E., and R. Weil. 2009. "Brassica Cover Crops for Nitrogen Retention in the Mid-Atlantic Coastal Plain." Journal of Environmental Quality 38 (2): 520–28.
- Day, R. H. 1965. "Probability Distributions of Field Crops." Journal of Farm Economics 47: 713–41.
- Di Falco, S., and J. Chavas. 2009. "On Crop Biodiversity, Risk Exposure, and Food Security in the Highlands of Ethiopia." *American Journal of Agricultural Economics* 91 (3): 599–611.
- Di Falco, S., and M. Veronesi. 2014. "Managing Environmental Risk in Presence of Climate Change: The Role of Adaptation in the Nile Basin of Ethiopia." *Envi*ronmental and Resource Economics 57 (4): 553–77.
- Ding, G., X. Liu, S. Herbert, J. Novak, D. Amarasiriwardena, and B. Xing. 2006. "Effect of Cover Crop Management on Soil Organic Matter." *Geoderma* 130 (3–4): 229–39.
- Ding, Y., K. Schoengold, and T. Tadesse. 2009. "The Impact of Weather Extremes on Agricultural Production Methods: Does Drought Increase Adoption of Conservation Tillage Practices?" *Journal of Agricultural and Resource Economics* 34: 395–411.
- Dong, F., D. Hennessy, and H. Jensen. 2010. "Contract and Exit Decisions in Finisher Hog Production." American Journal of Agricultural Economics 92 (3): 667–84.
- Dong, F., D. Hennessy, H. Jensen, and R. Volpe. 2016. "Technical Efficiency, Herd Size, and Exit Intentions in US Dairy Farms." *Agricultural Economics* 47 (5): 533–45.

- Dong, S., Y. Jiang, Y. Dong, L. Wang, W. Wang, Z. Ma, C. Yan, C. Ma, and L. Liu. 2019. "A Study on Soybean Responses to Drought Stress and Rehydration." *Saudi Journal of Biological Sciences* 26 (8): 2006–2017.
- Dube, E., C. Chiduza, and P. Muchaonyerwa. 2012. "Conservation Agriculture Effects on Soil Organic Matter on a Haplic Cambisol after Four Years of Maize– Oat and Maize–Grazing Vetch Rotations in South Africa." Soil and Tillage Research 123: 21–28.
- Ebelhar, S. A., W. W. Frye, and R. L. Blevins. 1984. "Nitrogen from Legume Cover Crops for No-Tillage Corn 1." *Agronomy Journal* 76 (1): 51–55.
- Fleming, P. 2017. "Agricultural Cost Sharing and Water Quality in the Chesapeake Bay: Estimating Indirect Effects of Environmental Payments." *American Journal* of Agricultural Economics 99 (5): 1208–1227.
- Groff, S. 2015. "The Past, Present, and Future of the Cover Crop Industry." *Journal* of Soil and Water Conservation 70 (6):130A–133A.
- Hardaker, J. B., J. Richardson, G. Lien, and K. Schumann. 2004. "Stochastic Efficiency Analysis with Risk Aversion Bounds: A Simplified Approach." *Australian Journal of Agricultural and Resource Economics* 48 (2): 253–70.
- He, W., X. Luo, and G. Sun. 2014. "The Trend of GIS-Based Suitable Planting Areas for Chinese Soybean under the Future Climate Scenario." In *Ecosystem Assessment and Fuzzy Systems Management*, edited by B. Y. Cao, S. Q. Ma, and H. H. Cao, 325–38. Cham, Switzerland: Springer International Publishing.
- Hunter, M., M. Schipanski, M. Burgess, J. LaChance, B. Bradley, M. Barbercheck, J. Kaye, and D. Mortensen. 2019. "Cover Crop Mixture Effects on Maize, Soybean, and Wheat Yield in Rotation." *Agricultural and Environmental Letters* 4 (1).
- Jiang, P., and K. Thelen. 2004. "Effect of Soil and Topographic Properties on Crop Yield in a North-Central Corn–Soybean Cropping System." *Agronomy Journal* 96: 252–58.
- Karl, T. 1986. "The Sensitivity of the Palmer Drought Severity Index and Palmer's Z-Index to their Calibration Coefficients Including Potential Evapotranspiration." Journal of Climate and Applied Meteorology 25 (1): 77–86.
- Kaspar, T. C., D. J. Pulido, T. Fenton, T. Colvin, D. Karlen, D. Jaynes, and D. Meek. 2004. "Relationship of Corn and Soybean Yield to Soil and Terrain Properties." *Agronomy Journal* 96: 700–709.
- Kendall, M. G., and A. Stuart. 1977. The Advanced Theory of Statistics, Vol. 1, 168. New York: Macmillan.
- Kessler, A., S. Archontoulis, and M. Licht. 2020. "Soybean Yield and Crop Stage Response to Planting Date and Cultivar Maturity in Iowa, USA." *Agronomy Journal* 112: 382–94.
- Kim, K., J. P. Chavas, B. Barham, and J. Foltz. 2014. "Rice, Irrigation and Downside Risk: A Quantile Analysis of Risk Exposure and Mitigation on Korean Farms." *European Review of Agricultural Economics* 41 (5): 775–815.
- Koundouri, P., C. Nauges, and V. Tzouvelekas. 2006. "Technology adoption under production uncertainty: theory and application to irrigation technology." *American Journal of Agricultural Economics* 88 (3): 657–670.
- Kravchenko, A. N., D. Bullock, and C. Boast. 2000. "Joint Multifractal Analysis of Crop Yield and Terrain Slope." Agronomy Journal 92: 1279–1290.
- Laloy, E., and C. Bielders. 2010. "Effect of Intercropping Period Management on Runoff and Erosion in a Maize Cropping System." *Journal of Environmental Quality* 39 (3): 1001–1008.
- Langdale, G. W., R. L. Blevins, D. L. Karlen, D. K. McCool, M. A. Nearing, E. L. Skidmore, A. W. Thomas, D. D. Tyler, and J. R. Williams. 1991. "Cover Crop

Effects on Soil Erosion by Wind and Water." In *Cover Crops for Clean Water*, edited by W. L. Hargrove, 15–21. Ankeny, IA: SWCS.

- Le, D. T., R. Nishiyama, Y. Watanabe, M. Tanaka, M. Seki, K. Yamaguchi-Shinozaki, K. Shinozaki, L. Tran. 2012. "Differential Gene Expression in Soybean Leaf Tissues at Late Developmental Stages under Drought Stress Revealed by Genome-Wide Transcriptome Analysis." *PLoS One* 7: e49522.
- Lee, D. 2005. "Agricultural Sustainability and Technology Adoption: Issues and Policies for Developing Countries." *American Journal of Agricultural Economics* 87: 1325–1334.
- Lichtenberg, E., H. Wang, and D. Newburn. 2018. "Uptake and Additionality in a Green Payment Program: A Panel Data Study of the Maryland Cover Crop Program." Paper presented at the annual meeting of the Agricultural and Applied Economics Association, August 5–7, Washington, DC.
- Linkemer, G., J. Board, and M. Musgrave. 1998. "Waterlogging Effects on Growth and Yield Components in Late-Planted Soybean." Crop Science 38: 1576–1584.
- Lu, Y. C., K. B. Watkins, J. R. Teasdale, and A. A. Abdul-Baki. 2000. "Cover crops in sustainable food production." *Food Reviews International* 16 (2): 121–157.
- Major, D. J., D. Johnson, J. Tanner, and I. Anderson. 1975. "Effects of Day Length and Temperature on Soybean Development." Crop Science 15: 174–79.
- McDaniel, M. D., L. Tiemann, and A. Grandy. 2014. "Does Agricultural Crop Diversity Enhance Soil Microbial Biomass and Organic Matter Dynamics? A Meta-analysis." *Ecological Applications* 24 (3): 560–70.
- Myers, R., A. Weber, and S. Tellatin. 2019. "Cover Crop Economics: Opportunities to Improve Your Bottom Line in Row Crops." SARE Technical Bulletin. Accessed June 12, 2022. https://www.sare.org/wp-content/uploads/Cover-Crop -Economics.pdf.
- Miguez, F. E. and G. A. Bollero. 2005. "Review of Corn Yield Response under Winter Cover Cropping Systems Using Meta-analytic Methods." *Crop Science* 45 (6): 2318–2329.
- Miller, L., J. Chin, and K. Zook. 2012. "Policy Opportunities to Increase Cover Crop Adoption on North Carolina Farms." Masters Project, Duke University, Durham, NC.
- Mitchell J. P., A. Shrestha, and S. Irmak. 2015. "Trade-offs between Winter Cover Crop Production and Soil Water Depletion in the San Joaquin Valley, California." *Journal of Soil and Water Conservation* 70 (6):430–40,
- Motta, A. C., D. Wayne Reeves, C. Burmester, and Y. Feng. 2007. "Conservation Tillage, Rotations, and Cover Crop Affecting Soil Quality in the Tennessee Valley: Particulate Organic Matter, Organic Matter, and Microbial Biomass." Communications in Soil Science and Plant Analysis 38 (19–20): 2831–2847.
- Munawar, A., R. Blevins, W. Frye, and M. Saul. 1990. "Tillage and Cover Crop Management for Soil Water Conservation." Agronomy Journal 82 (4): 773–77.
- Muñoz, J. D., J. Steibel, S. Snapp, and A. Kravchenko. 2014. "Cover Crop Effect on Corn Growth and Yield as Influenced by Topography." *Agriculture, Ecosystems* & *Environment* 189: 229–39.
- National Drought Mitigation Center (NDMC). 2021. "What Is the USDM." Accessed November 11, 2021. https://droughtmonitor.unl.edu/About/WhatistheUSDM .aspx.
- National Research Council, 2007. Understanding American Agriculture: Challenges for the Agricultural Resource Management Survey. Washington, DC: The National Academies Press.
- Nelson, K. A., and C. Meinhardt. 2011. "Soybean Yield Response to Pyraclostrobin and Drainage Water Management." *Agronomy Journal* 103: 1359–1365.

- Nielsen, D. C., D. Lyon, R. Higgins, G. Hergert, J. Holman, and M. Vigil. 2016. "Cover Crop Effect on Subsequent Wheat Yield in the Central Great Plains." Agronomy Journal 108 (1): 243–56.
- OECD. 2011. Managing Risk in Agriculture: Policy Assessment and Design. Paris: Organization for Economic Cooperation and Development.
- Paz, J., W. D. Batchelor, G. L. Tylka, and R. G. Hartzler. 2001. "A Modeling Approach to Quantify the Effects of Spatial Soybean Yield Limiting Factors." *Transactions of the ASAE* 44 (5): 1329–1334.
- Pedersen, P., and M. Licht. 2014. "Soybean Growth and Development." PM 19948. Ames, IA: Iowa State University Extension and Outreach.
- Radočaj, D., M. Jurišić, V. Zebec, and I. Plaščak. 2020. "Delineation of Soil Texture Suitability Zones for Soybean Cultivation: A Case Study in Continental Croatia." Agronomy 10 (6): 823.
- Reddy, K. 2001. "Effects of Cereal and Legume Cover Crop Residues on Weeds, Yield, and Net Return in Soybean (Glycine max)." Weed Technology 15 (4): 660–68.
- Reddy, P. P. 2017. "Cover/Green Manure Cropping." In Agro-ecological Approaches to Pest Management for Sustainable Agriculture, 91–107. Springer, Singapore.
- Reuters. 2022. "Off-Season 'Cover' Crops Expand as U.S. Growers Eye Low-Carbon Future." Accessed January 5, 2022. https://www.reuters.com/markets /commodities/farming-climate-off-season-cover-crops-expand-us-growers-eye -low-carbon-future-2022-01-04/.
- Robertson, G. P., T. Bruulsema, R. Gehl, D. Kanter, D. Mauzerall, C. Rotz, and C. Williams. 2013. "Nitrogen–Climate Interactions in US Agriculture." *Biogeochemistry* 114 (1): 41–70.
- Sackett, J. L. 2013. "An NCR-SARE Cover Crop Project: Farmer-Cooperator Motivation and Agronomic Practices." Journal of the NACAA 6 (2).
- Sainju, U., B. Singh, and W. Whitehead. 2002. "Long-term effects of tillage, cover crops, and nitrogen fertilization on organic carbon and nitrogen concentrations in sandy loam soils in Georgia, USA." Soil & Tillage Research 63: 167–179.
- Sainju, U., B. Singh, F. Whitehead, and S. Wang. 2006. "Carbon supply and storage in tilled and nontilled soils as influenced by cover crops and nitrogen fertilization." *Journal of Environmental Quality*: 1507–17.
- Schnepf, M., and C. Cox. 2006. "Environmental benefits of conservation on cropland: the status of our knowledge." In *Environmental Benefits of Conservation* on Cropland: The Status of Our Knowledge, edited by M. Schnepf and C. Cox. Ankeny, IA: Soil and Water Conservation Society.
- Shaheen, T., M. Rahman, M. Shahid Riaz, Y. Zafar, and M. Rahman. 2016. "Soybean Production and Drought Stress." In *Abiotic and Biotic Stresses in Soybean Production*, edited by M. Miransari. San Diego, CA: Academic Press.
- Shane, W. W., and K. Barker. 1986. "Effects of Temperature, Plant Age, Soil Texture, and Meloidogyne incognita on Early Growth of Soybean." *Journal of Nematology* 18 (3): 320–26.
- Shanks L., D. Moore, and C. Sanders. 1998. "Soil erosion." In *Cover Cropping* in Vineyards. A Grower's Handbook, edited by C. A. Ingels, R. L. Bugg, G. T. McGourty, and L. P. Christensen, 80–85. Oakland, CA: University of California. Publication 3338.
- Smith, R. G., L. Atwood, and N. Warren. 2014. "Increased Productivity of a Cover Crop Mixture Is Not Associated with Enhanced Agroecosystem Services." *PloS One* 9 (5): p.e97351.
- Snapp, S. S., S. Swinton, R. Labarta, D. Mutch, J. Black, R. Leep, J. Nyiraneza, and K. O'Neil. 2005. "Evaluating Cover Crops for Benefits, Costs and Performance within Cropping System Niches." *Agronomy Journal* 97 (1): 322–32.

- Specht, J. E., D. Hume, and S. Kumudin. 1999. "Soybean Yield Potential— A Genetic and Physiological Perspective." Crop Science 39: 1560–1570.
- Steele, M., F. J. Coale, R. L. Hill. 2012. "Winter Annual Cover Crop Impacts on No-Till Soil Physical Properties and Organic Matter." Soil Science Society of America. Journal 76: 2164–2173.
- SARE (Sustainable Agriculture Research and Education Program). 2007. Managing Cover Crops Profitably, 3rd edition. Accessed on March 2, 2023. https://www.sare .org/Learning-Center/Books/Managing-Cover-Crops-Profitably-3rd-Edition.

——. 2017. "Cover Crops. What Is Sustainable Agriculture?" UC Division of Agriculture and Natural Resources. Accessed January 5, 2022.https://sarep.ucdavis.edu/sustainable-ag/cover-crops.

- Tack, J., A. Harri, and K. Coble. 2012. "More Than Mean Effects: Modeling the Effect of Climate on the Higher Order Moments of Crop Yields." *American Journal of Agricultural Economics* 94 (5): 1037–1054.
- Tilman, D. K., J. Fargione, B. Wolff, C. D'Antonio, A. Dobson, R. Howarth, D. Schindler, W. Schlesinger, D. Simberloff, and D. Swackhamer. 2001. "Forecasting Agriculturally Driven Global Environmental Change." Science 292: 281–84.
- Unger, P. W., and M. F. Vigil. 1998. "Cover Crop Effects on Soil Water Relationships." Journal of Soil and Water Conservation 53 (3): 200–207.
- USDA. 2022. "USDA Offers Expanded Conservation Program Opportunities to Support Climate Smart Agriculture in 2022." Accessed January 15, 2022. https://www.usda.gov/media/press-releases/2022/01/10/usda-offers-expanded -conservation-program-opportunities-support.
- USDA Climate Hubs. 2021. "Drought Resistant Practices." Accessed October 20, 2021. https://www.climatehubs.usda.gov/hubs/northeast/topic/drought-resistant -practices.
- USDA ERS. 2000. "Farm Resource Regions." Agricultural Information Bulletin Number 760. Accessed on September 5, 2021. https://www.ers.usda.gov /webdocs/publications/42298/32489_aib-760_002.pdf?v=42487#:~:text=The %20Farm%20Resource%20Regions%20are%20derived%20from%20four,Land %20Resource%20Regions%2C%20and%20NASS%20Crop%20Reporting %20Districts.
- USDA NRCS. 2018. Technical Note: Iowa Agronomy Technical Note 38 (Cover Crop Management). Accessed November 1, 2021. https://efotg.sc.egov.usda.gov /references/public/IA/Cover_Crop_Management_38_AGR_TN_2018_08.pdf.
 ______. 2019. NRCS Cover Crop Termination Guidelines Version 4.
- USDA Risk Management Agency. 2015. "Conservation Compliance—Highly Erodible Land and Wetlands. Risk Management Agency Fact Sheet." Accessed January 15, 2022. https://www.rma.usda.gov/en/Fact-Sheets/National-Fact -Sheets/Conservation-Compliance-Highly-Erodible-Land-and-Wetlands#:~: text=Highly%20erodible%20land%20is%20any%20land%20that%20can,total %20field%20acreage%20that%20contains%20highly%20erodible%20soils.
- Vitale, J. D., C. Godsey, J. Edwards, and R. Taylor. 2011. "The Adoption of Conservation Tillage Practices in Oklahoma: Findings from a Producer Survey." *Journal* of Soil and Water Conservation 66 (4): 250–64.
- Wallander, S., D. Smith, M. Bowman, and R. Claassen. 2021. Cover Crop Trends, Programs, and Practices in the United States. Economic Information Bulletin number 222, U.S. Department of Agriculture, Economic Research Service.
- Wandel, J., and J. Smithers. 2000. "Factors Affecting the Adoption of Conservation Tillage on Clay Soils in Southwestern Ontario, Canada." *American Journal of Alternative Agriculture* 15 (4): 181–88.

- Westcott, P., and M. Jewison. 2013. "Weather Effects on Expected Corn and Soybean Yields." FDS-13g-01. Economic Research Service/USDA.
- Wulanningtyas, H., Y. Gong, P. Li, N. Sakagami, J. Nishiwaki, and M. Komatsuzaki. 2021. "A Cover Crop and No-Tillage System for Enhancing Soil Health by Increasing Soil Organic Matter in Soybean Cultivation." Soil and Tillage Research 205.