

NBER WORKING PAPER SERIES

HEALTH CAPITAL AND THE PRENATAL ENVIRONMENT:  
THE EFFECT OF MATERNAL FASTING DURING PREGNANCY

Douglas Almond  
Bhashkar Mazumder

Working Paper 14428  
<http://www.nber.org/papers/w14428>

NATIONAL BUREAU OF ECONOMIC RESEARCH  
1050 Massachusetts Avenue  
Cambridge, MA 02138  
October 2008

We gratefully acknowledge comments from seminar participants at the NBER Cohorts meeting Spring 2007, UC Davis, Chicago Harris, the National Poverty Center Life Events Conference, Brown University, Stockholm University (IIES), the NBER Childrens Program, the University of Illinois (Chicago), the Federal Reserve Bank of San Francisco, Notre Dame, the Tinbergen Institute, and the University of Chicago (Booth School of Business). We also thank Lena Almond, Hoyt Bleakley, Janet Currie, Carlos Dobkin, Phoebe Ellsworth, Andrew Gelman, Jon Guryan, Jim Heckman, Hilary Hoynes, Darren Lubotsky, Doug Miller, Kevin Milligan, Diane Whitmore-Schanzenbach, and Kosali Simon for helpful comments. We thank Ana Rocca, Kenley Barrett, Sarena Goodman, and Shyue-Ming Loh for excellent research assistance and Christine Pal for careful copy editing. The views expressed here do not reflect the views of the Federal Reserve system, nor those of the National Bureau of Economic Research.

© 2008 by Douglas Almond and Bhashkar Mazumder. All rights reserved. Short sections of text, not to exceed two paragraphs, may be quoted without explicit permission provided that full credit, including © notice, is given to the source.

Health Capital and the Prenatal Environment: The Effect of Maternal Fasting During Pregnancy  
Douglas Almond and Bhashkar Mazumder  
NBER Working Paper No. 14428  
October 2008, Revised July 2009  
JEL No. I1,I12,J1,J14

**ABSTRACT**

We use the Islamic holy month of Ramadan as a natural experiment in fasting and fetal health. In Michigan births 1989-2006, we find prenatal exposure to Ramadan among Arab mothers results in lower birthweight and reduced gestation length. Exposure to Ramadan in the first month of gestation is also associated with a sizable reduction in the number of male births. In Census data for Uganda, Iraq, and the US we find strong associations between in utero exposure to Ramadan and the likelihood of being disabled as an adult. Effects are particularly large for mental (or learning) disabilities. We also find significant effects on proxies for wealth, the sex composition of the adult population, and more suggestive evidence of effects on schooling and earnings. We find no evidence that negative selection in conceptions during Ramadan accounts for our findings, suggesting that avoiding Ramadan exposure during pregnancy is costly or the long-term effects of fasting unknown.

Douglas Almond  
Department of Economics  
Columbia University  
International Affairs Building, MC 3308  
420 West 118th Street  
New York, NY 10027  
and NBER  
da2152@columbia.edu

Bhashkar Mazumder  
Federal Reserve Bank of Chicago  
230 S. LaSalle Street  
Chicago, IL 60604  
bmazumder@frbchi.org

# 1 Introduction

The early childhood environment, including that experienced before birth, can have permanent effects on health capital and skill formation [Cunha and Heckman, 2007]. Recent studies by economists have utilized exogenous shocks “caused by conditions outside the control of the mother” to provide compelling evidence in support of the fetal origins hypothesis [Currie, 2009]. These studies have typically leveraged relatively uncommon historical events, such as exposure to famine or infectious disease, for identification. An unresolved question is whether more commonly encountered early-life exposures also exert significant long-term effects on health and human capital.

In this study, we consider a common early-childhood exposure that is ongoing today: disruptions to the timing of prenatal nutrition from meal skipping during pregnancy.<sup>1</sup> Specifically we consider the effects of maternal fasting during daylight hours during the month of Ramadan. Three in four Muslim pregnancies overlap with Ramadan and surveys indicate the majority of pregnant Muslims observe the fast. There is currently no consensus on whether prenatal fasting harms newborn health. We provide new evidence on fasting’s effects on birth outcomes and the first evidence of effects later in life using large-sample microdata on Muslims in the United States, Iraq, and Uganda.

Our methodological approach addresses a key flaw in previous studies of Ramadan fasting and birth outcomes. These epidemiological studies have compared pregnant women who fasted to those who did not at a point in time, under the basic assumption that the decision to fast is exogenous.<sup>2</sup> Instead, we compare births over many years where Ramadan overlaps with pregnancy to those where Ramadan does not and estimate the reduced form effect of Ramadan’s timing.<sup>3</sup> That is, we estimate an “intent to treat” (ITT) effect without relying on the decision whether to fast for identification.<sup>4</sup> This approach yields distinct ITT estimates for specific months of gestation; Muslim births where Ramadans

---

<sup>1</sup>Nearly 1 in 4 women report skipping meals during pregnancy in the US [Siega-Riz et al., 2001].

<sup>2</sup>Pre-pregnancy BMI, along with other characteristics, has been found to predict fasting observance [Kavehmanesh and Abolghasemi, 2004].

<sup>3</sup>We do not observe whether mothers fasted in our data. See Section 6.

<sup>4</sup>We draw an analogy with research designs where there is random assignment to treatment and control groups but where compliance may be endogenous. In our case we assume that the timing of Ramadan relative to pregnancy is exogenous, but that the decision to fast is endogenous and not be universally observed.

falls in the early postnatal period serve as the control group. Because Ramadan follows a lunar calendar, its occurrence moves forward by roughly 11 days each year. Thus, we can disentangle the effect of prenatal overlap with Ramadan from season of birth, which is also related to health in adulthood [Doblhammer and Vaupel, 2001, Costa and Lahey, 2005, Costa et al., 2007, Buckles and Hungerman, 2008].

Using natality data from Michigan, we find that prenatal exposure to Ramadan lowers birth weight and reduces gestation length. Further, the likelihood of a male birth is about 10% lower when Ramadan falls very early in pregnancy and occurs during the peak period of daylight fasting hours. Using Census data for the United States, Iraq, and Uganda we find long-term effects on adult health and some economic outcomes. We generally find the largest effects on adults when Ramadan falls early in pregnancy. Rates of adult disability are roughly 20% higher, with specific mental disabilities showing substantially larger effects. Our estimates are conservative to the extent that Ramadan is not universally observed. Importantly, we detect no corresponding outcome differences when the same design is applied to non-Muslims as a falsification test.

As we discuss in Section 2, our findings are plausible in the context of the biomedical literature where studies have documented that even relatively short fasts lasting 12 hours result in dramatic changes in the metabolic biochemical profiles of pregnant mothers. Such changes have specifically been found in the context of Ramadan fasting. One recent study has documented heightened levels of the hormone cortisol among pregnant women fasting during Ramadan [Dikensoy et al., 2009], which has been associated with adverse long-term health effects through fetal programming of the neuro-endocrine system. Our results are also consistent with studies of the Dutch famine which found particularly large long-term health effects associated with early-pregnancy exposure. Several studies have also documented that maternal nutrition during pregnancy varies positively with male births (in the cross-section).

Our identifying assumption is that pregnancies are not timed relative to Ramadan along unobserved determinants of health. We present evidence that pre-determined maternal and paternal characteristics are not systematically related to the timing of conception relative to Ramadan. In our Michigan data, we observe maternal education, whether previous birth was low birth weight, maternal diabetes, whether mother smoked, and

whether Medicaid paid for the delivery: each is unrelated to the timing of pregnancy relative to Ramadan. Not surprisingly, controlling for these factors has a negligible effect on our ITT estimates.

The remainder of the paper is organized as follows. Section 2 describes previous epidemiological work on Ramadan and health, referencing additional supporting material in the Appendix. Section 3 describes our natality and Census data, ITT measures, and econometric model. Section 4 presents our results for birth outcomes in Michigan and Section 5 describes our findings for adult outcomes in Uganda, Iraq, and the US. Section 6 interprets our results and discusses future research.

## 2 Previous Literature

We briefly discuss previous studies of maternal fasting and health outcomes. In the interest of space, we refer the reader to additional background material in Appendix, Section A.<sup>5</sup>

Previous studies of fasting and health have evaluated: 1) measures of maternal and fetal health during pregnancy, and; 2) health at birth. In contrast to prenatal health, measurement of newborn health is relatively standardized (e.g., by birth weight or infant mortality). However, studies of maternal and fetal health allow for comparisons over time for the same pregnancy – in and out of the fasting state – addressing the potential endogeneity of the fasting decision.

### 2.1 Ramadan and Health During Pregnancy

Writing in *The Lancet*, Metzger et al. [1982] documented a set of divergent biochemical measures among pregnant women who skipped breakfast in the second half of pregnancy. Relative to twenty-seven non-pregnant women with similar characteristics, “circulating fuels and glucoregulatory hormones” changed profoundly in twenty-one pregnant women when the “overnight fast” was extended to noon on the following day (relative to post-prandial baseline). Further, plasma glucose and alanine was lower in the pregnant women

---

<sup>5</sup>Appendix Section A summarizes the effects of fasting on caloric intake and weight gain; the rates of observance of Ramadan fasting by pregnant women; the potential health effects of maternal biochemical changes on offspring; fasting and fetal programming; studies of Ramadan fasting’s effect on birth outcomes; and our hypotheses relating specific periods of exposure to particular outcomes.

than in the non-pregnant women after 12 hours of fasting while levels of free fatty acids and beta-hydroxybutyrate, a ketone, were significantly higher. This set of biochemical changes, also known as “accelerated starvation”, occurred after only “minor dietary deprivation” for both lean and obese women. Metzger et al. [1982] concluded that meal-skipping “should be avoided during normal pregnancy.” Meis and Swain [1984] found that *daytime* fasts during pregnancy caused significantly lower glucose concentrations than nighttime fasts.

According to survey data, most pregnant Muslims observe the Ramadan fast (See Appendix Section A.1.2 for details). Following the study of breakfast skipping by Metzger et al. [1982], Ramadan fasting was likewise found to cause accelerated starvation among pregnant women in Gambia [Prentice et al., 1983] and in England [Malhotra et al., 1989]. Mirghani et al. [2004] found that maternal glucose levels were lower in the fasting state compared to the postprandial baseline, a difference accentuated by the number days fasted: “the effect on maternal glucose levels during Ramadan fasting is cumulative.” Several studies of maternal fasting during Ramadan have found adverse effects carried over to measures of fetal health: fetal breathing movements and fetal heart rate accelerations [Mirghani et al., 2004, 2005].

Recently, Dikensoy et al. [2009] reported that Ramadan fasting is associated with increases in cortisol levels during pregnancy, but not for non-fasting pregnant women (both relative to pre-pregnancy levels). This finding is of interest because cortisol is a stress hormone frequently invoked as a potential mechanism through which prenatal experiences may “program” adult health [Kapoor et al., 2006].

To summarize, there is fairly consistent evidence that fasting during pregnancy has a “first-stage” effect on maternal and fetal health measures. We summarize the literature on potential fasting sequelae in Appendix Section A. Despite uncertainty whether these first stage effects carry over to birth outcomes and longer-term effects (See Section 2.2 below), the Institute of Medicine nevertheless recommends pregnant women should “eat small to moderate sized meals at regular intervals, and eat nutritious snacks” [Institute of Medicine, 1992:45]. Similarly, the American College of Obstetricians and Gynecologists recommends that pregnant women avoid skipping meals.<sup>6</sup>

---

<sup>6</sup>[http://www.acog.org/publications/patient\\_education/bp087.cfm?printerFriendly=yes](http://www.acog.org/publications/patient_education/bp087.cfm?printerFriendly=yes)

## 2.2 Ramadan and Perinatal Health

The effect of fasting on birth outcomes is less clear. Most previous studies have drawn comparisons over only a single Ramadan season. Since the panel-data dimension is generally absent for analyses of birth outcomes, studies have resorted to strong assumptions on the comparability of fasters and non-fasters. These two groups are likely different in ways that would generate differences in birth outcomes absent any causal effect of fasting.<sup>7</sup> This basic weakness in design has been exacerbated by: 1) small sample sizes that in general would only be able to distinguish quite large effects from zero; 2) consideration of Ramadan fasts observed exclusively in mid or late gestation. We refer the reader to the more detailed discussion of these studies in Appendix A.4.1.

No previous study has exploited idiosyncratic variation across birth cohorts in the timing of Ramadan relative to birth. As Ramadan’s forward movement through the western calendar is slow, the separation of Ramadan from seasonal effects on birth outcomes (e.g., Doblhammer and Vaupel [2001], Costa and Lahey [2005]) requires data across many birth years. Data availability, therefore, may have precluded implementation of an ITT analysis. Similarly, no previous study has exploited the number of daylight hours during the Ramadan fast for identification (which is clearly not feasible for populations living near the equator, e.g., in Uganda or Indonesia).

Finally, we have not found any previous studies that trace adult health (or socioeconomic) outcomes back to *in utero* Ramadan exposure. The study closest to ours in this respect is by Azizi et al. [2004] who found no significant difference in the IQ’s of school-age children by maternal fasting behavior during the third trimester (please see Appendix Section A.4.2 for details). Subsequent to our study, Ewijk [2009] analyzed IFLS data from Indonesia, finding evidence of long-term effects of fasting.<sup>8</sup>

## 3 Data and Methodology

Our analysis requires microdata with information on:

---

<sup>7</sup>See footnote 2.

<sup>8</sup>Ewijk [2009] graciously notes that we are “the first to systematically examine [Ramadan’s] long-term effects.”

1. a substantial number of Muslims;
2. precise information on birth date (i.e., more detailed than age in years);
3. coverage of many birth cohorts (i.e., birth years);
4. health outcomes.

In datasets with a large number of both Muslims and non-Muslims, we would also like to distinguish between these groups. We summarize the handful of suitable datasets below (see Appendix B for details), followed by our econometric approach.

### 3.1 Michigan Natality Files

From Michigan’s Division for Vital Records and Health Statistics, we obtained birth certificate microdata for 1989 to 2006 in Michigan – approximately 2.5 million birth records.<sup>9</sup> Although, there is no information on religion, ancestry of the mother is reported which allows us to construct a proxy for whether the mother is Muslim based on reported “Arab” ancestry (or reported ancestries from predominantly Muslim countries).<sup>10</sup> Compared to other US states, Michigan has a relatively large Arab population.<sup>11</sup> There are a total of about 50,000 births to mothers of Arab ancestry (about 2.2 percent of MI births) over this period. While there is a large population of Arabs around Detroit, the Arabs are reasonably dispersed throughout the State (see Appendix Figure A2, Panel A).

Since a large fraction of Arabs in Michigan are actually Chaldeans – a sect of Christianity – our proxy may misclassify many mothers and thereby attenuate our estimated effects.<sup>12</sup> We use the 2000 US Census SF3 (1 in 6 sample) data to identify Michigan zipcodes with heavy concentrations of Chaldeans – who presumably do not observe the fast – relative to Arabs (see Appendix Figure A2, Panel B). In some specifications, we will drop observations from these zipcodes to compare ITT estimates. Appendix Table A2 provides summary statistics for Michigan’s natality data.

---

<sup>9</sup>Special thanks to Michael Beebe and Glenn Copeland in Michigan’s Vital Statistics Office.

<sup>10</sup>See Section B.1 of the Appendix for more detail.

<sup>11</sup>We thank Carlos Dobkin (UCSC) for suggesting we focus on Michigan’s Muslim population.

<sup>12</sup>According to the 2000 Census, about a quarter of those of an Arabic speaking ancestry in Michigan are Chaldean Christians. Our estimates based on the Detroit Arab American Study (DAAS) suggest that about 47% of those who self-identify as “Arab American” in the Detroit area are Chaldean.



## 3.2 Census Data

To consider whether health in adulthood is affected by prenatal Ramadan exposure, we analyze Census microdata for three countries. Data from the Uganda 2002 Census are best suited for our analysis because religion is reported, there are large numbers of both Muslims and non-Muslims in Uganda, month of birth is reported, and a host of disability measures are queried. While less well-suited to our analysis, 1997 Iraq Census data and US Census data allow us to assess whether the basic Uganda results are replicated in other settings.

### 3.2.1 Uganda Census 2002

Our sample of Muslim adults includes approximately 80,000 men and women between the ages of 20 and 80 in 2002. Muslims constitute about 11% of Uganda's population and have more schooling and lower rates of disability than non-Muslims (Appendix Table A3). Both Muslims and non-Muslims share a strong seasonality in the number of births. Muslims tend to live in the southeastern portion of the country.

The Uganda Census asks about specific disabilities including: blindness or vision impairments, deafness or hearing impairments, being mute, disabilities affecting lower extremities, disabilities affecting upper extremities, mental/learning disabilities, and psychological disabilities (lasting six months or longer). As only about 5% of adults report a disability compared to over 10% in the US Census, disabilities recorded in the Uganda Census may be more severe. Further, there is information on the *origin* of disabilities: congenital, disease, accident, aging, war injury, other or multiple causes. In the absence of direct measures of economic status we use home ownership. We also consider several other socioeconomic outcomes such as literacy, schooling, and employment.

### 3.2.2 Iraq Census 1997

Although religion is not reported in the Iraq Census, roughly 97% of the population is Muslim, minimizing concerns about mis-classification. Our sample contains over 288,000 individuals between the ages of 20 and 80. The large sample size is offset by low rates of reported disabilities. At 1.5%, Iraqis are substantially less likely than Americans (around

12%) or Ugandans (around 5%) to report a disability. Along with a general disability question, there are specific questions about disabilities involving sight, hearing, lower extremities, upper extremities, and psychological disabilities. In contrast to Uganda, there is no variable to assess mental/learning disabilities.

In addition to home ownership, we consider polygyny as a second proxy for wealth/status. Under Iraqi law, courts may only allow polygyny if husbands are able to support financially multiple wives and if they are able to maintain separate households for each wife [Iraq Legal Development Project, 2005].<sup>13</sup> More generally, polygyny reflects high male status [Edlund, 1999]. We also examine whether individuals attained more than primary schooling and employment status at enumeration. Sample means for our outcomes are shown with the regression results in Table 7.

### 3.2.3 US Census 1980, ACS 2005-2007

Our third Census sample is composed of immigrants to the US who were born in predominantly Muslim countries.<sup>14</sup> We use a 6% sample from the 1980 Census along with a pooled sample of the American Community Surveys (ACS) for the years 2005 through 2007 (3% sample). The ACS is modelled on the long form of the decennial Census. We use these years because they provide the quarter of birth.<sup>15</sup> Not observing birth month

---

<sup>13</sup>Under Iraqi Personal Status Code Number 188, Article 3(4):

Marriage of more than one wife is prohibited in the absence of judicial permission on two conditions: (a) The husband has financial sufficiency to marry more than one wife. (b) He should have a legal interest.

Iraqi Personal Status Code Number 188, Article 26 states:

The husband should not house his second wife in the same house with the first one without her approval, and should not house any other relative with her without her approval, except his minor child.

Roughly 2% of Iraqi men report polygynous unions.

<sup>14</sup>Specifically, we use countries with at least an 80 percent Muslim population. To assess the magnitude of misclassification attributable to this birth-county proxy, the same proxy variable was created in the 2001 Canada Census, which includes self-reported religion. 67% of Canadian immigrants from these 80% Muslim countries reported being Muslim. We thank Kevin Milligan (UBC) for this information. See Appendix Section B.5 for additional details.

<sup>15</sup>In addition to not knowing birth month, we do not know the exact birth *year* in the ACS since it is not asked and age is not reported as of a specific enumeration date as it is in the decennial Census. Given that Ramadan exposure shifts by only 11 days from year to year, using `survey year - age` provides a good approximation of birth year for the purposes of constructing Ramadan exposure measures at the quarterly level. The correlation between Ramadan exposure using survey year minus age, and survey

dulls the empirical comparisons that we can make. On the positive side, we obtain a large, national sample of US immigrants from Muslim countries in which to implement our ITT analysis. Data quality is high, and includes additional outcomes beyond disability, such as earnings.<sup>16</sup> Selected sample means are shown at the bottom of Table 8.

### 3.3 Ramadan Measures

We record start and end dates for the 104 Ramadans in the 20th century and construct the following measures of prenatal Ramadan exposure:<sup>17</sup>

**exp<sub>pct</sub>** calculates the fraction of days over the subsequent 30 days that overlap with Ramadan.

**exp<sub>hrspct</sub>** utilizes the number of daylight hours in each day to proxy for the length of the Ramadan fast.<sup>18</sup> The numerator is the number of daylight hours over the next 30 days that overlap with Ramadan; the denominator is the maximum number of daylight hours over any 30 day period over the entire sample period (which depends only on latitude). Daylight hours in Michigan vary from a low of around 9 to a high of over 15 at the summer solstice when the effects of accelerated starvation may be most evident. Please see Appendix Figure A3 for an illustrative example of *exp<sub>hrspct</sub>* versus *exp<sub>pct</sub>* coding from 1989 (and the associated description in Appendix Section B.1).

**ramp<sub>pct</sub>** is the fraction of days in each month that overlap with Ramadan.<sup>19</sup> *ramp<sub>pct</sub>* is used when we observe just the birth month, as in our Uganda and Iraq samples and

---

year minus age minus 1, is about 0.93.

<sup>16</sup>Specifically we take the log of annual earnings and use the CPI for inflation adjustment. For consistency, we restrict the sample to ages between 20 and 80. Earnings are restricted to a sample between the ages of 20 and 60.

<sup>17</sup>Many websites translate dates from the Islamic (Hijri) calendar. We used the following website <http://www.oriold.unizh.ch/static/hegira.html>, but verified the dates from a second source.

<sup>18</sup>The beginning of the Ramadan fast actually precedes sunrise and begins at the time of the morning prayer (*fajr*). The precise timing of the morning prayer may vary across mosques and typically depends on a rule regarding the angle of the sun relative to the horizon. For this reason we actually understate the number of fasting hours in our data. Daylight hours are measured for the city of Dearborn which contains a large share of the state's Arab population.

<sup>19</sup>We opted to use this measure, rather than a simple dummy variable since it provides a continuous measure of treatment (more power).

is calculated for each month (not each day as for measures above). As Uganda is at the equator, the number of daylight hours is fairly constant over the year at 12.

Each of the above measures is calculated at up to nine different points in time corresponding to the gestation months (ten in some specifications where we include the month prior to conception). *exppect* and *exphrspect* are calculated for the day beginning each pregnancy month. This feature of our research design is useful as specific health outcomes may be more or less susceptible to nutrition depending on the stage of pregnancy (see Appendix Table A1 and accompanying text in Appendix Section A). Finally, we construct a fourth exposure measure by taking the average of *rampct* for each quarter of the year. This quarterly measure is used for our US sample, where we observe just the quarter of birth.

In general, we assign the Ramadan ITT by counting backwards from the birth date. We consider alternatives to this approach where we have estimates of gestation length (Michigan). In the rest of our data, we do not observe gestation length. Furthermore, measurement error in gestation length, commonly based on date of last menstrual period, may be substantial [Pearl et al., 2007]. For Michigan, we consider the following three assignment rules:

1. The first approach uses the exact date of birth and simply assumes that all births have a normal gestation length of 40 weeks since last menstrual period. The Ramadan exposure measures are assigned by going backwards from the birth date in 30 day increments and using *daily* exposure measures (*exphrspect* and *exppect*) from 30 days prior to birth to 270 days prior to birth. Using this approach the measure of Ramadan exposure 9 months prior to birth is a proxy for the actual exposure during the “first month” of gestation.
2. Our second approach incorporates the measures of gestation from the natality data to match each individual to an estimated date of conception.<sup>20</sup> We then assign Ramadan exposure for the first month based on the daily exposure measures for the

---

<sup>20</sup>Gestation based on last menstrual period (LMP) is used except if it is missing or if it differs with the physician estimated gestation by more than 14 days, in which case the physician estimated measure is substituted. The conception date is estimated as occurring 14 days after LMP.

date that is 4 days prior to the estimated date of conception.<sup>21</sup> We then proceed to assign Ramadan exposure measures forward in 30-day increments. Using this approach a child born premature after say, 34-35 weeks of gestation would only have been *in utero* for about 8 months and therefore only the first 8 exposure variables are actually relevant for birth outcomes.

3. Our third approach ignores information on exact birth date and actual gestation and mimics what we can do with our Census samples where we only know month of birth. Here we match individuals to the *rampct* measure for each of the 9 months prior to birth.

### 3.4 Econometric Model

We regress each outcome,  $y_i$ , on a set of nine Ramadan exposure measures (either *exhrspct*, *exppct* or *rampct*). Separate exposure measures for each gestation month  $k$  are included simultaneously in each regression, even though an individual will only be exposed to Ramadan in at most two (adjacent) months of gestation. The effects of Ramadan exposure in a given month of gestation, therefore, are measured relative to no prenatal exposure to Ramadan – i.e., when Ramadan falls in the two to three months after birth.

$$y_i = \alpha + \sum_{k=1}^9 \theta_k \cdot \text{exhrspct}_k + \text{year}_y + \text{month}_m + \text{geog}_g + \varepsilon_i \quad (1)$$

Additional controls include birth year dummies, a set of calendar birth month dummies (or conception month dummies if gestation information is used), and a set of dummies that measure geographic location at the time of birth.<sup>22</sup> In the Michigan analysis we also include a number of largely predetermined variables as additional controls: mother’s age, mother’s age squared, mother’s years of education, father’s age, father’s age squared, father’s education, a dummy for missing father’s education, parity, tobacco use during pregnancy, alcohol use during pregnancy, the number of previous pregnancies where the

---

<sup>21</sup>This ensures that we have lined up the gestation month exposure in a parallel way to that used in the first approach. In other words, if a birth has exactly 40 weeks or 280 days of gestation, using either approach we will start measuring gestation exactly 270 days prior to birth.

<sup>22</sup>In Michigan we use 84 county dummies. In Uganda we use 56 district of birth dummies. In Iraq we use 18 governates of birth and in the US we use 38 countries of birth.

child was born dead, and whether the birth was paid for by Medicaid (an income proxy).<sup>23</sup> In our pooled samples of adult men and women in Uganda, Iraq, and the US we also include a female dummy.

In a typical specification where we include the nine exposure measures simultaneously, we also run an  $F$ -test on the joint significance of all nine coefficients. This tests the overall effect of Ramadan exposure during any point in gestation. In addition, since our hypotheses for some outcomes (Table A1) suggest an effect only in specific gestation months, we also run tests of equality of all coefficients.

When we use a non-Muslim sample as a falsification check (in a separate regression), the birth timing and birth location effects are allowed to vary across groups (along with the other parameters). For estimates on population counts by month we use aggregate measures at the cell level where cells are defined by each of the distinct conception or birth months over the sample period. For Michigan, this yields 216 cells (18 years  $\times$  12 calendar months).

For the adult US sample where we only observe quarter of birth, there is an added difficulty in identifying a control period with no *in utero* Ramadan exposure. If we assume that all births were of normal gestation length, then individuals were *in utero* for each of the two quarters prior to the birth quarter, along with some portion of their birth quarter. This would suggest that we use the quarter three quarters prior to the birth quarter as the omitted category. A more conservative approach would treat only those with Ramadan exposure in the two quarters prior to birth as treated and omit both of the other two birth quarters. We will show results using each of the two approaches.

## 4 Michigan Results

### 4.1 Birthweight Weight and Gestation

Table 1 presents the results for birthweight. In the first four columns we assign births their Ramadan exposures by counting backwards from the date of birth (i.e., ignoring the reported information on gestation and assuming pregnancies all had normal gestation

---

<sup>23</sup>Parity is defined as the number of previous live births. Alcohol and tobacco use are arguably endogenous since their use may be reduced during the month of gestation that overlaps with Ramadan.

length). We designate the coefficients on Ramadan exposure according to the number of months prior to birth.

Columns (1) and (2) use measures of Ramadan exposure that incorporate hours of daylight (*exphrspct*). The first entry of column (1) indicates that birthweight was about 40 grams lower if an Arab mother was in the first month of pregnancy (i.e. nine months before birth) during the peak period of daylight hours and Ramadan coincided with the first month of gestation. This result is significant at the 5 percent level. In addition, gestation months 2, 4, and 6 also show significant negative effects of around 40 grams. We also find that the  $F$ -test on the joint importance of all the prenatal Ramadan exposure measures is significant at the 8 percent level. The test of the equality of coefficients is not rejected at conventional significance levels.

In column (2), we also include Ramadan exposure 10 months prior to birth as a falsification and robustness check. As expected there is no detectable effect on birth weight and its inclusion does not substantially affect the point estimates for other months. In column (3) we measure Ramadan exposure by the fraction of days in the gestation month that overlap with Ramadan. This treats a Ramadan falling in winter the same as one during summer. It also changes the interpretation of the coefficient to reflect the average effect of Ramadan exposure across all seasons. As expected, this reduces the magnitude of the estimated effects to around 25-30 grams, which remain statistically different from zero. In column (4) we again include exposure in the month prior to the beginning of pregnancy and find no effect. Column (5) ignores information about the exact date of birth and only utilizes the birth month to assign exposure measures of the fraction of days in each month that overlap with Ramadan. With this approach we now find only the second month of gestation to be significant and that the coefficients are far from jointly significant or significantly different from one another. Nevertheless, the effect size remains in the 30 gram range. Using only birth month re-allocates some Ramadan ITT assignments to adjacent months.<sup>24</sup>

In columns (6)-(9) of Table 1, we assign Ramadan exposure using the time elapsed since conception, and exclude the 7% of births that are pre-term (< 37 weeks gestation).

---

<sup>24</sup>For example, the effect of Ramadan falling nine months prior to birth in column (4) appears smoothed into the adjacent month (8 months before birth) in column (5).

This ensures that at least the first 8 exposure measures are reasonably interpretable since all of these births would have been *in utero* for 8 months.<sup>25</sup> Columns (6) and (7) utilize the daylight hours index of exposure. We find effects that are statistically significant at the 5 percent level for exposure in month 1. As in columns (1) and (2), the size of these effects are between 35 and 40 grams. In column (7) we again show no effect in the month prior to conception. Thus, the pattern of birth weight effects we find are not an artifact of assigning the Ramadan ITT using date of birth. Columns (8) and (9) show smaller effects from first month exposure when seasonal differences in the length of the diurnal fast is ignored. The smaller sample used in columns (6) through (9) that ignores gestation length have considerably higher significance levels on our joint test of an effect across all gestation months (i.e., when we drop premature births and estimate the specifications in columns (1)-(5)). This suggests that part of the difference between the two panels of Table 1 may be due to selection on nearly full-term births. For example, the magnitude of the effects for Ramadans that fall 4 months prior to birth are attenuated in the full term sample. We explore the role of conditioning on different gestation lengths in Appendix Table A4. The associated text discusses impacts on gestation length itself, which are modest compared to those for intrauterine growth retardation (IUGR).

## 4.2 Refutability Check Using Non-Arab Births

Identification of our ITT estimates comes from an idiosyncratic, non-linear function of date of birth. If this function somehow introduced a mechanical relationship between birth date and health outcomes (e.g. picking up a time trend), it should also be observed among non-Muslims. As a check on the validity of the results above, we apply the same ITT to our non-Arab sample in Table 2. The first six columns assign exposure assuming normal gestation for the whole sample, while the seventh through ninth columns utilize the gestation data and present the results for the sample of full-term births. Within each of these sets of results, we use either all non-Arab mothers or a “super control” group of non-Arab mothers living in zipcodes without any Arabs (according to the 2000 census

---

<sup>25</sup>In our tables we use italics to signify estimates that may include post natal exposure for some sample members.



data).<sup>26</sup> We find no birthweight effects among non-Arabs comparable to the results in Table 1.<sup>27</sup>

### 4.3 Summary and Discussion of Birth Weight Results

Overall, *in utero* exposure to Ramadan is associated with lower birth weight among Michigan’s Arab mothers. A full month of exposure to Ramadan during the peak period of daylight hours could lead to a reduction in birth weight of about 40 grams especially when Ramadan falls in the first month of gestation. Nevertheless, the size of this effect is relatively small: 40 grams is only about 1.2 percent of the mean birth weight for Arabs. These effects are population averages and do not account for the fact that some fraction of these women are not actually fasting and more importantly, perhaps, that we may be including a sizable fraction of Non-Muslim women among the Arabs.<sup>28</sup> In terms of the birth weight distribution, it appears that most of the estimated effect for early pregnancy exposure is in the middle of the distribution (see Appendix Figure A4), rather than a disproportionate increase in the likelihood of low birth weights, which may be more closely tied to other measures of newborn health than variation at higher birth weights [Almond et al., 2005]. In any case, we take a conservative view of the birth weight findings and emphasize that they provide evidence of a “first stage” effect of Ramadan fasting.

Finally, if Ramadan observance during pregnancy varied by socioeconomic or health status, treatments effects would presumably also show a corresponding gradient, other things equal. We observe no systematic gradient in the size of the birth weight effects by maternal education, Medicaid use, or month prenatal care was initiated (results available from authors). If treatment effects are relatively homogeneous, this suggests that fasting observance is high or fairly uniform across socioeconomic groups by month of gestation.

---

<sup>26</sup>The results are unaffected if we use the natality data to identify non-Arab zipcodes. However, given that we have more confidence in the reporting of ancestry detail in the Census we opted to use those figures.

<sup>27</sup>In column (3) of Table 2 we obtain two estimates of about -4 grams that are only significant at the 10 percent level, which may be chance occurrences among the 81 reported point estimates.

<sup>28</sup>In Appendix Table A4 we show that these effects rise are somewhat larger when we exclude zipcodes with a higher Chaldean/Arab ratio.

## 4.4 Fetal Death and the Sex Ratio at Birth

Mathews et al. [2008] found that poor maternal nutrition (possibly due to breakfast skipping), around the time of conception skews the sex ratio in favor of girls, most likely through the selective attrition of male conceptuses. Similarly, Almond et al. [2009] found that severe morning sickness in early pregnancy is associated with female births, but also a 50% fetal death rate.<sup>29</sup> More generally, maternal nutrition among mammals close to conception is positively associated with the likelihood of male offspring Cameron [2004].

We consider Ramadan’s effect on sex at birth and the number of lives births in Table 3. Using our full sample of Arab mothers (column 1) we find a large effect of -3.7 percentage points ( $p$ -value = 0.06) on the likelihood of a male birth from exposure to Ramadan during the longest diurnal fast in month 1 of pregnancy. In column (2) when we restrict the zipcodes to those with fewer Chaldeans relative to Arabs, this point estimate rises substantially to -6.6 percentage points and is significant at the 1 percent level. Interestingly, we find some suggestive evidence of an effect in the month prior to conception possibly reflecting the persistent effect of fasting just prior to conception. Columns (3) and (4) show no analogous effects for our non-Arab samples.

To shed light on the cause of the sex ratio change, we aggregate counts of total births, male births, and female births based on the month of *conception*. This results in 216 observations for conceptions between April 1988 and March 2006. The mean fraction of births (weighted) that are male is 51.8 percent for Arabs over this period and the mean sex ratio (male to female births) is 1.086. For non-Arabs, the respective means are 51.3 and 1.054. We regressed the log of the number of births on the *exphrspect* measures of Ramadan exposure including dummies for month and year of conception and our full set of controls.

Results for the more restricted sample of Arabs are shown in columns (5) through (7). Peak exposure to the Ramadan fast in the month after conception is associated with a 16 percent decline in total births. If male vulnerability [Kraemer, 2000] is the culprit, this drop should be concentrated among male births. When we examine this by sex, we indeed find this is driven by a 26 percent drop in male births ( $p$ -value = 0.005), while female

---

<sup>29</sup>By fetal death, we mean any attrition between conception and live birth. This could include attrition during embryonic development before a fetus is fully formed.

births fall by a statistically insignificant 4 percent.<sup>30</sup> This decline in births associated with fasting around the time of conception is probably not due to other behavioral changes associated with Ramadan since it is difficult to imagine an alternative mechanism which impacts *sex-specific* fertility.

Among those conceived shortly after Ramadan, (i.e., exposure in month zero, the month prior to pregnancy) there is a statistically insignificant increase of 6.3 percent in total live births associated with full exposure with some suggestive evidence of a female skew.<sup>31</sup> The fact that total births rise somewhat after Ramadan is not so surprising since the end of Ramadan (*Eid ul-Fitr*) is a major event for Muslims and is celebrated with a three day period of festivities.<sup>32</sup> Further, the end of Ramadan marks the end of a restriction on sex during daytime hours and the end of a period of piety. Nevertheless, the increase in fertility is small, statistically insignificant and perhaps skewed toward female births. We address potential selection issues that might arise from fertility changes in section 4.6.

## 4.5 Other Birth Outcomes

We briefly summarize results for nine other birth outcomes: low birth weight, infant death, APGAR scores, maternal weight gain, NICU unit, C-section, abnormal conditions, and congenital anomalies. We used three different specifications and two different samples but found that many of the results were sensitive to these choices.<sup>33</sup> Appendix Table A5 shows results with our more restricted sample of zipcodes and the broadest sample in terms of gestation (25-42 weeks) where only the first 5 gestation month coefficients are readily interpretable (due to to compositional effects). The most consistent finding concerned lower maternal weight gain due to exposure in the 7th month of gestation that is apparent in all of our samples, though not always statistically significant. While maternal

---

<sup>30</sup>Several other gestation months show larger drops for female births associated with Ramadan exposure.

<sup>31</sup>The increase in male births (one third of one percent) lags female births (9 percent). When we use our full sample of Arab births (not shown) the increase in total births is 4 percent with male births rising by 0.5 percent and female births rising by 6.5 percent.

<sup>32</sup>In Hong Kong natality microdata, the lunar new year celebration is associated with a comparable increase in births approximately nine months later (results available from authors).

<sup>33</sup>We varied gestation length (25-42 weeks, 37-42 weeks, and 39-42 weeks) and whether we used the full sample of Arabs or the more restricted sample of zipcodes.

weight gain might appear to be a desirable “first stage” measure, it has at least one (unobserved) self-reported component (pre-pregnancy weight) and thereby suffers from substantially more measurement error than newborn’s birth weight [Schieve et al., 1999].

## 4.6 Selective Timing of Conceptions Around Ramadan

Our identifying assumption is that the composition of Muslim parents does not change systematically by their children’s *in utero* exposure to Ramadan. A general concern could be that mothers of higher socioeconomic status (SES) seek to avoid having pregnancies overlap with Ramadan by concentrating conceptions during the two to three months just after Ramadan.<sup>34</sup> Another concern could be if less healthy or less educated women are more likely to conceive in the month prior to Ramadan which might account for our consistent findings on first month exposure. Finally, although we did not detect a statistically significant increase in conceptions following Ramadan, one might be concerned that there is selection associated with which parents conceive just after Ramadan due to general behavioral changes in society during this festive period.

We assess whether Ramadan exposure during pregnancy and the month prior to conception is associated with a set of pre-determined characteristics of the pregnancy that may be correlated with birth outcomes.<sup>35</sup> Table 4 estimates equation (1) with twelve “outcome” variables: mothers’ education, whether the pregnancy was paid for by Medicaid (income proxy), mother’s age, father’s age, father’s education, tobacco use during pregnancy, alcohol use during pregnancy, parity, whether a previous child was born dead, an indicator for missing father’s education, whether the mother had previously delivered a small baby and whether diabetes was considered a risk factor for the mother. Out of the 120 estimates, only 1 coefficient was significant at the 5 percent level. We found exposure during the last month of pregnancy was associated with *lower* alcohol use. We found only 4 coefficients that were significant at the 10 percent level and three of them suggested Ramadan exposure was positively selected. For example mothers who had high exposure in

---

<sup>34</sup>This might not alter our conclusions concerning differences due to exposure *within* the gestation period.

<sup>35</sup>Because we only observe those conceptions which result in a live birth, effects of post-conception Ramadan exposure may be manifested in pre-determined characteristics if Ramadan-induced fetal mortality has a gradient in these same characteristics.

the fourth month of gestation had higher education than mothers whose pregnancies did not overlap with Ramadan. Most importantly we found no evidence indicating positive selection in mothers who conceive in the month after Ramadan and no evidence suggesting that mothers who conceive in the month before Ramadan are negatively selected. In an additional check, we have run all of our birthweight and gestation length results dropping mothers who conceived in the month after Ramadan so that our effects are estimated only relative to mother's who conceived two to three months after Ramadan but whose pregnancies did not overlap with Ramadan, and found very similar results.<sup>36</sup>

## 5 Census Results

### 5.1 Results from Uganda Census

Our presentation of potential long-term effects begins with Uganda, where self-reported religion, birth month, and various health outcomes are available for a sizable number of adult Muslims and non-Muslims. As in data from other countries (e.g., the US Census), disability is the primary measure of health in the Uganda Census.

#### 5.1.1 Disability Outcomes

Table 5 shows disability outcomes for Muslims and non-Muslims. Because these outcomes have a low incidence rate we have multiplied the coefficients and standard errors by 100 to make them easier to read. The effects are therefore measured in percentage points. In the first column we show the effects of Ramadan exposure over each of the nine months preceding birth. In column (1) we find a statistically significant increase in the likelihood of a disability (of any kind) for Muslims born nine months after Ramadan (point estimate of 0.819 and  $p$ -value of 0.02). Relative to the mean disability rate of 3.8 percent, the effect is substantial at 22 percent. We find that no other month prior to birth is statistically significant and the  $p$ -value on the joint test of all nine coefficients does not approach

---

<sup>36</sup>If anything, point estimates tend to increase in absolute value. For example the Table 1 column (6) estimate for the first month of gestation rises to -53.3 from -37.9 with a  $p$ -value of 0.02. The Table A4 column (6) estimate of the 2nd month effect on weeks of gestation is -0.16 rather than -0.11 with a  $p$ -value of 0.02.

statistical significance. We cannot reject that all of the coefficients are equal.

Turning to specific disabilities (columns (2) to (5)), the most striking finding is the increased incidence of a mental or learning disability (column (4)) when Ramadan occurs during the first month pregnancy. The point estimate is 0.250 with a  $p$ -value of 0.001. Given the mean rate of 0.14 percent this implies that the occurrence of Ramadan early in pregnancy nearly doubles the likelihood of a disability related to diminished cognitive function. Thus, the increase in mental/learning disabilities from month-one Ramadan exposure would account for about 15% of all mental/learning disabilities among Muslims. Furthermore, those with exposure in month 8 have a 100% increase (significant at the 5% level) and those with Ramadan exposure in months 5 or 6 show smaller increases (significant at the 10% level). The joint test on all gestation months of no effect is rejected at the 4 percent significance level.

We also find that the incidence of sight/blindness and hearing/deafness are higher for those born 9 months after Ramadan. Specifically, the magnitude of the effects relative to those not *in utero* are 33 percent for blindness ( $p$ -value = 0.07) and 64 percent for deafness ( $p$ -value = 0.04). For hearing/deafness we also find a marginally significant effect for those exposed to Ramadan in the fifth month of gestation.

We run the same specifications on our sample of non-Muslims in columns (6) - (10). We find no cases of a corresponding significant result for Muslims also occurring for Non-Muslims for these outcomes. We tested the sensitivity of the results for Muslims to also including exposure during the 10th month prior to birth and found that the results were unaffected and that in no case was the coefficient on the 10th month statistically significant (see Appendix Table A6). We also ran our specifications separately for men and women (not shown) and found that the results were qualitatively similar though the estimates were much less precise.

### 5.1.2 Causes of Disability

Previous falsification tests have considered Ramadan exposure outside of pregnancy and Ramadan exposure during pregnancy for non-Muslims. Information on the causes of disabilities provides a third falsification test. We group these reported causes – accident,

occupational injury, war injury, aging, disease, or congenital – by whether they can reasonably be linked to fasting via the mechanisms discussed earlier. Clearly, disabilities that arise from accidents, occupational injuries, or war injuries are postnatal and should not be related to maternal fasting during Ramadan. On the other hand, the fetal origins hypothesis suggests that extended periods of nutritional restriction may be associated with a reprogramming of the body’s systems that result in poor health outcomes later in life (see Appendix for additional discussion). This would be consistent with those who report “aging” as the source of a disability. Since it is conceivable that fasting might contribute to a weakened immune system, respondents who report disabilities due to “disease” could plausibly be related to the timing of Ramadan. Finally, whether maternal nutrition affects congenital disabilities (those present at birth) is not clear-cut.<sup>37</sup>

We find no significant effects from accidents, occupational injury or war injuries for Muslims or non-Muslims in any gestation month (Appendix Table A7). In contrast, Muslims born nine months after Ramadan have an increased incidence of disabilities due to aging of 0.37 percentage points ( $p$ -value = 0.006). We find no evidence linking Ramadan exposure to disease-related or congenital disabilities (consistent with Michigan results for congenital anomalies). We found no comparable effect of first month exposure to Ramadan on disabilities caused by aging for non-Muslims.<sup>38</sup>

In order to address possible concerns about selective timing of pregnancy in Uganda, we used a sample of children aged 17 or under and living with their parents and regressed parent characteristics (education, illiteracy, and disability) on the child’s Ramadan exposure using equation (1). As with Michigan, we found no statistically significant effects of negative selection on parent characteristics. Finally, we also found that the results were insensitive to excluding outlier cohorts that had extremely large or small disability rates. If anything, excluding outliers slightly increased the point estimates and their precision.

---

<sup>37</sup>If the disability is epigenetic then it may be associated with maternal fasting.

<sup>38</sup>Among non-Muslims the only significant effect is that those exposed to Ramadan one month before birth are 0.12 percentage points ( $p$ -value = 0.017) more likely to have a congenital disability. This is a 20 percent effect relative to the mean.

### 5.1.3 Sex Composition of Adult Population

With the Uganda data we explore the possibility that maternal fasting may influence the sex composition of the adult population. This could arise either from alterations to the sex composition at birth or because of selective mortality by sex after birth as implied by some of the fetal origins literature (see Appendix Section 1). To assess this, we conduct an analysis parallel to our Michigan analysis. First we simply regress male as an outcome in equation (1). Second, we aggregate the population by cells constructed by birth month both for the pooled sample as well as separately by sex and take the log of the population counts as an outcome.

Results are shown in the left most panel of Table 6. In column (1) we find that every implied gestational month has a negative coefficient and that the 1st, 4th and 7th months of gestation are statistically significant at the 5 percent level. The joint test of all the exposure months is significant at the 10 percent level. In column (2) we find only slight evidence cohort size is related to Ramadan exposure when we pool men and women. When we look at the log of population counts of males in column (3), seven of the nine months have negative coefficients and the 7th month of gestation has a particularly large and statistically significant effect (15%). The effects on the sex in column (1) appear to be driven by reductions in the number of males. In column (4) we show the analogous results for women where the effects are all positive but only significant in one month. In other results (not shown) we find no comparable effects on the sex composition for non-Muslims.

### 5.1.4 Other Outcomes in Uganda

The remaining columns of Table 6 show results for non-health outcomes. Unfortunately preferred economic outcomes, such as wages, income, and wealth, are not available. In column (5) we examine whether home ownership, a proxy for wealth, is affected. We restrict the sample to men since they are the vast majority of property owners in Uganda.<sup>39</sup> We find that men exposed to Ramadan in the first month of gestation are 2.6 percentage

---

<sup>39</sup>Uganda is a patriarchal society where land is passed down through sons. Although women are not prevented from owning land, by one estimate, 93 percent of Ugandan land is owned by men. (<http://www.womensenews.org/article.cfm/dyn/aid/1456/context/archive>).



points less likely to own their home ( $p$ -value=0.027) and that men exposed in the 2nd month of gestation are 2.1 percentage points less likely to own their home ( $p$ -value=0.051). Given the high rate of male home ownership (73.4 percent), these effects are not especially large. We can reject that there is no effect of Ramadan exposure over all gestation months on home ownership at the 5 percent level. In contrast, we find no statistically significant effects of Ramadan exposure on home ownership for non-Muslims.

In columns (6) through (9) we examine illiteracy, completed years of schooling, a dummy for no schooling, and employment status at the time of the Census. We find no statistically significant effects that associate greater Ramadan exposure with higher illiteracy or lower schooling. In fact those born 8 months after Ramadan appear to have *higher* human capital levels by both of these measures. The magnitude of these effects, however, is small. For example, the increase in years of schooling for these individuals is only about a tenth of a year, or 1.6 percent of the sample mean. When we split the sample by gender, we only found these positive effects for men so we speculate that these results might reflect a selective effect on surviving males, who seem to bear the brunt of Ramadan-related attrition. In developing countries where average mortality rates are high, the selective effect of health disruptions operating through early-life mortality may overwhelm “fetal origins” effects on adult survivors [Deaton et al., 2009]. Those born 5 months after Ramadan have lower employment rates of about 1.8 percentage points.

## 5.2 Results from Iraq Census

We replicate the basic Uganda results using 1997 Iraq Census data. Columns (1) to (9) of Table 7 show the effects on disability. Full exposure to Ramadan nine months before birth is associated with a 0.3 percentage point increase in the probability of having a disability ( $p$ -value =0.025). While in Uganda the overall disability rate was 3.8 percent, in Iraq it is just 1.5 percent. However, the effect size relative to the mean in Iraq is about 19 percent, comparable to the 22 percent effect size that we estimated in Uganda. In Iraq the rates of disabilities involving sight and hearing, however, are a much smaller fraction of the reported rates for Uganda and this may explain why we detect no effect on these

measures in columns (2) and (3).<sup>40</sup>

“Insane” is the sole mental disability queried, which IPUMS relabeled as “psychological” disability. Interestingly, at 0.35 percent, Iraq’s psychological disability rate is actually higher than the combined rate of 0.28 percent for mental/learning *plus* psychological disabilities in Uganda (despite Iraq’s lower overall disability rate). This suggests that mental/learning disabilities that are related to cognitive impairments may be subsumed in the psychological disability measure for Iraq. In column (4) we find strong effects on psychological disabilities just as we did for mental/learning disabilities in Uganda. First month exposure to Ramadan is associated with 0.19 percentage point increase in the likelihood of a psychological disability or 56 percent effect relative to the mean ( $p$ -value =0.002). We also estimate positive but insignificant effects in 7 of the other 8 gestation months. As was the case in Uganda with mental/learning disabilities, the joint test of zero effect across all gestation months is easily rejected at the 5 percent level, as is the test of equality of coefficients. The fact that both overall disability as well as disabilities that likely capture cognitive impairments appear to be impacted in precisely the same period of fetal development in two different societies is remarkable and reinforces evidence that our findings are not due to chance.

In columns (5) through (9) of Table 7 we turn to socioeconomic outcomes. The 1997 Iraqi Census asks about instances of men having multiple wives which we use to proxy for wealth (as described earlier). For this measure, shown in column (5) we find strong effects throughout much of fetal development: the joint test of all gestation month exposure coefficients is significant at the 8 percent level. Men with first month exposure are more than half a percentage point ( $p$ -value =0.07) less likely to have multiple wives. Strong effects are also found during months 4, 5 and 6 of gestation. Similarly, for home ownership (column 6), we see highly significant effects of exposure throughout the *in utero* period and the joint test of all gestation month coefficients is significant at the 2 percent level.

In column (7) we see no effects on the sex composition of the adult population. Many

---

<sup>40</sup>For vision/blindness only 0.14 percent report this disability which is only about one-tenth of the share reporting a comparable disability in Uganda. For deaf/hearing only 0.02 percent report this disability which is only one-sixteenth of the rate found in Uganda.

gestation months actually show positive coefficients but none are significant nor are they jointly significant. There is suggestive evidence that Ramadan exposure affects the likelihood of completing primary school or less (column 8). Three coefficients are significant at the 10 percent level but the implied effects sizes are only around 2 percent. Finally, in column (9) we find tiny positive (perverse) effects of Ramadan exposure on employment that are consistently significant. We note that among males, home owners are less likely to be employed (73%) than non-home owners (82%) suggesting that employment may be a poor proxy for economic status in Iraq. Further, controlling for home ownership and multiple wives eliminates the effects of Ramadan exposure on employment for men.

As with our Uganda results, we have also run all of these estimates including exposure 10 months prior to birth and in no case did it meaningfully alter the results.

### 5.3 Results from US Census

The US data provide potentially higher quality measures of disability as well as direct measures of labor market outcomes (unavailable for Iraq and Uganda). As mentioned in Section 3.2.3, the data's chief drawback is absence of month of birth information, which dulls assignment of the Ramadan ITT. On the other hand, findings of long-term effects in the US may address concerns that the Uganda and Iraq results do not generalize to Muslims living in a high-income Western society.

Panel A of Table 8 presents results for four outcomes across three samples. In each regression we estimated effects of Ramadan exposure in one's birth quarter and in the prior two birth quarters, omitting Ramadan exposure three quarters prior to the birth quarter and thereby treating that cohort as the control group. Since we have averaged our monthly exposure measures within each birth quarter, our quarterly exposure measure never exceeds one third (0.33) in any quarter. Therefore, our coefficients which represent the effect of a unit change in exposure should be scaled down by one-third to be interpretable as the effect of fasting for 30 days within the birth quarter. For this table we have also multiplied all the dichotomous outcomes by 100.

The first four columns pool samples drawn from the 1980 Census and the 2005-2007 ACS. We find a significant effect of exposure in the quarter prior to birth on the incidence

of disability (column (1)). The coefficient of 3.50 suggests that Ramadan's occurrence in a particular month within that quarter would raise the likelihood of a disability by about 1.17 percentage points. Since the mean rate of disability in the sample is 6.07 percent, this implies about a 19 percent effect at the mean, again quite similar to the corresponding magnitudes in Uganda and Iran. The  $p$ -value on the joint test of the three quarters is 0.015. However, the timing of exposure appears to be concentrated more in the second trimester rather than early in gestation.<sup>41</sup>

In column (4), we find that exposure both in the birth quarter and two quarters prior to birth is associated with a reduction in log annual earnings of about 5.5 to 6 percent (rescaled). Both estimates are significant at the 5 percent level and the effects are jointly significant at the 0.11 level. To measure human capital we began by examining completed years of schooling as an outcome (not shown) and found a statistically significant effect of about 0.15 years from one month's exposure that occurs two quarters prior to birth. Since this was a relatively small sized effect, we then looked for whether this was concentrated at the low end of the education distribution by looking at failure to complete 12 years of schooling as an outcome ("less than HS"). Column (2) shows a coefficient of 2.513 for Ramadans falling two quarters prior to the birth quarter, significant at the 11 percent level. The point estimate implies that one month of exposure to Ramadan fasting is associated with about a 0.84 percentage point increase in the high school dropout rate which implies a 6.4 percent effect size evaluated at the mean.

We find negative point estimates for the likelihood of being male (column (3)) that are not statistically significant, but nevertheless potentially meaningful quantitatively. Complete exposure in the quarter two quarters prior to the birth quarter is associated with a roughly 1 percentage point decline in the male share of the adult Muslim population.

In columns (5) through (8) we restrict the sample to those observed in the 1980 Census, about a third of the full sample, and observe broadly similar effects. With this sample we also observe that the other two quarters of Ramadan appear to have strong effects on adult disability. In columns (9) through (12) we restrict the sample to the pooled

---

<sup>41</sup>Depending on when in the quarter an individual is born, the prior quarter could reflect effects anywhere from the 4th gestation month to the 9th. If births were uniformly distributed within a quarter then the mean exposure month would be the fifth month of gestation.

ACS sample. We again see disability effects of a similar magnitude and timing. For both subsamples the effects on earnings are insignificant but if anything appear to be stronger earlier in pregnancy (two quarters prior to the birth quarter).

Panel B takes a more conservative approach in designating the control group. Although we know that the vast majority of pregnancies that overlap with Ramadan's occurrence in the two quarters prior to birth will actually be exposed to Ramadan for the entirety of the three months of the quarter, this is much less likely for the quarter of birth itself and the quarter three quarters prior to the birth quarter. Therefore, here both of these quarters are omitted and serve as the reference group. As can be seen in columns (1) through (4) of Panel B, this does not alter our basic results. The effect size on disability is a bit smaller but still highly significant. The effect on having fewer than 12 years of schooling actually rises, becomes significant at the 5 percent level, and the joint test of both exposure coefficients is now significant at the 8 percent level. On the other hand the effect on earnings drops considerably and is no longer statistically significant.

We have also run all of these specifications on a sample of US born whites that are presumably non-Muslims based on reported ancestry and found no effects of similar magnitudes.

To summarize, in the US Census we find: 1) strong evidence of elevated rates of disability from *in utero* exposure to Ramadan toward the middle of pregnancy; 2) evidence of lower earnings with Ramadan exposure early in pregnancy (e.g., first trimester); and 3) suggestive evidence linking Ramadan exposure to a lower likelihood of completing high school.

## 6 Discussion and Future Research

As rates of fasting by pregnant women during Ramadan approach unity, our ITT estimate approaches the treatment effect of fasting (which cannot be said of previous comparisons between fasters and non-fasters). Fasting observance may be highest in early pregnancy, both because mothers may be unaware they are pregnant and the burden of pregnancy

is lower.<sup>42</sup> Thus, the estimated health damage attributable to Ramadan falling in the first month of pregnancy may approximate the treatment effect of fasting during this period. Correspondence between our ITT estimate and fasting’s effect is likely higher in Iraq and Uganda where we have little classification error in Muslim status. In our Michigan data, our proxy for Muslims will include a higher fraction of non-Muslims due to the likely presence of Chaldeans who report Arab ancestry. This is also likely to be the case, albeit to a lesser extent, with our US Census data where we may include some non-Muslim immigrants from predominantly Muslim countries. As compliance (fasting during Ramadan) is presumably zero for non-Muslims, our US ITT estimates are likely attenuated.

Ideally, we would observe fasting behavior by month of pregnancy and subsequent health or human capital outcomes for a large sample of Muslims. With this information and a sufficiently long span of birth years, we could construct Wald estimates of the effect of fasting on health during each pregnancy month. Ramadan’s coincidence with pregnancy month would be the binary instrumental variable for fasting observance. As long as Muslims are not fasting for other reasons during the month of Ramadan (as seems reasonable), this Wald estimate could be interpreted as the effect of fasting on fasters (i.e., the treatment on the treated rather than simply a *LATE* estimate, see Angrist and Pischke [2009]). Failing this, data on fasting behavior and pregnancy month could be used to estimate the first stage effects of Ramadan timing (preferably for the US, Uganda, or Iraq), and combined with our ITT estimates in a two-sample IV procedure. This approach would integrate potential heterogeneity in fasting rates by pregnancy month.

The most compelling previous studies of fetal origins have relied on exogenous shocks “caused by conditions outside the control of the mother” [Currie, 2009]. Our study departs from these in considering a treatment that to a greater degree is within the control of the mother (but still identified by exogenous timing). That obtaining a dispensation to postpone fasting until after pregnancy is apparently the exception rather than the norm (see Appendix A.1.2) suggests two possibilities. First, the cost of requesting the dispen-

---

<sup>42</sup>The only study that we are aware of that documented differences in fasting behavior across pregnancy was by Arab and Nasrollahi [2001] who found that of the 4,343 women delivering in hospitals in Hamadan, Iran in 1999, fasting was only slightly more common when Ramadan fell in the first trimester (77% ) than in the second trimester (72%) or third trimester (65%).

sation may be high – in part because mothers usually become aware of their pregnancies after the first month [Floyd et al., 1999]. Alternatively, it may be that the full health consequences of Ramadan fasting during pregnancy are unknown. This explanation also seems plausible as ours is the first study to find long-term effects.

An alternative approach families could adopt is to time pregnancies to commence shortly after Ramadan, and thereby avoid the overlap. That we do not observe this behavior could suggest that timing pregnancies is costly or unreliable,<sup>43</sup> or again that fasting during pregnancy is not considered teratogenic.

Future research should assess whether other commonly-experienced disruptions to prenatal nutrition exert similar effects as Ramadan fasting. Most US pregnancies are not recognized until after the first month of gestation [Floyd et al., 1999]. Given the results of this study, maternal behavior during the first month of pregnancy can have permanent impacts on offspring health. Roughly 40% of US women of childbearing age are attempting to lose weight [Cohen and Kim, 2009] and 24% of women reported meal-skipping during pregnancy [Siega-Riz et al., 2001].<sup>44</sup> Even in relatively well-nourished populations, prenatal nutrition (and at a minimum its timing) may be sub-optimal for fetal development. Future research should employ new identification strategies to evaluate both short and long-term health effects of nutrition in early pregnancy on health and other end points, e.g., test scores.

## References

- Douglas Almond, Kenneth Y. Chay, and David S. Lee. The costs of low birth weight. *The Quarterly Journal of Economics*, 120(3):1031–1084, August 2005.
- Douglas Almond, Lena Edlund, Michael Joffe, and Mårten Palme. Is hyperemesis gravidarum an adaptive condition? manuscript, Stockholm University, February 2009.
- Joshua D. Angrist and Jörn-Steffen Pischke. *Mostly Harmless Econometrics: An Empiricist's Companion*. Princeton University Press, Princeton, New Jersey, 2009.

---

<sup>43</sup>Dickert-Conlin and Chandra [1999] found a responsiveness to tax incentives in the timing of deliveries, not conceptions.

<sup>44</sup>Furthermore, approximately 5% of pregnant women manifest eating disorders [Turton et al., 1999].

- Malihe Arab and Shahla Nasrollahi. Interrelation of ramadan fasting and birth weight. *Medical Journal of the Islamic Academy of Sciences*, 14(3):91–95, 2001.
- Fereidoun Azizi, Hossein Sadeghipour, Behnam Siahkollah, and Nasrollah Rezaei-Ghaleh. Intellectual development of children born to mothers who fasted during pregnancy. *International Journal of Vitamin and Nutrition Research*, 74(5):374–380, September 2004.
- Kasey Buckles and Daniel M. Hungerman. Season of birth and later outcomes: Old questions, new answers. manuscript, University of Notre Dame, July 2008.
- Elissa Z. Cameron. Facultative adjustment of mammalian sex ratios in support fo the trivers-willard hypothesis: evidence for a mechanism. *Proceedings of the Royal Society B: Biological Sciences*, 271(1549):1723–1728, 2004.
- Jennifer H. Cohen and Hyoshin Kim. Sociodemographic and health characteristics associated with attempting weight loss during pregnancy. *Preventing Chronic Disease: Public Health Research, Practice, and Policy*, 6(1):1–9, January 2009. Centers for Disease Control and Prevention.
- Dora L. Costa and Joanna N. Lahey. Predicting older age mortality trends. *Journal of the European Economic Association*, 3(2-3):487–493, April-May 2005.
- Dora L. Costa, Lorens A. Helmchen, and Sven Wilson. Race, infection, and arteriosclerosis in the past. *Proceedings of the National Academy of Sciences*, 104(33):13219–13224, 2007.
- Flavio Cunha and James J. Heckman. The technolgy of skill formation. *American Economic Review*, 97(2):31–47, May 2007.
- Janet Currie. Healthy, wealthy, and wise: Is there a causal relationship between child health and human capital development? *Journal of Economic Literature*, XLVII(1): 87–122, March 2009.
- Angus Deaton, Carlos Bozzoli, and Climent Quintana-Domeque. Child mortality and adult height. *Demography*, forthcoming 2009.



- Stacy Dickert-Conlin and Amitabh Chandra. Taxes and the timing of births. *Journal of Political Economy*, 1999, vol. 107, no. 1, 107(1):161–177, 1999.
- E. Dikensoy, O. Balat, B. Cebesoy, A. Ozkur, H. Cicek, and G. Can. The effect of Ramadan fasting on maternal serum lipids, cortisol levels and fetal development. *Archives of gynecology and obstetrics*, 279(2):119, 2009.
- Gabriele Doblhammer and James W. Vaupel. Lifespan depends on month of birth. *Proceedings of the National Academy of Sciences*, 98(5):2934–2939, February 2001.
- Lena Edlund. Son preference, sex ratios, and marriage patterns. *Journal of Political Economy*, 107(6):1275–1304, December 1999.
- Reyn Van Ewijk. Long-term health effects on the next generation of ramadan fasting during pregnancy. April 2009. CEP Discussion Paper No 926.
- RL Floyd, P Decoufle, and DW Hungerford. Alcohol use prior to pregnancy recognition. *American Journal of Preventive Medicine*, 17:101–107(7), August 1999.
- Amita Kapoor, Elizabeth Dunn, Alice Kostaki, Marcus H. Andrews, and Stephen G. Matthews. Fetal programming of hypothalamo-pituitary-adrenal function: Prenatal stress and glucocorticoids. *Journal of Physiology*, 572(1):31–44, 2006.
- Zohreh Kavehmanesh and Hassan Abolghasemi. Maternal ramadan fasting and neonatal health. *Journal of Perinatology*, 24:748–750, 2004.
- Sebastian Kraemer. The fragile male. *BMJ*, 321(7276):1609–1612, 2000.
- Ashok Malhotra, P.H. Scott, J.S. Scott, H. Gee, and B.A. Wharton. Metabolic changes in asian muslim pregnant mothers observing the ramadan fast in britian. *British Journal of Nutrition*, 61(3):663–672, May 1989.
- Fiona Mathews, Paul J. Johnson, and Andrew Neil. You are what your mother eats: evidence for maternal preconception diet influencing foetal sex in humans. *Proceedings of the Royal Society B*, pages 1–8, April 2008.

- P. J. Meis and J.C. Rose and M. Swain. Pregnancy alters diurnal variation of plasma glucose concentration. *Chronobiol. Int.*, 1:145–149, 1984.
- Boyd E. Metzger, Veronica Ravnkar, Rita A. Vileisis, and Norbert Freinkel. ‘accelerated starvation’ and the skipped breakfast in late normal pregnancy. *The Lancet*, 1(8272): 588–592, March 1982.
- Hisham M. Mirghani, Sarath Weerasinghe, Shamsaa Al-Awar, Lolowa Abdulla, and Mutairu Ezimokhai. The effect of intermittent maternal fasting on computerized fetal heart tracing. *Journal of Perinatology*, 25:90–92, 2005.
- H.M. Mirghani, S.D. Weerasinghe, J.R. Smith, and M. Ezimokhai. The effect of intermittent maternal fasting on human fetal breathing movements. *Journal of Obstetrics and Gynaecology*, 24(6):635–637, September 2004.
- Institute of Medicine, editor. *Nutrition During Pregnancy and Lactation: An Implementation Guide*. National Academy Press, Washington D.C., 1992.
- M Pearl, ML Wier, and M Kharrazi. Assessing the quality of last menstrual period date on california birth records. *Paediatric and perinatal epidemiology*, 21 Suppl 2:50, 2007.
- AM Prentice, A Prentice, WH Lamb, PG Lunn, and S. Austin. Metabolic consequences of fasting during ramadan in pregnant and lactating women. *Human nutrition. Clinical nutrition*, 37(4):283–94, July 1983.
- Laura A. Schieve, Geraldine S. Perry, Mary E. Cogswell, Kelley S. Scanlon, Deborah Rosenberg, Suzan Carmichael, and Cynthia Ferre. Validity of self-reported pregnancy delivery weight. *American Journal of Epidemiology*, 150(9), 1999.
- Anna Maria Siega-Riz, Tracy Hermann, David A Savitz, and John M Thorp. Frequency of eating during pregnancy and its effect on preterm delivery. *American Journal of Epidemiology*, 153(7):647–652, 2001.
- P. Turton, P. Hughes, H. Bolton, and P. Sedgwick. Incidence and demographic correlates of eating disorder symptoms in a pregnant population. *The International Journal of Eating Disorders*, 26(4):448–452, 1999.

**Table 1: Effects of Ramadan Exposure on Birth Weight by Gestation Month, Michigan Arabs**

Dependent Variable is Birthweight, Coefficients are on Ramadan Exposure measures (% of daylight hours, % of days)

Months prior to Birth	<i>Ramadan Exposure Assuming Normal Gestation</i>					<i>Ramadan Exposure Using Gestation Data (37-42 wks)</i>				
	(1) Birth Day Hours	(2) Birth Day Hours	(3) Birth Day Days	(4) Birth Day Days	(5) Birth Mo. Days	Gestation Month exposure	(6) Birth Day Hours	(7) Birth Day Hours	(8) Birth Day Days	(9) Birth Day Days
10	--	6.4 (19.7)	--	4.3 (13.8)	--	0	--	6.0 (18.1)	--	5.5 (12.4)
9	-39.8 ** (18.5)	-38.9 ** (18.7)	-26.7 ** (12.7)	-25.9 ** (13.0)	-10.8 (12.7)	1	-37.9 ** (16.9)	-36.9 ** (17.1)	-22.3 * (11.5)	-21.3 * (11.7)
8	-38.0 ** (17.7)	-35.7 * (19.1)	-24.5 ** (12.4)	-22.8 * (13.6)	-33.2 *** (12.4)	2	-27.2 * (16.1)	-25.0 (17.5)	-16.9 (11.1)	-14.7 (12.2)
7	7.6 (18.3)	9.6 (19.2)	7.5 (12.8)	9.0 (13.6)	-8.8 (12.8)	3	-4.7 (16.6)	-2.8 (17.6)	-1.7 (11.5)	0.1 (12.3)
6	-37.6 ** (18.5)	-35.6 * (19.5)	-25.4 ** (13.0)	-23.9 * (13.9)	-10.8 (12.8)	4	-13.4 (16.6)	-11.5 (17.7)	-8.9 (11.6)	-6.9 (12.5)
5	1.1 (18.8)	3.0 (19.7)	2.5 (13.2)	4.0 (14.0)	-18.5 (13.2)	5	-16.0 (16.9)	-14.1 (17.8)	-9.8 (11.9)	-7.9 (12.6)
4	-33.8 * (18.7)	-32.0 * (19.5)	-23.6 * (13.2)	-22.2 (13.9)	-8.2 (13.2)	6	-21.9 (17.0)	-20.2 (17.8)	-14.8 (11.9)	-13.0 (12.6)
3	-19.7 (19.1)	-17.9 (19.9)	-13.0 (13.0)	-11.6 (13.7)	-16.2 (13.3)	7	-17.3 (16.7)	-15.6 (17.5)	-14.1 (11.7)	-12.4 (12.4)
2	-13.5 (18.6)	-11.9 (19.3)	-7.6 (12.5)	-6.3 (13.1)	-12.2 (12.6)	8	12.4 (16.0)	14.0 (16.7)	8.1 (11.2)	9.7 (11.8)
1	10.1 (19.0)	12.3 (20.1)	7.8 (12.8)	9.5 (13.9)	-5.9 (13.0)	9	-15.5 (16.4)	-13.4 (17.6)	-10.7 (11.4)	-8.6 (12.5)
joint test, coefficients on months 1 to 9 equal to 0										
<i>p</i> -value	0.078	0.100	0.251	0.076	0.478		0.289	0.350	0.437	0.513
joint test, coefficients on months 1 to 9 are equal										
<i>p</i> -value	0.166	0.251	0.280	0.121	0.844		0.405	0.396	0.557	0.518
<i>N</i>	44112	44112	32101	44112	44112		41106	41106	41106	41106

*Notes:* Columns 1-4 use exact birth date but do not utilize gestation data. Column 5 only uses birth month. Columns 6-10 use either self reported or physician estimated gestation (see text). Entries show the coefficient on the relevant Ramadan exposure measure. Controls include mother's age, mother's age squared, mother's education, tobacco use, alcohol use, parity, father's education, dummy for missing father's education, father's age, father's age squared, number of previous pregnancies that resulted in death at birth, conception (or birth) month dummies, county dummies and birth year dummies. Italicized estimates may represent postnatal exposure for some sample members.

Standard errors in parentheses, \*significant at 10%; \*\* significant at 5%; \*\*\* significant at 1%

**Table 2: Effects of Ramadan Exposure on Birth Weight by Gestation Month, Michigan Non-Arabs**

Months prior to Birth	<i>Exposure Assuming Normal Gestation</i>						<i>Exposure Using Gestation Data</i>			
	<i>All Non Arabs</i>			<i>Exc. Arab Zip</i>			<i>All Non Arabs</i>	<i>Ex Arab Zip</i>		
	(1) Birth Day Hours	(2) Birth Day Days	(3) Birth Month Days	(4) Birth Day Hours	(5) Birth Day Days	(6) Birth Month Days	Gestation Month exposure	(7) Birth Day Hours	(8) Birth Day Days	(9) Birth Day Hours
9	-3.3 (3.3)	-2.2 (2.3)	-1.4 (2.4)	14.1 (11.9)	11.3 (8.3)	4.8 (8.3)	1	-3.2 (3.3)	-0.4 (2.3)	21.8 (11.9) *
8	-0.5 (3.2)	-0.8 (2.3)	-2.3 (2.3)	8.4 (11.4)	4.6 (8.1)	9.7 (8.0)	2	2.5 (3.2)	3.0 (2.2)	10.9 (11.2)
7	-2.9 (3.3)	-2.5 (2.3)	-4.0 (2.3) *	9.3 (11.7)	7.2 (8.3)	3.2 (8.3)	3	-1.6 (3.2)	-1.1 (2.3)	16.1 (11.5)
6	0.5 (3.3)	-0.1 (2.3)	0.5 (2.3)	5.3 (11.6)	4.7 (8.3)	13.5 (8.3)	4	-0.8 (3.2)	0.1 (2.3)	10.1 (11.4)
5	-3.7 (3.3)	-3.2 (2.4)	-4.2 (2.4) *	8.8 (11.8)	7.1 (8.4)	-4.4 (8.3)	5	0.6 (3.2)	0.7 (2.3)	5.8 (11.5)
4	1.0 (3.3)	0.6 (2.4)	-1.9 (2.3)	2.9 (11.8)	2.5 (8.5)	11.7 (8.4)	6	0.1 (3.2)	0.8 (2.3)	5.5 (11.5)
3	-3.8 (3.3)	-2.7 (2.3)	0.5 (2.3)	9.6 (11.8)	7.1 (8.3)	2.9 (8.4)	7	-0.8 (3.2)	0.5 (2.3)	-2.2 (11.3)
2	-0.8 (3.2)	-1.5 (2.2)	-3.5 (2.2)	8.8 (11.5)	7.2 (8.0)	9.1 (8.0)	8	-1.4 (3.1)	-1.1 (2.2)	8.5 (11.0)
1	0.8 (3.3)	-0.2 (2.3)	-1.4 (2.3)	-4.5 (11.9)	-6.2 (8.3)	-3.3 (8.4)	9	-4.8 (3.2)	-3.1 (2.3)	-2.5 (11.5)
joint test, coefficients on months 1 to 9 equal to 0										
<i>p</i> -value	0.933	0.921	0.561	0.943	0.834	0.703	0.857	0.747	0.677	
joint test, coefficients on months 1 to 9 are equal										
<i>p</i> -value	0.934	0.958	0.743	0.979	0.892	0.783	0.813	0.655	0.768	
<i>N</i>	2E+06	2E+06	2E+06	122844	122844	122844	1E+06	1E+06	84675	

*Notes:* All entries are coefficients on Ramadan exposure to daylight hours over subsequent 30 days as fraction of peak daylight hours. Columns 1 and 2 assume normal gestation. Columns 3 and 4 use either self reported or physician estimated gestation and use births between 39 and 42 weeks gestation. Columns 1 and 3 use a broad sample of imputed non-Arab mothers based on reported ancestry or race. Columns 2 and 4 further restrict this sample to mothers living in zipcodes with no Arabs according to the 2000 Census. Controls include mother's age, mother's age squared, mother's education, tobacco use, alcohol use, parity, father's education, dummy for missing father's education, father's age, father's age squared, number of previous pregnancies that resulted in death at birth, conception (or birth) month dummies, county dummies and birth year dummies. Standard errors in parentheses, \*significant at 10%; \*\* significant at 5%; \*\*\* significant at 1%

**Table 3: Effects of Ramadan Exposure on Sex at Birth and Live Births, Michigan Arabs and Non Arabs**

Gestation Month exposure	<i>Dependent Variable is Male</i>				<i>Dependent Variable is Log Live Births (Total, Male Female)</i>					
	(1)	(2)	(3)	(4)	Arab Sample			Non-Arab Sample		
	All Arabs	Arabs Exc. Chaldean Zip	All Non-Arabs	Non-Arab Zipcodes	Total	Male	Female	Total	Male	Female
0	-0.032 (0.021)	-0.033 (0.025)	0.003 (0.003)	0.007 (0.012)	0.063 (0.068)	0.003 (0.087)	0.090 (0.097)	0.030 (0.040)	0.050 (0.049)	0.044 (0.047)
1	-0.037* (0.020)	-0.066*** (0.023)	0.002 (0.003)	0.011 (0.012)	-0.164** (0.063)	-0.258*** (0.082)	-0.043 (0.087)	0.042 (0.035)	0.051 (0.044)	0.031 (0.042)
2	-0.008 (0.020)	-0.008 (0.024)	0.001 (0.003)	-0.019 (0.012)	-0.007 (0.066)	0.017 (0.086)	0.021 (0.093)	-0.016 (0.041)	-0.005 (0.049)	0.022 (0.048)
3	-0.002 (0.020)	-0.007 (0.024)	0.002 (0.003)	-0.009 (0.012)	-0.088 (0.065)	-0.143* (0.083)	-0.060 (0.093)	0.029 (0.039)	-0.010 (0.048)	0.076 (0.046)
4	-0.022 (0.020)	-0.030 (0.024)	0.002 (0.003)	-0.007 (0.012)	0.047 (0.070)	-0.037 (0.090)	0.084 (0.096)	0.045 (0.042)	0.054 (0.049)	0.066 (0.047)
5	0.005 (0.020)	-0.015 (0.024)	0.001 (0.003)	0.014 (0.012)	-0.035 (0.069)	-0.055 (0.084)	0.004 (0.097)	0.023 (0.038)	0.032 (0.048)	0.021 (0.046)
6	-0.013 (0.020)	-0.022 (0.024)	0.001 (0.003)	-0.011 (0.012)	0.037 (0.066)	-0.021 (0.085)	0.078 (0.090)	-0.012 (0.038)	-0.018 (0.048)	0.024 (0.045)
7	-0.012 (0.020)	-0.009 (0.023)	0.002 (0.003)	0.018 (0.012)	-0.113 (0.069)	-0.083 (0.084)	-0.148 (0.094)	0.012 (0.039)	0.039 (0.048)	-0.014 (0.047)
8	-0.026 (0.019)	-0.031 (0.023)	0.001 (0.003)	-0.005 (0.011)	0.008 (0.063)	-0.063 (0.078)	0.088 (0.087)	0.056 (0.035)	0.038 (0.045)	0.077* (0.042)
9	<i>-0.030</i> (0.020)	<i>-0.030</i> (0.024)	<i>0.003</i> (0.003)	<i>0.001</i> (0.012)	<i>0.003</i> (0.068)	<i>-0.063</i> (0.083)	<i>0.060</i> (0.097)	<i>0.007</i> (0.039)	<i>0.020</i> (0.049)	<i>0.006</i> (0.046)
joint test, coefficients on months 1 to 9 equal to 0										
p-value	0.52	0.34	1.00	0.21	0.19	0.34	0.63	0.69	0.80	0.58
N	41153	29955	1434014	112516	216	216	216	216	216	216

*Notes:* All entries are coefficients on Ramadan exposure to daylight hours over subsequent 30 days as fraction of peak daylight hours during sample period. Gestation is either self reported or physician estimated. All samples use births with gestation between 37 and 42 weeks and include controls for mother's age, mother's age squared, mother's education, tobacco use, alcohol use, parity, father's education, dummy for missing father's education, father's age, father's age squared, number of previous pregnancies that resulted in death at birth, conception month dummies, county dummies and birth year dummies. The Arab sample in columns 5 through 7 drops Arab mothers living in zipcodes with a large share of Chaldeans. The non-Arab sample in columns 8 through 10 drops presumed Non Arab mothers living in zipcodes with an Arab presence. Standard errors in parentheses. *Italicized estimates may represent postnatal exposure for some sample members.* \*significant at 10%; \*\* significant at 5%; \*\*\* significant at 1%

**Table 4: Effects of Ramadan Exposure on Characteristics of Pregnancies Resulting in Live Births, Michigan Arabs**

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
Gestation Month Exposure	Mother's Education	Medicaid	Mother's Age	Father's Age	Father's Education	Tobacco	Alcohol	Parity	Previous Child Born Dead	Father's Educ. Miss.	Previous Small Baby	Diabetes Risk Factor
0	0.030 (0.138)	0.001 (0.019)	-0.111 (0.226)	-0.256 (0.261)	-0.007 (0.134)	0.001 (0.008)	-0.002 (0.002)	0.040 (0.060)	-0.029 (0.026)	-0.007 (0.010)	0.001 (0.003)	-0.008 (0.006)
1	0.014 (0.130)	-0.008 (0.018)	0.269 (0.213)	-0.011 (0.246)	0.050 (0.126)	-0.003 (0.007)	0.002 (0.001)	0.002 (0.057)	-0.035 (0.024)	-0.005 (0.009)	-0.003 (0.003)	-0.006 (0.006)
2	-0.099 (0.133)	0.008 (0.018)	0.194 (0.217)	0.080 (0.251)	-0.085 (0.129)	0.006 (0.007)	-0.001 (0.001)	-0.033 (0.058)	-0.028 (0.025)	0.018* (0.009)	-0.003 (0.003)	-0.009 (0.006)
3	0.079 (0.133)	-0.017 (0.018)	0.089 (0.219)	-0.172 (0.253)	0.135 (0.130)	-0.005 (0.007)	-0.001 (0.002)	0.021 (0.058)	-0.049* (0.025)	-0.013 (0.009)	-0.006* (0.003)	0.003 (0.006)
4	0.234* (0.134)	-0.007 (0.018)	0.310 (0.220)	-0.023 (0.253)	0.164 (0.130)	-0.012 (0.007)	0.000 (0.002)	-0.030 (0.059)	-0.012 (0.025)	-0.011 (0.009)	-0.001 (0.003)	-0.004 (0.006)
5	0.142 (0.135)	0.022 (0.019)	0.081 (0.221)	-0.087 (0.255)	0.160 (0.131)	0.008 (0.007)	-0.002 (0.002)	0.024 (0.059)	0.011 (0.025)	-0.004 (0.010)	-0.002 (0.003)	0.003 (0.006)
6	-0.064 (0.134)	0.004 (0.018)	0.174 (0.220)	0.214 (0.254)	-0.042 (0.131)	-0.010 (0.007)	0.000 (0.002)	0.011 (0.059)	-0.013 (0.025)	-0.013 (0.009)	0.001 (0.003)	0.008 (0.006)
7	0.030 (0.132)	-0.004 (0.018)	0.158 (0.217)	-0.139 (0.251)	-0.028 (0.129)	0.000 (0.007)	0.000 (0.001)	-0.079 (0.058)	-0.027 (0.025)	-0.012 (0.009)	-0.001 (0.003)	-0.004 (0.006)
8	-0.087 (0.127)	-0.012 (0.017)	0.260 (0.208)	0.048 (0.241)	0.009 (0.124)	-0.010 (0.007)	-0.001 (0.001)	0.060 (0.056)	-0.007 (0.024)	-0.007 (0.009)	-0.001 (0.003)	0.005 (0.006)
9	<i>0.039</i> (0.134)	<i>-0.019</i> (0.018)	<i>-0.012</i> (0.219)	<i>-0.345</i> (0.254)	<i>0.155</i> (0.130)	<i>0.000</i> (0.007)	<i>-0.003**</i> (0.002)	<i>-0.042</i> (0.059)	<i>-0.020</i> (0.025)	<i>-0.005</i> (0.009)	<i>0.000</i> (0.003)	<i>0.006</i> (0.006)

joint test, coefficients on months 1 to 9 equal to 0

*p*-value 0.48 0.72 0.81 0.88 0.56 0.52 0.32 0.84 0.56 0.21 0.79 0.24

joint test, coefficients on months 1 to 9 are equal

*p*-value 0.39 0.64 0.9 0.83 0.49 0.47 0.26 0.78 0.68 0.2 0.8 0.18

*N* 42441 43116 43719 42489 40951 43000 42971 43427 43466 43728 42971 42971

*Notes:* All entries are coefficients on Ramadan exposure to daylight hours over subsequent 30 days as fraction of peak daylight hours during sample period. Gestation is either self reported or physician estimated. All samples use births with gestation between 37 and 42 weeks. Regressions include dummies for conception month, county and birth year. Italicized estimates may represent post natal exposure for some sample members. Standard errors in parentheses. \*significant at 10%; \*\* significant at 5%; \*\*\* significant at 1%

**Table 5: Effects of Ramadan Exposure in Months Prior to Birth on Disability Outcomes in Uganda**

Months Prior to Birth	<i>Muslims</i>					<i>Non-Muslims</i>				
	(1) Disability	(2) Sight/ Blind	(3) Hearing/ Deaf	(4) Mental/ Learning	(5) Psych.	(6) Disability	(7) Sight/ Blind	(8) Hearing/ Deaf	(9) Mental/ Learning	(10) Psych.
9	0.819** (0.359)	0.349* (0.193)	0.243** (0.117)	0.250*** (0.071)	-0.098 (0.072)	-0.023 (0.146)	-0.052 (0.080)	0.028 (0.052)	-0.037 (0.028)	0.045 (0.030)
8	0.087 (0.337)	-0.078 (0.180)	0.162 (0.110)	0.103 (0.066)	-0.068 (0.067)	-0.015 (0.137)	-0.043 (0.075)	0.043 (0.049)	-0.005 (0.026)	-0.028 (0.028)
7	-0.132 (0.349)	-0.022 (0.187)	0.13 (0.114)	0.028 (0.069)	0.058 (0.069)	-0.074 (0.142)	-0.142* (0.078)	-0.006 (0.051)	-0.006 (0.027)	0.010 (0.029)
6	0.197 (0.353)	0.074 (0.189)	0.161 (0.115)	0.100 (0.070)	-0.098 (0.070)	-0.091 (0.144)	0.082 (0.079)	-0.007 (0.051)	-0.017 (0.027)	0.017 (0.029)
5	0.085 (0.348)	-0.004 (0.187)	0.197* (0.114)	0.129* (0.069)	-0.058 (0.069)	0.209 (0.143)	-0.111 (0.079)	0.051 (0.051)	0.034 (0.027)	0.006 (0.029)
4	0.273 (0.352)	0.039 (0.189)	0.072 (0.115)	0.117* (0.070)	-0.049 (0.070)	-0.090 (0.144)	-0.030 (0.079)	0.048 (0.051)	-0.004 (0.027)	-0.017 (0.029)
3	0.104 (0.364)	0.124 (0.195)	0.099 (0.119)	0.039 (0.072)	-0.009 (0.073)	0.003 (0.147)	0.115 (0.081)	-0.018 (0.053)	-0.004 (0.028)	0.010 (0.030)
2	-0.266 (0.350)	-0.272 (0.187)	0.026 (0.114)	0.144** (0.069)	-0.019 (0.070)	0.039 (0.142)	-0.015 (0.078)	0.065 (0.051)	-0.043 (0.027)	0.036 (0.029)
1	-0.103 (0.366)	0.018 (0.196)	0.086 (0.120)	0.089 (0.072)	-0.034 (0.073)	0.208 (0.148)	-0.061 (0.082)	0.035 (0.053)	0.010 (0.028)	0.023 (0.030)
joint test, coefficients on months 1 to 9 equal to 0										
<i>p</i> -value	0.390	0.560	0.480	0.040	0.740	0.670	0.290	0.890	0.560	0.650
joint test, coefficients on months 1 to 9 are equal										
<i>p</i> -value	0.310	0.460	0.830	0.290	0.750	0.570	0.240	0.910	0.490	0.580
<i>Mean</i>	3.80%	1.06%	0.38%	0.14%	0.14%	5.21%	1.49%	0.61%	0.17%	0.20%
<i>N</i>	80924	80922	80923	80921	80921	640825	640789	640781	640777	640776

*Notes:* All entries are coefficients on Ramadan exposure measured as the percent of days overlapping with Ramadan during the nine months preceding birth (rampct). Each outcome is multiplied by 100, so that coefficients are in units of percentage points. All regressions include an indicator for female, birth month dummies, district of birth dummies and birth year dummies. Standard errors in parentheses, \*significant at 10%; \*\* significant at 5%; \*\*\* significant at 1%

**Table 6: Effects of Ramadan Exposure in Months Prior to Birth on Other Outcomes, Ugandan Muslims**

Months Prior to Birth	Sex Composition of Adult Population				Socioeconomic Outcomes				
	(1) Male	(2) Log Population	(3) Log Males	(4) Log Females	(5) Home Ownership	(6) Illiterate	(7) Years of Schooling	(8) No Schooling	(9) Employed
9	-0.020** (0.009)	0.001 (0.047)	-0.030 (0.059)	0.053 (0.065)	-0.026** (0.012)	0.008 (0.008)	-0.088 (0.068)	-0.004 (0.007)	0.000 (0.009)
8	-0.015* (0.009)	0.015 (0.044)	-0.034 (0.056)	0.081 (0.062)	-0.021* (0.011)	-0.015** (0.007)	0.119* (0.064)	-0.007 (0.007)	-0.001 (0.008)
7	-0.003 (0.009)	0.007 (0.045)	-0.055 (0.057)	0.083 (0.063)	-0.017 (0.011)	0.007 (0.008)	-0.009 (0.066)	0.001 (0.007)	-0.009 (0.009)
6	-0.021** (0.009)	-0.047 (0.045)	-0.081 (0.057)	0.01 (0.063)	0.008 (0.011)	-0.014* (0.008)	0.01 (0.067)	-0.013* (0.007)	0.013 (0.009)
5	-0.015 (0.009)	0.069 (0.045)	0.014 (0.057)	0.150** (0.064)	-0.018 (0.011)	0.012 (0.008)	-0.015 (0.067)	0.005 (0.007)	-0.019** (0.009)
4	-0.016* (0.009)	0.002 (0.045)	-0.03 (0.057)	0.036 (0.064)	-0.010 (0.011)	0.008 (0.008)	-0.045 (0.067)	0.006 (0.007)	-0.001 (0.009)
3	-0.026*** (0.010)	-0.085* (0.046)	-0.148** (0.057)	0.008 (0.064)	0.008 (0.012)	0.002 (0.008)	0.061 (0.069)	-0.002 (0.008)	0.005 (0.009)
2	-0.009 (0.009)	0.025 (0.045)	0.001 (0.056)	0.066 (0.063)	-0.005 (0.011)	0.009 (0.008)	0.069 (0.067)	0.009 (0.007)	-0.002 (0.009)
1	-0.009 (0.010)	-0.025 (0.047)	-0.031 (0.059)	0.012 (0.065)	0.000 (0.012)	0.005 (0.008)	-0.011 (0.069)	-0.005 (0.008)	0.001 (0.009)
joint test, coefficients on months 1 to 9 equal to 0									
<i>p</i> -value	0.100	0.460	0.420	0.520	0.050	0.100	0.440	0.390	0.460
joint test, coefficients on months 1 to 9 are equal									
<i>p</i> -value	0.640	0.360	0.570	0.770	0.070	0.070	0.380	0.300	0.380
<i>Mean</i>	0.506	4.205	3.554	3.399	0.734	0.30	6.94	0.25	0.66
<i>N</i>	81197	648	653	649	40463	78990	60117	80142	74348

*Notes:* All entries are coefficients on Ramadan exposure measured as the percent of days overlapping with Ramadan during the nine months preceding birth (rampct). All regressions include birth month and birth year dummies. Columns 2-4 use data on population counts aggregated to the level of birth year and birth month. Column 1 and columns 5-9 also include district of birth dummies. Column 5 is restricted to men. Columns 6-9 include a dummy for females.

Standard errors in parentheses, \*significant at 10%; \*\* significant at 5%; \*\*\* significant at 1%



**Table 7: Effects of Ramadan Exposure in Months Prior to Birth on Various Outcomes, Iraq**

Months Prior to Birth	Disability Outcomes				Socioeconomic Outcomes				
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
	Disability	Blind/ Vision	Deaf/ Hearing	Psych.	Multiple Wives	Home Ownership	Male	Primary School	Employed
9	0.299** (0.134)	0.030 (0.040)	-0.005 (0.016)	0.195*** (0.064)	-0.521* (0.289)	-1.577** (0.673)	0.433 (0.544)	-0.048 (0.619)	1.082*** (0.414)
8	-0.161 (0.121)	-0.018 (0.036)	-0.016 (0.015)	0.038 (0.058)	-0.148 (0.263)	-0.765 (0.607)	0.440 (0.490)	0.968* (0.561)	0.131 (0.374)
7	-0.088 (0.121)	0.026 (0.036)	-0.002 (0.015)	-0.101* (0.058)	-0.111 (0.270)	-2.060*** (0.611)	0.248 (0.492)	0.715 (0.565)	0.844** (0.375)
6	0.012 (0.118)	-0.011 (0.036)	-0.016 (0.014)	0.074 (0.057)	-0.428 (0.267)	-1.145* (0.597)	0.224 (0.482)	-0.139 (0.555)	1.078*** (0.367)
5	0.065 (0.117)	0.051 (0.035)	0.005 (0.014)	0.073 (0.056)	-0.494* (0.263)	-1.436** (0.590)	-0.018 (0.476)	0.820 (0.548)	0.827** (0.362)
4	0.083 (0.121)	0.029 (0.036)	-0.007 (0.015)	0.032 (0.058)	-0.585** (0.250)	-0.530 (0.609)	0.135 (0.491)	0.701 (0.569)	0.559 (0.374)
3	0.104 (0.124)	0.051 (0.037)	-0.012 (0.015)	0.029 (0.060)	-0.317 (0.262)	-0.882 (0.628)	-0.293 (0.506)	1.098* (0.584)	-0.629 (0.385)
2	0.007 (0.121)	0.045 (0.036)	-0.018 (0.015)	0.024 (0.058)	-0.169 (0.253)	-1.416** (0.613)	0.585 (0.494)	0.827 (0.568)	0.073 (0.377)
1	0.090 (0.128)	-0.032 (0.038)	0.008 (0.016)	0.021 (0.062)	0.193 (0.274)	-0.185 (0.650)	-0.535 (0.521)	1.103* (0.597)	-0.572 (0.397)
joint test, coefficients on months 1 to 9 equal to 0									
<i>p</i> -value	0.240	0.480	0.860	0.020	0.080	0.020	0.820	0.230	0.000
joint test, coefficients on months 1 to 9 are equal									
<i>p</i> -value	0.180	0.470	0.880	0.020	0.150	0.380	0.780	0.550	0.000
<i>Mean</i>	1.54%	0.14%	0.02%	0.35%	1.96%	73.46%	49.06%	57.50%	43.46%
<i>N</i>	288388	208711	287155	288388	75502	139516	288388	288388	287155

*Notes:* All entries are coefficients on Ramadan exposure measured as the percent of days overlapping with Ramadan during the nine months preceding birth (rampct). All regressions include birth month and birth year dummies. Each outcome is multiplied by 100, so that coefficients are in units of percentage points. Columns 5 and 6 are restricted to men. All regressions on pooled samples of men and women include a dummy for females.

Standard errors in parentheses, \*significant at 10%; \*\* significant at 5%; \*\*\* significant at 1%

**Table 8: Effects of Ramadan Exposure in Quarters Prior to Birth on Various Outcomes, US**

<i>Panel A: Effects relative to exposure 3 quarters prior to birth quarter</i>												
Quarters Prior to Birth	Pooled				1980 Census				ACS, 2005-2007			
	(1) Disability	(2) Less HS	(3) Male	(4) Log Earn.	(5) Disability	(6) Less HS	(7) Male	(8) Log Earn.	(9) Disability	(10) Less HS	(11) Male	(12) Log Earn.
2	1.382 1.089	2.457* 1.468	-1.371 2.268	-0.067 0.076	2.351 1.551	2.918 2.699	-2.193 3.695	-0.068 0.128	0.885 1.445	2.533 1.708	-3.426 2.948	-0.135 0.085
1	3.680*** (1.046)	-0.539 (1.409)	-2.844 (2.246)	-0.149* (0.079)	3.500** (1.524)	-1.393 (2.753)	-1.084 (3.625)	-0.022 (0.125)	3.603*** (1.386)	-2.330 (1.734)	-1.769 (2.826)	-0.002 (0.082)
0	1.437 (1.077)	-0.367 (1.453)	0.449 (2.177)	-0.115 (0.078)	2.710* (1.562)	2.004 (2.775)	-2.865 (3.727)	-0.2 (0.129)	1.108 (1.423)	-0.855 (1.660)	1.22 (2.905)	-0.078 (0.084)
joint test, coefficients on quarters 1 to 3 equal to 0												
p-value	0.000	0.110	0.400	0.280	0.130	0.390	0.880	0.390	0.050	0.020	0.340	0.320
Mean	6.07%	13.01%	57.25%	9.922	4.05%	15.95%	62.59%	8.996	7.09%	11.46%	54.45%	10.293
N	44121	45739	45474	24837	14313	15291	15174	7072	26707	30448	27169	16076
<i>Panel B: Effects relative to exposure in birth quarter and 3 quarters prior to birth quarter</i>												
Quarters Prior to Birth	Pooled				1980 Census				ACS, 2005-2007			
	(1) Disability	(2) Less HS	(3) Male	(4) Log Earn.	(5) Disability	(6) Less HS	(7) Male	(8) Log Earn.	(9) Disability	(10) Less HS	(11) Male	(12) Log Earn.
2	0.556 (0.896)	2.668** (1.209)	-1.596 (1.867)	-0.010 (0.065)	0.867 (1.294)	-2.850 (2.294)	-0.624 (3.080)	0.042 (0.106)	0.243 (1.186)	0.332 (1.414)	-4.133* (2.420)	-0.090 (0.070)
1	2.956*** (0.894)	-0.354 (1.205)	-3.102* (1.862)	-0.083 (0.065)	2.136 (1.305)	0.405 (2.316)	0.349 (3.110)	0.080 (0.106)	3.036** (1.179)	3.881*** (1.425)	-2.392 (2.405)	0.038 (0.070)
joint test, coefficients on quarters 1 to 3 equal to 0												
p-value	0.000	0.050	0.240	0.430	0.260	0.390	0.960	0.750	0.030	0.020	0.210	0.270
Mean	6.07%	13.01%	57.25%	9.922	4.05%	15.95%	62.59%	8.996	7.09%	11.46%	54.45%	10.293
N	44121	45739	45474	24837	14313	15291	15174	7072	26707	30448	27169	16076

*Notes:* All entries are coefficients on Ramadan exposure measured as the percent of days overlapping with Ramadan during the quarters preceding (or including) birth (rampct). All regressions include birth quarter, birth year and country of birth dummies. Dichotomous outcomes (disability, less HS and male) are multiplied by 100, so that coefficients are in units of percentage points. All regressions are on pooled samples of men and women include a dummy for females. Standard errors in parentheses, \*significant at 10%; \*\* significant at 5%; \*\*\* significant at 1%

# Contents

<b>A Biomedical Studies of Fasting</b>	<b>ii</b>
A.1 First Stage Effects of Ramadan . . . . .	ii
A.1.1 Caloric Intake and Weight Among Fasting Adults . . . . .	ii
A.1.2 Is Ramadan Observed by Pregnant Muslims? . . . . .	iii
A.2 Ramadan and Fetal Health . . . . .	iv
A.2.1 Pathways from Maternal to Fetal Health . . . . .	iv
A.2.2 Empirical Studies of Fetal Health . . . . .	v
A.3 Mechanisms of Fetal Programming . . . . .	v
A.4 Ramadan and Perinatal Health . . . . .	vii
A.4.1 Birth Outcomes . . . . .	vii
A.4.2 Longer-term Effects . . . . .	viii
A.5 Nutrition and the Sex Ratio at Birth . . . . .	viii
A.6 Hypotheses: Outcomes and Timing . . . . .	x
<b>B Data</b>	<b>x</b>
B.1 Michigan Natality Microdata . . . . .	x
B.2 Uganda Census 2002 . . . . .	xii
B.3 Iraq Census 1997 . . . . .	xiii
B.4 US Census 1980, ACS, 2005-2007 . . . . .	xiii
B.5 Other Suitable Datasets? . . . . .	xiv
<b>C Birth Outcomes by Gestation Length</b>	<b>xiv</b>
<b>D Distributional Effects on Birth Weight</b>	<b>xvi</b>

# E Appendix Tables & Figures

*Tables A1-A7, Figures A1-A4*

## A Biomedical Studies of Fasting

We begin by summarizing evidence on the “first stage” effect of fasting during Ramadan. That is, what is the existing evidence that Ramadan fasting can have a detectable effect on health? In Section A.1, we consider studies of caloric intake and weight change during intermittent fasting and summarize survey data on the prevalence of Ramadan fasting among pregnant women. Second, we discuss the potential impacts of maternal biochemical changes caused by fasting (accelerated starvation) on the fetus in Section A.2. Third, we examine potential pathways by which intermittent fasting could have lasting effects through “fetal programming” in Section A.3. Fourth, we review the empirical studies that have explicitly examined the effects of Ramadan on birth and early childhood outcomes in Section A.4. Fifth, we briefly summarize a separate literature on nutrition and the sex ratio at birth – which to date has not used Ramadan fasting for identification – in Section A.5. Finally, we distill the preceding into research hypotheses which we will apply to our data in Section A.6.

### A.1 First Stage Effects of Ramadan

#### A.1.1 Caloric Intake and Weight Among Fasting Adults

Ramadan fasting in the adult population (i.e. not conditioning on pregnancy) has been associated with modest but statistically significant declines in the weight of fasters of around 1 to 3 kg (Husain et al. [1987]; Ramadan et al. [1999]; Adlouni et al. [1998]; Mansi [2007]; Takruri [1989]) Reductions in weight are sometimes (but not always) accompanied by declines in caloric intake and likely depend on dietary customs in specific countries.<sup>1</sup>

Two studies are of particular relevance. First, in a study of 185 pregnant women, Arab [2003] found that over a 24 hour period encompassing the Ramadan fast, over 90 percent of the women had a deficiency of over 500 calories relative to the required energy intake and 68 percent had a deficiency of over 1000 calories. Second, in the only large scale population-based study we are aware of, Cole [1993] found striking evidence of sharp weight changes during Ramadan for women in Gambia. The study was notable because it used fixed effects with 11 years of panel data and controlled for calendar month, calendar year, and stage of pregnancy (or lactation). Appendix Figure A1, taken from the study, shows that relative to the rest of the year, there is an increase in weight during the four

---

<sup>1</sup>For example, Husain et al. [1987] found reductions in caloric intake of between 6 percent and 25 percent relative to nonfasting conditions among Malaysians. In contrast, Adlouni et al. [1998] found a 20 percent increase in calories per day among Moroccans.

weeks prior to Ramadan and a sharp increase in weight at the very beginning of Ramadan. This is followed by an abrupt fall in weight of over 1kg (2.2 pounds) during the subsequent 3 weeks of fasting. The figure provides striking visual evidence that daytime fasting during Ramadan is affecting weight gain.

In any case, as we discuss in section 2.1 of the paper, fasting may induce maternal biochemical changes and reprogramming of the neuro-endocrine system due to alterations in the the *timing* of nutritional intake even if overall caloric intake or weight change is unaffected.

### A.1.2 Is Ramadan Observed by Pregnant Muslims?

Although pregnant women may request an exemption from fasting, they are expected to “make up” the fasting days missed during pregnancy after delivery and this requirement may discourage pregnant women from seeking the exemption since they may be the only member of the household fasting [Hoskins, 1992, Mirghani et al., 2004].<sup>2</sup> Mirghani et al. [2004] noted: “Most opt to fast with their families rather than doing this later”:636. In addition, some Muslims interpret Islamic Law as requiring pregnant women to fast. For example, the religious leader of Singapore’s Muslims held that: “a pregnant woman who is in good health, capable of fasting and does not feel any worry about herself or to her foetus, is required and expected to fast like any ordinary woman” [Joosoph and Yu, 2004].<sup>3</sup> Furthermore, since fasting during Ramadan is one of the five pillars of Islam and is a central part of the culture of the Muslim community, many women fear a loss of connection with the community or would feel guilty about not observing Ramadan [Robinson and Raisler, 2005].

As far as we are aware, comprehensive data on Ramadan fasting during pregnancy do not exist. Various surveys of Muslim women suggest that fasting is the norm. For example, of the 4,343 women delivering in hospitals in Hamadan, Iran in 1999, 71% reported fasting at least 1 day, “highlighting the great desire of Muslim women to keep fasting in Ramadan, the holy month” [Arab and Nasrollahi, 2001]. In a study in Singapore, 87% of the 181 muslim women surveyed fasted at least 1 day during pregnancy, and 74% reported completing at least 20 days of fasting [Joosoph and Yu, 2004]. In a study

---

<sup>2</sup>There are some differences in interpretation of the Koran among Imams regarding whether pregnant women must make up the fasting days later or simply pay alms for the poor, or both. See, for example, [http://islam1.org/iar/imam/archives/2006/09/09/fasting\\_the\\_month\\_of\\_ramadaan.php](http://islam1.org/iar/imam/archives/2006/09/09/fasting_the_month_of_ramadaan.php)

<sup>3</sup>Similarly, Arab and Nasrollahi [2001] noted that “According the Islamic teaching pregnant women are allowed to fast if it is not harmful to them”; faculty at the Kurdistan Medical Science University in Iran noted that pregnant and breastfeeding women “who fear for the their well being or that of the foetus/child” may be exempted from fasting [Shahgheibi et al., 2005].

conducted in Sana'a City, Yemen, more than 90 percent fasted over 20 days [Makki, 2002]. At the Sorrento Maternity Hospital in Birmingham, England, three quarters of mothers fasted during Ramadan [Eaton and Wharton, 1982]. In a study conducted in Gambia, 90 percent of pregnant women fasted throughout Ramadan [Prentice et al., 1983]. In the US, a study of 32 Muslim women in Michigan found that 28 had fasted in at least one pregnancy and reported that 60-90 percent of women from their communities fast during pregnancy [Robinson and Raisler, 2005].

In summary, survey data indicate that most but not all women observe the Ramadan fast during pregnancy. To the extent that pregnant Muslim women do not fast, ITT estimates are conservative estimates of fasting's effect. As discussed in Section 6 of the main paper, fasting observance is likely highest in early pregnancy.

## A.2 Ramadan and Fetal Health

### A.2.1 Pathways from Maternal to Fetal Health

Does exposure to ketones during “accelerated starvation” (Section 2.1 of the main text) impair the neural development of the fetus? Controlled studies of mice and rats have shown that prenatal exposure to ketones result in impaired neurological development. [Hunter and Sadler, 1987, Moore et al., 1989, Sheehan et al., 1985]. Hunter and Sadler [1987] reference studies showing ketones “rapidly diffuse from the maternal circulation across extraembryonic membranes”:263. They also point out that in addition to the period of neurulation (3rd to 4th week of gestation in humans), the earliest stages of embryogenesis when the “primitive streak” is observed (the 13th day post-conception), may be especially susceptible to ketones. Moore et al. [1989] noted that “even a relatively brief episode of ketosis might perturb the development of the early embryo”:248. They also emphasize that the effects of ketones were to slow neurological development rather than to produce a malformation. This may explain why similar studies in human populations have not (for the most part) found evidence of congenital malformations [ter Braak et al., 2002]

A related literature has examined the effects of poor metabolic regulation during pregnancy in mothers with Type 1 diabetes. In this case although the primary concern is avoiding hyperglycemia (abnormally high blood glucose), this sometimes results in severe cases of hypoglycemia (abnormally low blood glucose). The latter case may be instructive for understanding the potential effects of accelerated starvation since blood glucose drops after a prolonged fast. Some studies of *in utero* exposure to hypoglycemia

among diabetic mothers have shown that fetal growth is reduced and that the key period is between the fourth to sixth weeks of gestation [ter Braak et al., 2002]). It has also been shown that hypoglycemia among *non-diabetic* mothers is also associated with lower birth weight [Scholl et al., 2001]. Studies of diabetic mothers have shown long-term effects of *in-utero* exposure to hypoglycemia on cognitive functioning during childhood (Rizzo et al. [1991], Langan et al. [1991]). Interestingly, Rizzo et al. [1991] only found effects on intelligence scores for those exposed during the first trimester.

### A.2.2 Empirical Studies of Fetal Health

Fetal health measures have the advantage of permitting panel data techniques to address selection in to maternal fasting but the disadvantage of not being standardized health metrics. Several studies of maternal fasting during Ramadan have found adverse effects on at least two of these fetal health indicators. Mirghani et al. [2004] found evidence of reduced fetal breathing movements where measures of fetal breathing were taken both before and after fasting on the same day. The same study, however, found no change in overall body movements, fetal tone or maternal appreciation.<sup>4</sup> Mirghani et al. [2005] found a significantly fewer heart rate accelerations among pregnant women who were fasting during Ramadan late in pregnancy compared to controls. This was observed despite relatively short diurnal fasts (less than 10 hours duration) and the absence of significant changes in glucose levels. DiPietro et al. [2007] found a strong association between variation in fetal heart rate *in utero* and mental and psychomotor development and language ability during early childhood. Finally, Mirghani et al. [2007] found no effect of Ramadan fasting on uterine arterial blood flow.

In contrast, studies of hypoglycemia in animals and humans have examined the fetal heart rate, fetal breathing movements, and limb and body movements in order to identify impairments to fetal development. A review of these studies in ter Braak et al. [2002] do not show much affect of moderate hypoglycemia on fetal conditions.

### A.3 Mechanisms of Fetal Programming

We now discuss how disruptions to fetal health can have permanent effects. In a review of epidemiological studies on the fetal origins of adult diseases, Jaddoe and Witteman [2006] describe two hypotheses related to our study. The first is described as “fetal under-nutrition.” According to this view, inadequate prenatal nutrition leads to developmental

---

<sup>4</sup>A significant reduction in upper limb movements was noted but there was a concern that this might be due to observer bias.

adaptations that are beneficial for short-term survival but lead to lower birth weight. However, by permanently reprogramming the physiology and metabolism of the fetus, this ultimately makes the body susceptible to heart disease and diabetes during adulthood.<sup>5</sup> Although most studies of fetal origins have relied on blunt measures such as birth weight to proxy for nutritional restriction during pregnancy, a recurring theme in many studies is that fetal programming may occur even in the absence of birth weight effects. For example, studies of the Dutch famine have showed that those exposed to the famine early in gestation had dramatically higher rates of heart disease but did not have lower birth weight [Painter et al., 2005]. Similarly animal studies have often found evidence of fetal programming without detecting significant changes in fetal weight. e.g. Nishina et al. [2004]

A second prominent hypothesis is that nutritional restrictions inhibit the development of a placental enzyme that is required to convert cortisol into inactive cortisone, thereby exposing the fetus to excessive amounts of cortisol. It is suggested that exposure to glucocorticoids such as cortisol *in utero* leads to a reprogramming of the hypothalamic–pituitary adrenal axis (HPA) which in turn, could lead to impaired fetal development and worse health during adulthood.

In controlled animal studies, researchers have linked nutritional restrictions very early in gestation to an altered neuro-endocrine system, e.g., Nishina et al. [2004]. With respect to humans, Herrmann et al. [2001] have shown an association between fasts of 13 hours or longer and higher levels of plasma corticotrophin-releasing hormone (CRH) which could reflect a reprogramming of the HPA axis. As noted in the main text, Dikensoy et al. [2009] show that Ramadan fasting is associated with elevated cortisol levels during pregnancy (relative to pre-pregnancy levels), but not for non-fasting mothers. Kapoor et al. [2006] describe how the effects of fetal programming of HPA in humans may result in cognitive impairment; due to the complex feedback mechanisms involved, these effects may not be evident “until adulthood or early old age”. The authors also emphasize that many of the long-term effects may be sex-specific.

The existing literature on fetal origins however, has made little use of quasi-experimental research designs to address potential confounding factors or to identify the underlying mechanisms. Jaddoe and Witteman [2006] recently concluded: “Thus far, it is still not known which mechanisms underlie the associations between low birth weight and diseases in adult life. The causal pathways linking low birth weight to diseases in later life seem

---

<sup>5</sup>Jaddoe and Witteman [2006] note that this view has evolved into a more “general developmental plasticity model in which various fetal and post-natal environmental factors lead to programming responses”:93.



to be complex and may include combined environmental and genetic mechanisms in various periods of life. Well-designed epidemiological studies are necessary to estimate the population effect size and to identify the underlying mechanisms” Jaddoe and Witteman [2006, 91].

## A.4 Ramadan and Perinatal Health

### A.4.1 Birth Outcomes

Existing studies of birth outcomes have relied on comparisons between mothers who reported fasting to those who did not. Kavehmanesh and Abolghasemi [2004] compared 284 births to mothers in Tehran with a “history of fasting during pregnancy” to 255 mothers who did not fast. Although there were no statistically significant differences with respect to maternal education or height, pre-pregnancy BMI’s were substantially higher in the fasting group. For such comparisons, the conditional independence assumption required for causal inference [Angrist and Pischke, 2009] is tenuous. Shahgheibi et al. [2005] studied 179 newborns for whom Ramadan fell in the third trimester of pregnancy. Among fasters, birth weight was lower by 33 grams, birth length was lower by about 0.2 centimeters while head circumference was larger by 0.08 centimeters. Since these differences were not statistically significant with the small sample used, the authors concluded that fasting during the third trimester had “no effect” on growth indices. Arab and Nasrollahi [2001] studied 4,343 pregnancies in the Hamdan province of Iran and concluded that fasting did not impact birth weight. They did note however, that the incidence of low birth weight ( $< 2500$  grams) was higher among fasters in the second trimester but that this was significant only at the 9 percent level.

The largest and perhaps most commonly cited study on the effects of Ramadan on birth weight conducted a retrospective analysis of 13,351 babies born at *full term* from 1964-84 in Birmingham, England Cross et al. [1990]. Babies were categorized as Muslim on the basis of the first three letters of the mother’s surname and were matched to control groups by age. However, this study did not compare the birthweights of Muslims *in utero* during Ramadan to Muslims who were not *in utero* during Ramadan but instead compared across groups of Muslims and Non-Muslims. In addition, by design the study did not look at the potential effects of Ramadan on gestation length. Although Cross et al. [1990] found no significant effects on mean birth weight, like Arab and Nasrollahi [2001], they also found a higher incidence of low birth weight among fasters during the second trimester. Opaneye et al. [1990] found that in Al-Kharj, Saudi Arabia, the incidence of

low birth weight increased during Islamic festivals, Ramadan in particular. 9.9% of the 415 births were below 2,500 grams during Ramadan, versus 6.3% for the 4,865 births in non-Ramadan months. Finally, Malhotra et al. [1989] and Mirghani and Hamud [2006] found no effects on birthweight and APGAR scores, even though they detected substantial biochemical changes.

A separate literature has found that skipping meals (not associated with Ramadan) has been associated with preterm delivery. Siega-Riz et al. [2001] studied diets during the second trimester of pregnancy for over 2000 women in North Carolina and found that women who did not follow the optimal guidelines of three meals and two snacks a day were 30 percent more likely to deliver preterm. They suggest that this is consistent with experimental evidence from animal studies. Herrmann et al. [2001] also reported that women who fasted for 13 hours or more were three times more likely to deliver preterm.

While most studies have focussed on birth weight, Mirghani and Hamud [2006] considered a broader range of birth outcomes. Specifically, they compared 168 pregnant fasters to a control group of 156 non-fasting mothers and found significantly higher rates of gestational diabetes, induced labor, cesarian sections, and admission to the special baby care unit.

#### **A.4.2 Longer-term Effects**

We are aware of just one previous study of on long-term effects of Ramadan. Azizi et al. [2004] surveyed outcomes among 191 children enrolled in 15 Islamic primary schools in Iran and their mothers about Ramadan fasting during pregnancy. Approximately half of the mothers selected for the analysis sample reported fasting. More than 1,600 mothers returned questionnaires regarding their fasting behaviour during pregnancy. However, the fraction of this initial sample who fasted during pregnancy is not reported by Azizi et al. [2004]. Among fasting mothers, those fasting during the third trimester were over-sampled. No significant difference in the IQ's of the children were found by maternal fasting behaviour. As mentioned in the main text, Ewijk [2009] analyzes long-term Ramadan effects using the Indonesian Family Life Study data. This work was inspired by ours and generally finds corroborative results.

### **A.5 Nutrition and the Sex Ratio at Birth**

Widely studied in evolutionary biology, the Trivers-Willard hypothesis posits that the reproductive success of sons is more sensitive to maternal condition than that of daughters

[Trivers and Willard, 1973]. Therefore, parents experiencing better conditions may favor male offspring. More generally, the sex ratio at birth and early childhood may proxy for unobserved health conditions given disproportionate male susceptibility to fetal and infant mortality [Kraemer, 2000, Mathews and Hamilton, 2005]. One proposed mechanism by which adjustment to the sex ratio may take place is through the nutritional status of the mother while pregnant [Cameron, 2004]. Roseboom et al. [2001] found that prenatal exposure to the Dutch famine of 1944-45 reduced the sex ratio of live births. Similarly, Almond et al. [2007] found the sex ratio in China was skewed toward females for cohorts born during the Great Leap Forward Famine. Askling et al. [1999] showed that women who experience severe morning sickness were much more likely to have girls.

A widely-publicized study by Mathews et al. [2008] has for the first time drawn a link between maternal nutrition prior to conception and the sex ratio at birth. The authors collected detailed information on food intake prior to pregnancy, early in pregnancy (14 weeks gestation) and late in pregnancy (28 weeks gestation) in Britain. They found no differences in the rates of male births arising from differences in nutritional intake either early or late in pregnancy but found a highly statistically significant positive relationship between high nutritional scores prior to conception and the birth of male offspring. They further examined the detailed data on sources of nutrition and found that among 133 food items consumed prior to pregnancy, only breakfast cereals was strongly associated with infant sex. The authors speculated that the mechanism underlying this connection is that the skipping of breakfast

*“extends the normal period of nocturnal fasting, depresses circulating glucose levels and may be interpreted by the body as indicative of poor environmental conditions.”*

Mathews et al. [2008] also referenced work by Larson et al. [2001] on *in vitro* fertilization of bovine embryos showing that glucose “enhances the growth and development of male conceptuses while inhibiting that of females.”

The study by Mathews et al. [2008] was observational and did not explore the source of dietary differences across mothers, nor did it control for some other factors known to influence the sex ratio (e.g., partnership status at the time of conception [Norberg, 2004]). Short of a controlled experiment, the research design utilized here has the advantage of leveraging plausibly exogenous differences in maternal fasting.

## A.6 Hypotheses: Outcomes and Timing

In this section, we distill findings from the biomedical literature most relevant to our Ramadan analysis. Appendix Table A1 summarizes the set of health outcomes we might expect to be affected by fasting (column 1), notes the mechanism (column 2), and lists the months of prenatal exposure that have been found or suggested to be particularly important (column 3). These hypotheses are based on either a clearly defined pathway linking fasting to a particular outcome, or an empirical result that has been established, irrespective of whether there is an explicit mechanism described in the study. In many of the studies, the period of *in utero* exposure was selected by design and therefore does not exclude effects in other periods.

In the case of birthweight, we describe four mechanisms through which fasting might operate and one empirical finding based on the Dutch famine. Two of the birthweight mechanisms are tightly linked to exposures occurring in early pregnancy. For several outcomes there are no clear hypotheses concerning timing that we could discern; a reasonable hypothesis would be to jointly test the effects of Ramadan exposure during all gestation months.

With respect to longer-term effects, in virtually all cases exposure to fasting during early pregnancy is the predominant hypothesis. For cognitive function, there are several arguably distinct channels through which prenatal fasting might be detrimental.

## B Data

### B.1 Michigan Natality Microdata

Our ancestry-based proxy for Muslim status is coded as follows. For births from 1989 to 1992, we include mothers who report their ancestry as “Arab/Middle Eastern” in the ITT (whose pregnancies also overlap with a Ramadan). Starting in 1993, several specific country codes for ancestry are reported. From 1993 to 2006 our ITT group includes mothers who report ancestry as: Arab/Middle Eastern, Arab/North African, Iran, Afghanistan, Mauritania, Somalia, Turkey or Western Sahara. Overall, 96% of our treatment group report their ancestry as Arab/Middle Eastern, hence we refer to the group as Arabs.

We also implement several other sample selection rules to minimize measurement error and misclassification of Muslims into the control group. We dropped births with no reported ancestry or where the ancestry might possibly include parents who are practicing

Muslims (e.g. Southeastern Asians). We also dropped non-Arab Blacks to avoid the possibility that there might be “Black Muslims” in our sample. We also dropped twin births and restricted the sample to births among mothers between the ages of 14 and 45.

The summary statistics are shown in Appendix Table A2. Arab mothers reported a year less education than non-Arab mothers on average, and are substantially more likely to receive Medicaid (46% versus 27%). Arab families are also larger (average parity is 18% higher for Arabs). Despite these differences in socioeconomic measures, birth outcomes are more similar. Rates of low birth weight and prematurity are actually slightly lower for Arabs than for non-Arabs. The geographic distribution of Arabs by zipcode in Michigan is shown in Appendix Figure A2. Somewhat surprisingly perhaps, Arabs do not appear noticeably more concentrated around the Dearborn and Detroit area than non-Arabs (Panel A).

The key variables for assigning *in utero* Ramadan exposure are birth date and gestation length. Michigan natality data include exact date of birth and a self-reported date of last menstrual period (LMP) for about 70 percent of the sample. The problem of selective reporting of LMP based on socioeconomic status is well known [Hediger et al., 1999]. There is also a field containing the physician’s estimate of gestation length, but we do not know how it is calculated or when during gestation.<sup>6</sup> We follow related epidemiological studies that utilize a simple algorithm for coding gestation (e.g., Siega-Riz et al. [2001], Herrmann et al. [2001]): gestation based on LMP is used except if it is missing or if it differs with physician estimated gestation by more than 14 days, in which case the physician estimated measure is substituted.

Appendix Figure A3 provides a hypothetical example to illustrate how our daily measures of Ramadan exposure are calculated. In 1989, Ramadan began on April 7th and ended on May 6th. For someone who was conceived on April 6th, his or her entire first month of gestation would overlap with Ramadan, i.e. *exppct*=1. Since during this Ramadan, daylight hours averaged about 13.7 hours per day, compared to 15.2 during the summer solstice, the hours exposure measure (*exphrspct*) peaks at about 0.9.

---

<sup>6</sup>A key concern is that this could be endogenous to Ramadan exposure. For example, if Ramadan affects fetal size and if physician estimates of LMP are based on measures of fetal size, this could lead to mis-measurement of the timing of Ramadan exposure. In addition, this measure might not be calculated uniformly and may depend on the timing of the first doctor visit and could therefore, be correlated with mother’s socioeconomic status.

## B.2 Uganda Census 2002

The Uganda Census contains roughly 2.5 million records (10% sample). Our main analysis sample includes men and women ages 20 to 80. Individuals whose birth month or birth year were imputed are dropped.<sup>7</sup> For each outcome measure, we recoded those with imputed data to missing. The disability question in the Uganda survey instrument asks: “Does (name) have any difficulty in moving, seeing, hearing, speaking difficulty, mental or learning difficulty, which has lasted or is expected to last 6 months or more?” The following specific disabilities are recorded in the dataset: blind or vision impaired, deaf or hearing impaired, mute, disability affecting lower extremities, disability affecting upper extremities, mental/learning disabilities and psychological disabilities. The original unharmonized variables label the last two variables “mental retardation” and “mental illness” while IPUMS-I relabelled them as “mental” and “psychological”. Our own reading of the instructions to the Uganda Census enumerators suggests that this relabelling was indeed appropriate. The former measure appears to identify those with “mental or learning disabilities” while the latter identifies those exhibiting “strange behaviors”. A subsequent question asks about the origin of the reported disability. The responses are coded into the following categories: congenital, disease, accident, aging, war injury, other or multiple causes.

The summary statistics are reported in Appendix Table A3. In contrast to Michigan, Uganda Muslims tend to have higher average SES. Muslims are less likely to be illiterate than non-Muslims (30% versus 36%) and completed more schooling. Disability rates for Muslims are also lower – 3.8% versus 5.2% for non-Muslims. Both Muslims and non-Muslims share a strong seasonality in the frequency of births by month. For both groups, birth in June was more than 50% more likely than birth in December. The frequency distribution across Ramadan ITT gestation months is much more uniform, and similar between Muslims and non-Muslims.

ITT assignment is determined by the reported birth month. We found age heaping: spikes in the number of respondents reporting of ages ending in zeroes (e.g. 20, 30, 40), suggesting measurement error. We therefore excluded records reporting these round-number ages.

---

<sup>7</sup>The IPUMS-I “unharmonized” variables contain imputation flags. We allowed records with “logical imputations” but dropped records imputed by “hot-deck”.

### B.3 Iraq Census 1997

The Iraq Census is also a 10 percent sample. We dropped individuals who reported ages ending in seven and those aged 50, 60 and 70 due to heavy age heaping at those values. We also drop those reporting birth months of January and July because of heaping at those months. For example, there were nearly three times as many respondents reporting a July birth as a June birth. This leaves us with a sample of over 288,000 individuals between the ages of 20 and 80.

### B.4 US Census 1980, ACS, 2005-2007

We assigned our Muslim proxy to respondents born in a country which currently has at least 80 percent Muslims.<sup>8</sup> We used the following 38 countries: Saudi Arabia, Somalia, Afghanistan, Maldives, Western Sahara, Turkey, Iran, Algeria, Mauritania, Yemen, Tunisia, Oman, Comoros, Djibouti, Morocco, Pakistan, Libya, Iraq, Iraq/Saudi Arabia, Tajikistan, Jordan, Qatar, Senegal, Azerbaijan, Egypt, Mali, Niger, Gambia, Uzbekistan, Turkmenistan, Indonesia, Bangladesh, Syria, Guinea, Kuwait, Bahrain, West Bank and Gaza. We excluded individuals who reported languages that would suggest they were non-Muslims.<sup>9</sup> We also dropped Iranian immigrants living in California who are predominantly Jewish.

In order to roughly gauge the extent of classification error, we can consider data from the 2001 Canada Census, which includes measures both of religion and country of birth. 67% of Canadian immigrants born in our predominantly-Muslim countries reported Muslim as their religion.<sup>10</sup> If the immigration patterns were assumed to be identical, then this would imply that roughly one in three of our sample of Census sample will not observe the Ramadan fast because they are not Muslim. Our control sample was restricted to whites born in the United States who did not report ancestries outside of the US or Western Europe. We chose not to use reported ancestry for assigning the Muslim proxy because of high rates of underreporting and the fact that it would combine first and second generation immigrants. We might suspect that that Ramadan observance is higher for first generation immigrants.

For earnings we use the CPI to convert values into 1997 dollars and restrict the sample to those aged 20 to 60 years.

---

<sup>8</sup>We took estimates derived from US State Departments's International Religious Freedom Report, the CIA World Factbook and Adherents.com as reported on the following Wikipedia URL:

[http://en.wikipedia.org/wiki/Islam\\_by\\_country](http://en.wikipedia.org/wiki/Islam_by_country)

<sup>9</sup>For example, some individuals born in Indonesia may report "Dutch" as their language.

<sup>10</sup>We thank Kevin Milligan (UBC) for providing these estimates.

## B.5 Other Suitable Datasets?

The Uganda and Iraq Census microdata were obtained from the Integrated Public Use Microdata Series - International (IPUMS-I). Other potentially relevant IPUMS-I samples are those for Egypt, Jordan, and Malaysia. Each has a large population of Muslims with Census data that purportedly include birth month.<sup>11</sup> Religion is not reported for Egypt and Jordan, but like Iraq, are overwhelmingly Muslim. However, in Egypt 85% of the sample is missing birth month. 40% are missing birth month in Malaysia, and only .5% of adults report a work disability. In Jordan's data, birth year and place of birth are missing.

In the US, month of birth is not reported in the decennial Census. While the National Health Interview Survey (NHIS) reports birth month, it does not disclose religion, detailed ethnicity, or country of birth.

## C Birth Outcomes by Gestation Length

As noted in the main text, variation in gestation length can generate compositional effects for pregnancies exposed to Ramadan late in gestation. For example, pregnancies with exposure to Ramadan in month 9 of pregnancy are exclusively full-term, but ITT estimates of the month 9 effect include premature births where Ramadan fell after delivery. Even for earlier months of pregnancy, the relationship to the ITT may differ depending on gestation length. Finally, gestation itself may respond to the ITT.

In Appendix Table A4, we vary the samples used to estimate birthweight effects based on the length of gestation. In column (1) we start with a sample of all births with gestation length 25 weeks to 42 weeks allowing us to include preterm births. The coefficients for months 1 to 5 are interpretable for all sample members.<sup>12</sup> We then progressively tighten the sample restriction by increasing the lower bound on gestation age producing samples of length 31-42 weeks in column (2), 35-42 weeks in column (3) and 39-42 weeks in column (4). This gradually increases the number of exposure month coefficients that can be interpreted. Interestingly, this does not appear to have any pronounced effects on the results. With the larger samples that include some preterm births, exposure during the sixth months is significant at the 10 percent level in two of the three specifications. The gestation samples that include pre-term births now provide additional evidence (on top

---

<sup>11</sup>Birth month and religion are available in the census of South Africa (unharmonized variables in IPUMS-I), but South Africa's Muslim population is relatively small (roughly 1.5% of the population).

<sup>12</sup>For example, exposure in months 6 to 9 will include the effects of Ramadan exposure in the postnatal period for the small subsample of those with only 25 weeks gestation.



of the “no gestation” sample results from Table 1 of the main text), that the coefficients are jointly different from zero with significance levels ranging from 6.1 percent to 12.4 percent.

In column (5) we attempt to address the fact that our ITT estimates may be attenuated by the inclusion of Chaldeans, who are not Muslim but might self identify as Arab. Specifically we use the sample of full-term births and drop mothers whose zipcode of residence has a ratio of Chaldeans to Arabs of greater than 0.5 according to the 2000 Census. This drops our sample by more than 25 percent and raises the standard errors considerably. However, there is some suggestive evidence of a stronger ITT effect. The most dramatic change is for exposure in the 7th month of gestation where we now see an effect: nearly a 60 gram drop in birthweight that is significant at the 1 percent level. The  $p$ -value on the  $F$ -test of joint significance also falls to 0.071 from 0.277 in column (4). In other specifications we have not always found much of a difference between the overall sample and the sample that drops heavily Chaldean zipcodes. This may be due to the large geographic overlap in the concentrations of Arabs and Chaldeans zipcodes and the relatively coarse approach dictated by the available data. In this sub-sample, we are arguably, estimating the effects of Ramadan exposure among Arabs in less concentrated areas where observance rates may be lower.

In columns (6) through (9) we estimate the effects of *in utero* Ramadan exposure on gestation weeks using the same approach on the full sample of Arabs. Among those with 25-42 weeks gestation, we estimate that Ramadan exposure during the second and fifth months of pregnancy reduces gestational age by a little more than 0.1 weeks or by roughly a day ( $p < 0.10$ ). However, these are the only months where exposure is statistically significant and we cannot reject that the first five gestation months jointly, have no effect. The effect from month 5 exposure weakens a bit as the sample is gradually restricted. Interestingly, column (9) shows that when the sample is confined to only full term births, four gestation months (2, 3, 5 and 8) are significant at at least the 10 percent level. Finally, in column (10) with the more restricted sample of non-Chaldean zipcodes we find a highly significant effect from second month exposure. In order to assess what portion of the birthweight effect might be attributed to decreased gestation, we ran a separate regression of birthweight on gestation for the sample of non-Arab mothers and found that each additional week of gestation adds about 165 grams. This suggests that a reduction in gestation of 0.1 weeks is associated with a 16 gram reduction in birthweight. From this rough exercise, it appears that less than half of the overall reduction in birthweight of around 40 grams that we typically estimate in Tables 1 and Appendix Table A4 is due to

reduced gestation as opposed intrauterine growth restriction (IUGR). For those exposed in the first month of pregnancy, it appears that virtually all of the birthweight reduction is due to IUGR.

## D Distributional Effects on Birth Weight

In Figure A4 we show which parts of the birthweight distribution are affected by early exposure to Ramadan fasting. We plot kernel densities comparing the birthweight distribution of those Arabs with no *in utero* exposure to those who had a significant exposure to Ramadan in the first month of gestation ( $exphrspect1 > 0.5$ ).<sup>13</sup> Most of the effect is in the middle part of the distribution. Specifically, those with first month exposure to Ramadan are more likely to have birthweight between 2800 and 3200 grams and less likely to have birthweight between 3250 and 3900 grams.<sup>14</sup> This suggests that little of the effect is at the low end of the distribution among those classified as “low birthweight” (<2500 grams).

## References

- A Adlouni, N Ghalim, R Sale, N Had, and A Bensliman H Parra and. Beneficial effect on serum apo ai, apo b and lp ai levels of ramadan fasting. *Clinica Chimica Acta*, 271 (2):179–189, March 1998.
- Douglas Almond, Lena Edlund, Hongbin Li, and Junsen Zhang. Long-term effects of the 1959-61 china famine: Mainland china and hong kong. *NBER Working Paper No. 13384*, September 2007.
- Joshua D. Angrist and Jörn-Steffen Pischke. *Mostly Harmless Econometrics: An Empiricist’s Companion*. Princeton University Press, Princeton, New Jersey, 2009.
- M Arab. Ketonuria and serum glucose of fasting pregnant women at the end of a day in ramadan. *The Journal of Physiology*, 553(2):637–647, 2003.
- Malihe Arab and Shahla Nasrollahi. Interrelation of ramadan fasting and birth weight. *Medical Journal of the Islamic Academy of Sciences*, 14(3):91–95, 2001.

---

<sup>13</sup>We also ran linear probability regressions using indicator variables of being in specific intervals of the birthweight distribution (e.g. 300-600 grams, 600-900 grams etc.) as an outcome including our other controls and obtained very similar results.

<sup>14</sup>There are also small differences between the samples in the interval from 2100 grams to 2700 grams, and for birthweight greater than 4500 grams.

- J. Askling, G. Erlandsson, M. Kaijser M, O. Akre, and A. Ekblom. Sickness in pregnancy and the sex of the child. *Lancet*, 354(9195):2053, December 1999.
- Fereidoun Azizi, Hossein Sadeghipour, Behnam Siahkolah, and Nasrollah Rezaei-Ghaleh. Intellectual development of children born to mothers who fasted during pregnancy. *International Journal of Vitamin and Nutrition Research*, 74(5):374–380, September 2004.
- Elissa Z. Cameron. Facultative adjustment of mammalian sex ratios in support fo the trivers-willard hypothesis: evidence for a mechanism. *Proceedings of the Royal Society B: Biological Sciences*, 271(1549):1723–1728, 2004.
- TJ Cole. *Seasonality and Human Ecology*, chapter Seasonal Effects on Physical Growth and Human Development, pages 89–106. Cambridge University Press, Cambridge, 1993.
- JH Cross, J Eminson, and BA Wharton. Ramadan and birth weight at full term in asian moslem pregnant women in birmingham. *Archives of Disease in Childhood*, 653:1053–56, 1990.
- E. Dikensoy, O. Balat, B. Cebesoy, A. Ozkur, H. Cicek, and G. Can. The effect of Ramadan fasting on maternal serum lipids, cortisol levels and fetal development. *Archives of gynecology and obstetrics*, 279(2):119, 2009.
- Janet A. DiPietro, Kathleen Costigan, Marc H. Bornstein, Chun-Shin Hahn, and Aristide Achy-Brou. Fetal heart rate and variability: Stability and prediction to developmental outcomes in early childhood. *Child Development*, 78(6):1788–1798, November/December 2007.
- PM Eaton and PA Wharton. *Applied Nutrition*, chapter Sorrento Study of Nutrition in Pregnancy, pages 89–91. John Libbey, London, 1982.
- Reyn Van Ewijk. Long-term health effects on the next generation of ramadan fasting during pregnancy. April 2009. CEP Discussion Paper No 926.
- Mary L. Hediger, Mary D. Overpeck, Andrea McGlynn, Robert J. Kuczmarski, Kurt R. Maurer, and William W. Davis. Growth and fatness at three to six years of age of children born small- or large-for-gestational age. *Pediatrics*, 104(3):e33, 1999.
- Tracy S Herrmann, Anna Maria Siega-Riz, Calvin J Hobel, Chandra Aurora, and Christine Dunkel-Schetter. Prolonged periods without food intake during pregnancy increase risk for elevated maternal corticotropin-releasing hormone concentrations. *American Journal of Obstetrics and Gynecology*, 185(2):403–412, 2001.
- Ann Hoskins. Pregnancy and fasting during ramadan. *British Medical Journal*, 304(6836):1247, May 1992.
- ES Hunter and TW Sadler. D-(-)-beta hydroxybutyrate-induced effects on mouse embryos in vitro. *Teratology*, 36:259–64, 1987.

- R Husain, MT Duncan, SH Cheah, and SL Chng. Effects of fasting in ramadan on tropical asiatic moslems. *British Journal of Nutrition*, 5833(1):41–48, 1987.
- Vincent W.V. Jaddoe and Jacqueline C.M. Witteman. Hypotheses on the fetal origins of adult diseases: Contributions of epidemiological studies. *European Journal of Epidemiology*, 21:91–102, 2006.
- J Joosop and S.L. Yu. A survey of fasting during pregnancy. *Singapore Medical Journal*, 45(12):583–586, 2004.
- Amita Kapoor, Elizabeth Dunn, Alice Kostaki, Marcus H. Andrews, and Stephen G. Matthews. Fetal programming of hypothalamo-pituitary-adrenal function: Prenatal stress and glucocorticoids. *Journal of Physiology*, 572(1):31–44, 2006.
- Zohreh Kavehmanesh and Hassan Abolghasemi. Maternal ramadan fasting and neonatal health. *Journal of Perinatology*, 24:748–750, 2004.
- Sebastian Kraemer. The fragile male. *BMJ*, 321(7276):1609–1612, 2000.
- SJ Langan, IJ Deary IJ, DA Hepburn, and BM Frier. Cumulative cognitive impairment following recurrent severe hypoglycaemia in adult patients with insulin-treated diabetes mellitus. *Diabetologia*, 34:337–344, 1991.
- Melissa A. Larson, Koji Kimura, Michael Kubisch, and R. Michael Roberts. Sexual dimorphism among bovine embryos in their ability to make the transition to expanded blastocyst and in the expression of the signaling molecule ifn-t. *PNAS*, 98(17):9677–9682, 2001.
- Abdulwahab M. Makki. Impact of ramadan fasting on birth weight in 4 hospitals in sanaa city,yemen. *Saudi Medical Journal*, 23(11):1419–1420, 2002.
- Ashok Malhotra, P.H. Scott, J.S. Scott, H. Gee, and B.A. Wharton. Metabolic changes in asian muslim pregnant mothers observing the ramadan fast in britian. *British Journal of Nutrition*, 61(3):663–672, May 1989.
- Kamal Mohmoud Saleh Mansi. Study the effects of ramadan fasting on the serum glucose and lipid profile among healthy jordanian students. *American Journal of Applied Sciences*, 3(8):565–569, 2007.
- Fiona Mathews, Paul J. Johnson, and Andrew Neil. You are what your mother eats: evidence for maternal preconception diet influencing foetal sex in humans. *Proceedings of the Royal Society B*, pages 1–8, April 2008.
- T.J. Mathews and Brady E. Hamilton. Trend analysis of the sex ratio at birth in the united states. *National Vital Statistics Reports*, 53(20), June 14 2005.
- Hisham M. Mirghani, Sarath Weerasinghe, Shamsaa Al-Awar, Lolowa Abdulla, and Mutairu Ezimokhai. The effect of intermittent maternal fasting on computerized fetal heart tracing. *Journal of Perinatology*, 25:90–92, 2005.

- Hisham M. Mirghani, Muntha Salem, and Sarath D. Weerasinghe. Effect of maternal fasting on uterine arterial blood flow. *Journal of Obstetrics and Gynaecology Research*, 33(2):151–154, April 2007.
- H.M. Mirghani and Omer A. Hamud. The effect of maternal diet restriction on pregnancy outcome. *American Journal of Perinatology*, 23(1):21–24, 2006.
- H.M. Mirghani, S.D. Weerasinghe, J.R. Smith, and M. Ezimokhai. The effect of intermittent maternal fasting on human fetal breathing movements. *Journal of Obstetrics and Gynaecology*, 24(6):635–637, September 2004.
- DCP Moore, M Stanisstreet, and CA Clarke. Morphological and physiological effects of beta-hydroxybutyrate on rat embryos grown in vitro at different stages. *Teratology*, 40:237–51, 1989.
- Hidenori Nishina, Lucy R. Green, Hugh H. G. McGarrigle, David E. Noakes, Lucilla Poston, and Mark A. Hanson. Effect of nutritional restriction in early pregnancy on isolated femoral artery function in mid-gestation fetal sheep. *Acta Medica Iranica*, 42(3):209–212, 2004.
- Karen Norberg. Partnership status and the human sex ratio at birth. *Proceedings of the Royal Society B: Biological Sciences*, 271(1555):2403–2410, 2004.
- A.A. Opaneye, D.D. Villegas, and Ali Abdel Azeim. Islamic festivals and low birthweight infants. *The Journal of the Royal Society for the Promotion of Health*, 110(3):106–107, June 1990.
- Rebecca C. Painter, Tessa J. Roseboom, and Otto P. Bleker. You are what your mother eats: evidence for maternal preconception diet influencing foetal sex in humans. *Reproductive Toxicology*, 20:345–352, 2005.
- AM Prentice, A Prentice, WH Lamb, PG Lunn, and S. Austin. Metabolic consequences of fasting during ramadan in pregnant and lactating women. *Human nutrition. Clinical nutrition*, 37(4):283–94, July 1983.
- J Ramadan, G Telahoun, NS Al-Zaid, and M Barac-Nieto. Responses to exercise, fluid, and energy balances during ramadan in sedentary and active males. *Nutrition*, 15(10):735–739, 1999.
- T Rizzo, BE Metzger, WJ Burns, and K Burns. Correlations between antepartum maternal metabolism and child intelligence. *New England Journal of Medicine*, 325:911–916, 1991.
- Trinka Robinson and Jeanne Raisler. Each one is a doctor for herself: Ramadam fasting among pregnant muslim women in the united states. *Ethnicity and Disease*, 15:S1–99 to S1–103, Winter 2005.
- Tessa J. Roseboom, Jan H.P. Meulen, Anita C.J. Ravelli, Clive Osmond, David J.P. Barker, and Otto P. Bleker. Effects of prenatal exposure to the dutch famine on adult disease in later life: An overview. *Twins Research*, 4(5):293–298, October 2001.

- Theresa O Scholl, MaryFran Sowers, Xinhua Chen, and Carine Lenders. Maternal glucose concentration influences fetal growth, gestation and pregnancy complications. *American Journal of Epidemiology*, 154(6):514–520, 2001.
- Shole Shahgheibi, Ebrahim Ghadery, Arash Pauladi, Sabah Hasani, and Syroos Shahsawari. Effects of fasting during the third trimester of pregnancy on neonatal growth indices. *Annals of Alquds Medicine*, 1426(2):58–62, 2005.
- EA Sheehan, F Beck, CA Clarke, and M Stanisstreet. Effects of beta-hydroxybutyrate on rat embryos grown in culture. *Experientia*, 41:273–75, 1985.
- Anna Maria Siega-Riz, Tracy Hermann, David A Savitz, and John M Thorp. Frequency of eating during pregnancy and its effect on preterm delivery. *American Journal of Epidemiology*, 153(7):647–652, 2001.
- Hamed R Takruri. Effect of fasting in ramadan on body weight. *Saudi Medical Journal*, 10(6):491–494, 1989.
- Edith W. M. T. ter Braak, Inge M. Evers, D. Willem Erkelens, and Gerard H. A. Visser. Maternal hypoglycemia during pregnancy in type 1 diabetes: maternal and fetal consequences. *Diabetes/Metabolism Research And Reviews*, 19:96–105, 2002.
- Robert L. Trivers and Dan E. Willard. Natural selection and the ability to vary the sex ratio of offspring. *Science*, 179(4068):90–92, January 1973.

**Table A1: Summary of Hypotheses Concerning Outcomes Affected by Fasting and Timing In Utero**

Outcome	Description of Mechanism (studies)	Gestation month
<b>Birth Outcomes</b>		
Birthweight	Direct effect of low blood glucose (Scholl et al, 2001)	6 to 7
Birthweight	Exposure to ketones, animal studies (Hunter, 1987; Moore, 1989)	1
Birthweight	HPA axis (various studies)	1 to 2
Birthweight	Low birthweight due to shorter gestation (Siega-Riz et al, 2001)	5 to 7
Birthweight	Empirical result --Dutch Famine (Painter et al 2005)	7 to 9
Low Birth Weight	Empirical result (Cross et al 1990; Arab and Nasrollahi, 2001)	4 to 6
Gestation	Fasting associated with high Plasma CRH (Siega-Riz et al, 2001)	5 to 7
NICU	empirical result (Mirghani and Hamud, 2006)	8
C-section	empirical result (Mirghani and Hamud, 2006)	8
Induced Labor	empirical result (Mirghani and Hamud, 2006)	8
Sex Ratio	Effect of low glucose, empirical result (Matthews et al, 2008)	0
<b>Long-Term Outcomes</b>		
Diabetes	Fetal nutrition (various studies)	1 to 3
Heart Disease	Fetal nutrition (various studies)	1 to 6
Cognitive Function	Exposure to ketones, animal studies (Hunter, 1987; Moore, 1989)	1
Cognitive Function	Low blood glucose (Rizzo et al, 1991)	1 to 3
Cognitive Function	HPA axis (Kapoor et al, 2006)	1 to 2
Cognitive Function	Fetal Heart Rate (Mirghani, 2005)	7 to 9
Adult Sex Ratio	HPA axis (Kapoor et al, 2006)	1 to 2

*Notes:* This table is based on a review of selected studies and does not include all relevant studies in the medical literature. Studies include both human and animal studies. In many of the studies, the period of in utero exposure was selected by design and therefore the fact that an effect was found in the chosen gestation period does not rule out possible effects in other periods.

**Table A2: Summary Statistics for Michigan Natality Data, 1989-2005**

	Arab			Non-Arab		
	mean	s.d.	N	mean	s.d.	N
Mother's Age	27.54	5.72	46979	27.41	5.73	1638059
Mother's Education	12.03	3.55	45584	13.18	2.37	1625226
Father's Age	33.81	6.48	45588	30.21	6.13	1462349
Father's Education	12.92	3.33	43931	13.40	2.39	1428050
Male Child	0.52	0.50	46983	0.51	0.50	1638213
Tobacco	0.04	0.19	46203	0.19	0.39	1611440
Alcohol	0.00	0.04	46170	0.02	0.12	1608527
Maternal Weight Gain	29.73	12.70	42216	31.04	13.03	1520595
No Prenatal Care	0.01	0.10	45068	0.01	0.08	1607940
Prenat. Care Begins 1st Trim.	0.86	0.34	45068	0.87	0.34	1607940
Prenat. Care Begins 2nd Trim.	0.10	0.29	45068	0.11	0.31	1607940
Prenat. Care Begins 3rd Trim.	0.03	0.17	45068	0.02	0.13	1607940
Medicaid	0.46	0.50	46315	0.27	0.45	1616231
Fraction Arab, Zipcode	0.21	0.25	46369	0.01	0.03	1612481
Birthweight	3325.08	513.65	46896	3427.71	565.23	1635183
Low Birthweight	0.04	0.21	46988	0.05	0.21	1638244
Infant Death	0.01	0.07	46988	0.01	0.08	1638244
Parity	1.64	1.74	46592	1.39	1.49	1628783
Preterm	0.06	0.23	46868	0.07	0.25	1633654
Gestation (author's calc.)	39.27	1.72	46868	39.29	1.85	1633654
Apgar 5 minute	8.94	0.56	46902	8.94	0.67	1632994
NICU	0.03	0.17	46915	0.04	0.19	1634113
Complication	0.25	0.43	46188	0.28	0.45	1618589
Abnormal Condition	0.06	0.24	46012	0.07	0.25	1611065
Medical Risk	0.20	0.40	46169	0.23	0.42	1618107
Medical Risk Diabetes	0.03	0.16	46169	0.03	0.17	1618107
Born January	0.077	0.27	46988	0.078	0.27	1638244
Born February	0.074	0.26	46988	0.077	0.27	1638244
Born March	0.083	0.28	46988	0.087	0.28	1638244
Born April	0.079	0.27	46988	0.084	0.28	1638244
Born May	0.084	0.28	46988	0.088	0.28	1638244
Born June	0.087	0.28	46988	0.086	0.28	1638244
Born July	0.089	0.29	46988	0.089	0.28	1638244
Born August	0.091	0.29	46988	0.088	0.28	1638244
Born September	0.087	0.28	46988	0.085	0.28	1638244
Born October	0.084	0.28	46988	0.083	0.28	1638244
Born November	0.081	0.27	46988	0.076	0.27	1638244
Born December	0.083	0.28	46988	0.078	0.27	1638244
Mo. 1, Ram Hrs Exposure	0.056	0.15	46868	0.056	0.15	1633654
Mo. 2, Ram Hrs Exposure	0.059	0.15	46868	0.058	0.16	1633654
Mo. 3, Ram Hrs Exposure	0.058	0.15	46868	0.059	0.16	1633654
Mo. 4, Ram Hrs Exposure	0.059	0.15	46868	0.060	0.16	1633654
Mo. 5, Ram Hrs Exposure	0.057	0.15	46868	0.060	0.16	1633654
Mo. 6, Ram Hrs Exposure	0.056	0.15	46868	0.060	0.16	1633654
Mo. 7, Ram Hrs Exposure	0.056	0.15	46868	0.061	0.16	1633648
Mo. 8, Ram Hrs Exposure	0.057	0.15	46865	0.061	0.16	1633617
Mo. 9, Ram Hrs Exposure	0.059	0.16	46861	0.060	0.16	1633475



**Table A3: Summary Statistics for Uganda Census Sample**

	Muslim			Non-Muslim		
	mean	s.d.	N	mean	s.d.	N
female	0.494	0.500	81197	0.498	0.500	643300
age	34.546	12.675	81197	36.697	13.907	643300
illiterate	0.304	0.460	78990	0.356	0.479	626473
years of schooling	6.944	3.269	60117	6.797	3.599	449968
no schooling	0.247	0.431	80142	0.290	0.454	635282
employed	0.660	0.474	74348	0.631	0.483	581842
elementary occupation	0.042	0.200	46284	0.042	0.200	347248
home ownership (males)						
# of wives (males)						
disability	0.0380	0.191	80924	0.0521	0.222	640825
blind/vision impaired	0.0106	0.102	80922	0.0149	0.121	640789
deaf/hearing impaired	0.0038	0.062	80923	0.0061	0.078	640781
mute/speech impaired	0.0009	0.030	80921	0.0015	0.038	640780
lower extremities	0.0125	0.111	80921	0.0161	0.126	640794
upper extremities	0.0039	0.062	80921	0.0056	0.075	640779
mental/learning	0.0014	0.037	80921	0.0017	0.041	640777
psychological	0.0014	0.038	80921	0.0020	0.045	640776
epilepsy	0.0005	0.023	80921	0.0009	0.031	640777
rheumatism	0.0009	0.030	80921	0.0016	0.039	640776
congen	0.0050	0.070	80921	0.0058	0.076	640778
disease	0.0203	0.141	80924	0.0283	0.166	640803
accident	0.0056	0.074	80921	0.0079	0.088	640782
occupational injury	0.0053	0.072	80921	0.0074	0.086	640786
war_injury	0.0007	0.027	80921	0.0013	0.036	640777
aging	0.0053	0.072	80921	0.0074	0.086	640786
Born January	0.105	0.306	81197	0.096	0.294	643300
Born February	0.076	0.265	81197	0.075	0.263	643300
Born March	0.072	0.258	81197	0.072	0.259	643300
Born April	0.110	0.313	81197	0.106	0.308	643300
Born May	0.070	0.256	81197	0.070	0.256	643300
Born June	0.102	0.302	81197	0.105	0.307	643300
Born July	0.094	0.292	81197	0.098	0.298	643300
Born August	0.079	0.269	81197	0.083	0.275	643300
Born September	0.079	0.269	81197	0.081	0.272	643300
Born October	0.078	0.268	81197	0.077	0.267	643300
Born November	0.069	0.253	81197	0.069	0.253	643300
Born December	0.067	0.250	81197	0.068	0.251	643300
rampct1	0.081	0.215	81197	0.081	0.216	643300
rampct2	0.079	0.214	81197	0.079	0.215	643300
rampct3	0.077	0.211	81197	0.078	0.212	643300
rampct4	0.084	0.219	81197	0.083	0.218	643300
rampct5	0.086	0.223	81197	0.085	0.221	643300
rampct6	0.084	0.217	81197	0.083	0.217	643300
rampct7	0.087	0.222	81197	0.085	0.221	643300
rampct8	0.090	0.226	81197	0.089	0.226	643300
rampct9	0.087	0.221	81197	0.087	0.221	643300

**Table A4: Effects of Ramadan Exposure on Birthweight and Gestation, by Varying Gestation, Michigan Arabs**

Gestation Month exposure	<i>Dependent variable is Birthweight</i>					<i>Dependent variable is Weeks of Gestation</i>				
	(1) 25-42 weeks	(2) 31-42 weeks	(3) 35-42 weeks	(4) 39-42 weeks	(5) Non-Chaldean 39-42 wks	(6) 25-42 weeks	(7) 31-42 weeks	(8) 35-42 weeks	(9) 39-42 weeks	(10) Non-Chaldean 39-42 wks
1	-31.7 * (18.4)	-32.4 * (17.7)	-41.6 ** (17.1)	-40.6 ** (18.9)	-35.4 (22.4)	-0.02 (0.06)	-0.03 (0.06)	-0.06 (0.05)	-0.01 (0.03)	-0.01 (0.04)
2	-43.2 ** (17.6)	-37.1 ** (16.9)	-29.2 * (16.3)	-33.6 * (17.9)	-46.5 ** (21.3)	-0.11 * (0.06)	-0.07 (0.05)	-0.04 (0.05)	-0.05 * (0.03)	-0.10 *** (0.04)
3	7.8 (18.2)	7.9 (17.4)	-7.5 (16.8)	-11.3 (18.4)	-21.1 (22.0)	0.02 (0.06)	0.03 (0.05)	-0.04 (0.05)	-0.07 ** (0.03)	-0.06 (0.04)
4	-11.3 (18.2)	-12.1 (17.4)	-11.7 (16.8)	-33.0 * (18.6)	-25.8 (22.3)	-0.03 (0.06)	-0.03 (0.05)	-0.04 (0.05)	-0.03 (0.03)	-0.04 (0.04)
5	-24.7 (18.5)	-15.7 (17.8)	-20.3 (17.2)	-23.7 (19.0)	-39.9 * (22.7)	-0.12 * (0.06)	-0.08 (0.06)	-0.08 * (0.05)	-0.06 * (0.03)	-0.08 * (0.04)
6	<i>-30.6</i> * (18.5)	<i>-31.6</i> * (17.8)	<i>-30.8</i> * (17.2)	-14.4 (19.0)	-25.8 (22.7)	<i>-0.08</i> (0.06)	-0.09 (0.06)	-0.08 (0.05)	-0.02 (0.03)	-0.03 (0.04)
7	-22.7 (18.2)	-18.2 (17.5)	-20.7 (16.9)	-29.5 (18.5)	-56.8 *** (21.8)	<i>0.07</i> (0.06)	0.10 * (0.05)	0.08 (0.05)	0.03 (0.03)	0.01 (0.04)
8	<i>32.8</i> * (17.6)	<i>28.5</i> * (16.8)	17.6 (16.2)	9.7 (17.9)	26.9 (21.3)	<i>0.08</i> (0.06)	<i>0.06</i> (0.05)	0.01 (0.05)	-0.06 * (0.03)	-0.07 * (0.04)
9	-19.9 (17.9)	-20.7 (17.2)	-18.4 (16.6)	-22.1 (18.2)	-26.6 (21.6)	<i>0.01</i> (0.06)	<i>0.00</i> (0.05)	<i>0.02</i> (0.05)	-0.01 (0.03)	-0.02 (0.04)
joint test, all shaded coeff equal to 0										
p-value	0.065	0.124	0.061	0.277	0.071	0.262	0.091	0.116	0.160	0.143
joint test, all shaded coeff are equal										
p-value	0.156	0.330	0.114	0.830	0.207	0.356	0.069	0.098	0.316	0.430
N	43626	43403	42874	30298	21890	43680	43619	43454	42923	21915

*Notes:* All entries are coefficients on Ramadan exposure to daylight hours over subsequent 30 days as fraction of peak daylight hours during sample period. Gestation is either self reported or physician estimated (see text). Shaded areas show coefficients that are relevant for given sample. Columns (5) and (10) exclude zipcodes with a Chaldean/Arab ratio  $\geq 0.5$ . Controls include mothers age, mother's age squared, mother's education, tobacco use, alcohol use, parity, father's education, dummy for missing father's education, father's age, father's age squared, number of previous pregnancies that resulted in death at birth, conception month dummies, county dummies and birth year dummies. Italicized estimates indicate that the estimate may represent post natal exposure for some sample members. Standard errors in parentheses, \*significant at 10%; \*\* significant at 5%; \*\*\* significant at 1%

**Table A5: Effects of Ramadan Exposure on Other Birth Outcomes by Gestation Month, Michigan Arabs**

Gestation Month exposure	(1) Low Birth Weight	(2) Infant Death Perinatal	(3) 5-minute APGAR score	(4) Maternal Weight Gain	(5) Neonatal Intensive Care Unit	(6) C-section	(7) Pregnancy Complication	(8) Abnormal Conditions	(9) Congenital Anomaly
1	-0.012 (0.009)	0.000 (0.001)	0.008 (0.021)	0.613 (0.549)	-0.019** (0.008)	-0.008 (0.015)	-0.033* (0.018)	-0.003 (0.010)	0.005 (0.006)
2	0.016* (0.009)	0.001 (0.001)	-0.047** (0.020)	0.380 (0.523)	0.008 (0.007)	0.013 (0.015)	0.040** (0.017)	-0.006 (0.010)	-0.006 (0.006)
3	-0.01 (0.009)	0.003** (0.001)	-0.015 (0.020)	-0.286 (0.543)	-0.006 (0.008)	-0.02 (0.015)	0.000 (0.018)	0.000 (0.010)	-0.007 (0.006)
4	-0.003 (0.009)	0.000 (0.001)	-0.008 (0.021)	-0.002 (0.546)	-0.013* (0.008)	0.011 (0.015)	-0.009 (0.018)	-0.018* (0.010)	-0.001 (0.006)
5	-0.004 (0.009)	0.001 (0.001)	0.004 (0.021)	0.27 (0.551)	-0.007 (0.008)	-0.026* (0.016)	0.015 (0.018)	-0.018* (0.011)	-0.003 (0.006)
6	0.001 (0.009)	0.000 (0.001)	-0.050** (0.021)	-0.006 (0.553)	-0.001 (0.008)	0.008 (0.016)	-0.001 (0.018)	0.015 (0.011)	0.005 (0.006)
7	0.003 (0.009)	0.000 (0.001)	-0.011 (0.020)	-1.085** (0.540)	-0.006 (0.007)	0.001 (0.015)	-0.011 (0.018)	-0.006 (0.010)	-0.006 (0.006)
8	-0.020** (0.009)	0.001 (0.001)	0.010 (0.020)	0.172 (0.524)	-0.004 (0.007)	-0.019 (0.015)	-0.024 (0.017)	-0.004 (0.010)	0.003 (0.006)
9	0.004 (0.009)	-0.001 (0.001)	-0.032 (0.020)	-0.387 (0.532)	0.008 (0.007)	0.011 (0.015)	0.007 (0.018)	0.017* (0.010)	0.008 (0.006)
joint test, coefficients on months 1 to 9 equal to 0									
p-value	0.170	0.410	0.120	0.440	0.110	0.550	0.120	0.090	0.370
joint test, coefficients on months 1 to 9 are equal									
p-value	0.130	0.400	0.160	0.350	0.090	0.470	0.080	0.060	0.280
N	31857	31857	31817	29285	31824	31726	31745	31709	31710

*Notes:* All entries are coefficients on Ramadan exposure to daylight hours over subsequent 30 days as fraction of peak daylight hours during sample period. Gestation is either self reported or physician estimated. All samples use births with gestation between 25 and 42 weeks. All regressions include controls for mothers age, mother's age squared, mother's education, tobacco use, alcohol use, parity, father's education, dummy for missing father's education, father's age, father's age squared, number of previous pregnancies that resulted in death at birth, conception month dummies, county dummies and birth year dummies. The sample drops Arab mothers living in zipcodes with a large share of Chaldeans. Italicized estimates indicate that the estimate may represent post natal exposure for some sample members. Standard errors in parentheses, \*significant at 10%; \*\* significant at 5%; \*\*\* significant at 1%

**Table A6: Effects of Ramadan Exposure on Disability Outcomes, Ugandan Muslims by Months Prior to Birth**

Months Prior to Birth	Disability	Sight/Blind	Hearing/Deaf	Mental/ Learning	Psychological	Lower Extremities	Upper Extremities
10	-0.127 (0.403)	0.000 (0.216)	-0.080 (0.132)	0.028 (0.079)	0.038 (0.080)	-0.112 (0.236)	0.032 (0.133)
9	0.793** (0.368)	0.349* (0.197)	0.227* (0.120)	0.255*** (0.073)	-0.091 (0.073)	0.312 (0.216)	-0.064 (0.122)
8	0.035 (0.375)	-0.078 (0.201)	0.129 (0.123)	0.115 (0.074)	-0.052 (0.075)	0.011 (0.220)	0.021 (0.124)
7	-0.176 (0.375)	-0.022 (0.201)	0.102 (0.123)	0.038 (0.074)	0.071 (0.075)	-0.355 (0.220)	0.088 (0.124)
6	0.152 (0.382)	0.074 (0.205)	0.132 (0.125)	0.11 (0.075)	-0.084 (0.076)	-0.191 (0.224)	0.096 (0.126)
5	0.04 (0.376)	-0.004 (0.202)	0.168 (0.123)	0.139* (0.074)	-0.044 (0.075)	-0.085 (0.221)	-0.025 (0.125)
4	0.229 (0.379)	0.039 (0.203)	0.044 (0.124)	0.127* (0.075)	-0.036 (0.076)	-0.007 (0.222)	0.19 (0.126)
3	0.059 (0.391)	0.124 (0.210)	0.071 (0.128)	0.049 (0.077)	0.004 (0.078)	-0.158 (0.229)	-0.017 (0.130)
2	-0.307 (0.373)	-0.272 (0.200)	0 (0.122)	0.153** (0.074)	-0.007 (0.074)	-0.095 (0.219)	0.012 (0.124)
1	-0.158 (0.405)	0.018 (0.217)	0.052 (0.132)	0.101 (0.080)	-0.017 (0.081)	-0.106 (0.237)	-0.042 (0.134)
joint test, coefficients on months 1 to 9 equal to 0							
<i>p</i> -value	0.380	0.580	0.680	0.060	0.780	0.370	0.830
<i>Mean</i>	3.80%	1.06%	0.38%	0.14%	0.14%	1.25%	0.39%
<i>N</i>	80924	80922	80923	80921	80921	80921	80921

*Notes:* All entries are coefficients on Ramadan exposure measured as the percent of days overlapping with Ramadan during the nine months preceding birth (rampct). Each outcome is multiplied by 100, so that coefficients are in units of percentage points. All regressions include an indicator for female, birth month dummies, district of birth dummies and birth year dummies.

Table A7: Effects of Ramadan Exposure on Causes of Disabilities, Ugandan Muslims, by Months Prior to Birth

Months Prior to Birth	<i>Unrelated to prenatal nutrition</i>			<i>Possibly Related to prenatal nutrition</i>		
	Accident	Occ. Injury	War Injury	Aging	Disease	Congenital
9	-0.060 (0.142)	0.059 (0.074)	0.054 (0.052)	0.373*** (0.136)	0.199 (0.267)	0.059 (0.074)
8	0.042 (0.133)	-0.023 (0.070)	0.001 (0.049)	0.137 (0.127)	-0.025 (0.250)	-0.023 (0.070)
7	-0.102 (0.137)	-0.063 (0.072)	0.000 (0.050)	-0.034 (0.132)	-0.248 (0.259)	-0.063 (0.072)
6	-0.025 (0.139)	0.050 (0.073)	0.043 (0.051)	0.222* (0.134)	-0.369 (0.262)	0.050 (0.073)
5	0.127 (0.137)	-0.009 (0.072)	-0.085* (0.050)	-0.022 (0.132)	0.100 (0.258)	-0.009 (0.072)
4	0.179 (0.139)	0.018 (0.073)	0.064 (0.051)	0.055 (0.133)	-0.252 (0.261)	0.018 (0.073)
3	-0.09 (0.144)	0.031 (0.075)	0.047 (0.053)	0.110 (0.138)	0.006 (0.270)	0.031 (0.075)
2	0.161 (0.138)	-0.063 (0.072)	0.021 (0.050)	-0.011 (0.132)	-0.158 (0.259)	-0.063 (0.072)
1	0.002 (0.144)	-0.086 (0.076)	0.057 (0.053)	0.051 (0.138)	-0.044 (0.271)	-0.086 (0.076)
joint test, coefficients on months 1 to 9 equal to 0						
<i>p</i> -value	0.710	0.730	0.460	0.210	0.750	0.730
joint test, coefficients on months 1 to 9 are equal						
<i>p</i> -value	0.640	0.640	0.400	0.210	0.730	0.640
<i>Mean</i>	0.56%	0.53%	0.07%	0.53%	2.03%	0.50%
<i>N</i>	80924	80922	80923	80921	80921	80921

*Notes:* All entries are coefficients on Ramadan exposure measured as the percent of days overlapping with Ramadan during the nine months preceding birth (rampct). Each outcome is multiplied by 100, so that coefficients are in units of percentage points. All regressions include an indicator for female, birth month dummies, district of birth dummies and birth year dummies.

Figure A1: Women's Weight Change Around Ramadan in Gambia





Fig. 8.5. Weight change in Gambian women during 3 months around the fast of Ramadan, expressed relative to mean weight for the other 9 months of the year. Each point has a standard error of about 0.15 kg. The data are adjusted for calendar month and year of measurement, and stage of pregnancy/lactation, using within-subject regression.


Source: Cole (1993)

Figure A2: Michigan Arab Population by Zipcode

Panel A: Quartiles of the Arab Population




Panel B: Ratio of the Chaldean to Arab Population



Source: Author's calculations using the 2000 Census SF3 file

**Figure A3:  
First Gestation Month Exposure to Ramadan**





**Figure A4:  
Distributional Effect of First Month Exposure on Birthweight**

