

NBER WORKING PAPER SERIES

ENDOGENOUS STRATIFICATION IN RANDOMIZED EXPERIMENTS

Alberto Abadie
Matthew M. Chingos
Martin R. West

Working Paper 19742
<http://www.nber.org/papers/w19742>

NATIONAL BUREAU OF ECONOMIC RESEARCH
1050 Massachusetts Avenue
Cambridge, MA 02138
December 2013

We thank Beth Akers, Josh Angrist, Gary Chamberlain, David Deming, Josh Goodman, Guido Imbens, Max Kasy, Larry Katz, Amanda Pallais, Paul Peterson, and Russ Whitehurst for helpful discussions. The views expressed herein are those of the authors and do not necessarily reflect the views of the National Bureau of Economic Research.

NBER working papers are circulated for discussion and comment purposes. They have not been peer-reviewed or been subject to the review by the NBER Board of Directors that accompanies official NBER publications.

© 2013 by Alberto Abadie, Matthew M. Chingos, and Martin R. West. All rights reserved. Short sections of text, not to exceed two paragraphs, may be quoted without explicit permission provided that full credit, including © notice, is given to the source.

Endogenous Stratification in Randomized Experiments
Alberto Abadie, Matthew M. Chingos, and Martin R. West
NBER Working Paper No. 19742
December 2013
JEL No. C01,C21,C9

ABSTRACT

Researchers and policy makers are often interested in estimating how treatments or policy interventions affect the outcomes of those most in need of help. This concern has motivated the increasingly common practice of disaggregating experimental data by groups constructed on the basis of an index of baseline characteristics that predicts the values that individual outcomes would take on in the absence of the treatment. This article shows that substantial biases may arise in practice if the index is estimated, as is often the case, by regressing the outcome variable on baseline characteristics for the full sample of experimental controls. We analyze the behavior of leave-one-out and repeated split sample estimators and show that in realistic scenarios they have substantially lower biases than the full sample estimator. We use data from the National JTPA Study and the Tennessee STAR experiment to demonstrate the performance of alternative estimators and the magnitude of their biases.

Alberto Abadie
John F. Kennedy School of Government
Harvard University
79 JFK Street
Cambridge, MA 02138
and NBER
alberto_abadie@harvard.edu

Martin R. West
Harvard Graduate School of Education
6 Appian Way, Gutman 454
Cambridge, MA 02138
martin_west@gse.harvard.edu

Matthew M. Chingos
Brown Center on Education Policy
Brookings Institution
1775 Massachusetts Ave. NW
Washington, DC 20036
mchingos@brookings.edu

1. Introduction

Recent years have seen rapid growth in the use of randomized experiments in social science research. In part, this has been motivated by the “credibility revolution” in which researchers have devoted much attention to the study of the conditions that allow estimation of treatment effects (Angrist and Pischke 2010; Murnane and Willett 2011). The main advantage of a large and well-executed randomized experiment is that the researcher can confidently rule out the possibility that unobserved differences between the treatment and control groups could explain the results of the study.

In addition to allowing estimation of average treatment effects for the entire population subjected to random assignment, experiments also make it possible to obtain unbiased estimates of treatment effects for subgroups of study participants. Subgroup effects are of particular interest to policymakers seeking to target policies on those most likely to benefit. As a general rule, subgroups must be created based on characteristics that are either immutable (e.g., race) or observed before randomization (e.g., on a baseline survey) so that they could not possibly have been affected by the treatment.

However, many researchers and policy makers are interested in estimating how treatments affect those most in need of help, that is, those who would attain extremely unfavorable outcomes in the absence of the treatment. Treatment parameters of this nature depend on the joint distribution of potential outcomes with and without treatment, which is not identified by randomization (see, e.g., Heckman et al., 1997). A solution to this problem is to combine baseline characteristics into a single index that reflects each participant’s predicted outcome and conduct separate analysis for subgroups of participants defined in terms of intervals of the predicted outcome.

A common implementation of this idea is the use of data on out-of-sample untreated units to estimate a prediction model for the outcome variable, which can then be applied to predict outcomes without treatment for the experimental units. This approach is common in medical research, where validated risk models are often available to stratify experimental subjects based on their predicted probability of a given health outcome (Kent and Hayward

2007). It is also the approach taken by Deming (2011) in his lottery-based study of the impact of enrollment in a choice school on criminal activity. His treatment effect estimates are based on a group of students that participated in a school choice lottery, but his *ex-ante* predicted outcomes for both the treatment and control groups are based on a group of students from the same school district who did not participate in the lottery.

However, experimental studies in the social sciences often lack data on a non-experimental control group that can be employed to predict the outcome that experimental units would attain in the absence of the treatment. A potential approach to this problem that is gaining popularity among empirical researchers is to use in-sample information on the relationship between the outcome of interest and covariates for the experimental controls to estimate potential outcomes without the treatment for all experimental units. We call this practice endogenous stratification, because it uses in-sample data on the outcome variable to stratify the sample.

Endogenous stratification is typically implemented in practice by first regressing the outcome variable on baseline characteristics using the full sample of experimental controls, and then using the coefficients from this regression to generate predicted potential outcomes without treatment for all sample units.

Unfortunately, as we demonstrate below, this procedure generates estimators of treatment effects that are substantially biased, and the bias follows a predictable pattern: results are biased upward for individuals with low predicted outcomes and biased downward for individuals with high predicted outcomes.

This bias pattern matches the results of several recent experimental studies that use this procedure and estimate strong positive effects for individuals with low predicted outcomes; in some cases, the studies also report negative effects for individuals with high predicted outcomes. For example, a 2011 working paper by Goldrick-Rab et al. reports that a Wisconsin need-based financial aid program for post-secondary education had no overall impacts on college enrollment or college persistence among eligible students as a whole. Looking separately at subgroups based on predicted persistence, however, the study finds

large positive effects on enrollment after three years for students in the bottom third of predicted persistence and equally large negative effects for students in the top third of predicted persistence.¹ A 2011 working paper by Dynarski et al. analyzing long-term impacts of the Project STAR experiment similarly finds that assignment to a small class in grades K-3 increased college enrollment rates among the quintile of students with the lowest ex-ante probability to enroll by 11 percentage points, but had no impact on students in the top four quintiles.² In a recent working paper prepared for the Institute of Education Sciences, Pane et al. (2013) report that the effect of a technology-based algebra curriculum on the test scores of middle and high school students differs significantly across quintiles of predicted test scores, with “potentially moderately large positive treatment effects in the lowest quintile and small negative effects of treatment in the highest two quintiles” (p. 21). Hemelt et al. (2012) find that an elementary school intervention with no average impact on college enrollment and degree receipt increased two-year college enrollment by 7 percentage points and associate’s degree completion by 3 percentage points for students in the bottom quartile of ex-ante probability of college-attendance.³ Rodriguez-Planas (2012) reports that a mentoring program for adolescents reduced risky behavior and improved educational attainment for students in the top half of the risk distribution but increased risky behavior in the bottom half.

Endogenous stratification also plays a supporting role in Angrist and Lavy’s (2009) experimental evaluation of a cash incentive program aimed at increasing matriculation

¹Table 5 of Goldrick-Rab et al. (2011) suggests that the grant increased full-time enrollment after three years by 17.0 percentage points for the bottom third but reduced full-time enrollment by 15.1 percentage points for the top third. Although the authors describe their subgroup analyses as “exploratory” and eliminated results based on endogenous stratification from a subsequent working paper on the same grant program (Goldrick-Rab et al. 2012), their initial findings received widespread media coverage, including articles in the *Chronicle of Higher Education*, *Inside Higher Ed*, and *Education Week*. In a related paper on the design of randomized experiments, Harris and Goldrick-Rab (2012) discuss potential explanations for the unexpected heterogeneity in their impact estimates based on full-sample endogenous stratification.

²The authors removed results based on endogenous stratification estimator from their published paper (Dynarski et al. 2013), which shows that assignment to a small class increased college enrollment (completion) by 2.7 (1.6) percentage points for the full Project STAR sample, with larger effects for African Americans.

³Hemelt et al. (2012) show that the same intervention increased two-year college enrollment by 4.8 and 6.3 percentage points for African American and low-income students, respectively.

certification rates for Israeli high school students. In order to test whether the program was most effective for girls on the certification margin, the researchers first group students by baseline test scores. They also, however, report results for students grouped by ex-ante certification probability based on a broader set of background characteristics as “a check on the notion that high lagged scores identify students who have a shot at classification” (p. 1396).

Finally, Giné et al. (2012) find that fingerprinting loan applicants in Malawi increased repayment rates only among the borrowers in the top quintile of predicted default risk. In contrast to many other studies that employ endogenous stratification, Giné et al. (2012) recognize that in-sample estimation of predicted outcomes may generate biases and provide evidence that their results are not artificially generated by those biases. This should indeed be a concern in empirical studies that use endogenous stratification. As we demonstrate in Section 3 using data from the National JTPA study and the Tennessee STAR experiment, biases resulting from endogenous stratification can completely alter the quantitative and qualitative conclusion of empirical studies.

In the remainder of this article, we first describe in more detail the increasingly popular procedure of stratifying experimental data by groups constructed on the basis of the predicted values from a regression of the outcome on baseline covariates for the full sample of experimental controls. We next explain why this method generates biases and describe the direction of those biases. We then describe leave-one-out and repeated split sample procedures that generate consistent estimators and show that the biases of these estimators are substantially lower than the bias of the full sample estimator in two realistic scenarios. We use data from the National JTPA Study and the Tennessee STAR experiment to demonstrate the performance of endogenous stratification estimators and the magnitude of their biases. We restrict our attention to randomized experiments, because this is the setting where endogenous stratification is typically used. However, similarly large biases may arise from endogenous stratification in observational studies.

2. Using Control Group Data to Create Predicted Outcomes

We begin by describing in detail the endogenous stratification method outlined above, which aims to classify study participants into groups based on their predicted value of the outcome variable in the absence of the treatment. Suppose that the sample consists of N observations of the triple (y, w, \mathbf{x}) , where y is an outcome variable, w is the treatment, and \mathbf{x} is a vector of baseline characteristics. When the object of interest is the average treatment effect, which in a randomized experiment is equal to $\tau = E[y|w = 1] - E[y|w = 0]$, researchers typically compare sample average outcomes for the treated and the control groups:

$$\hat{\tau} = \frac{\sum_{i=1}^N y_i w_i}{\sum_{i=1}^N w_i} - \frac{\sum_{i=1}^N y_i (1 - w_i)}{\sum_{i=1}^N (1 - w_i)}.$$

As discussed above, researchers sometimes aim to compare treated and non-treated after stratifying on a predictor of the outcome in the absence of the treatment. To our knowledge, most studies that use endogenous stratification implement it roughly as follows:

- (1) Regress the outcome variable on a set of baseline characteristics using the control group only. The regression coefficients are:

$$\hat{\boldsymbol{\beta}} = \left(\sum_{i=1}^N \mathbf{x}_i (1 - w_i) \mathbf{x}_i' \right)^{-1} \sum_{i=1}^N \mathbf{x}_i (1 - w_i) y_i.$$

- (2) Use the estimated coefficients to generate a predicted outcome value for all participants (both treatment and control groups), $\mathbf{x}_i' \hat{\boldsymbol{\beta}}$.
- (3) Divide participants into groups based on their predicted outcomes. Typically, unit i is assigned to group k if $\mathbf{x}_i' \hat{\boldsymbol{\beta}}$ falls in some interval delimited by c_{k-1} and c_k . The interval limits may be fixed or could be quantiles of the empirical distribution of $\mathbf{x}_i' \hat{\boldsymbol{\beta}}$. Many authors use a three-bin classification scheme of low, medium, and high predicted outcomes.

(4) Estimate treatment effects for each of the subgroups,

$$\hat{\tau}_k = \frac{\sum_{i=1}^N y_i I_{[w_i=1, c_{k-1} < \mathbf{x}'_i \hat{\beta} \leq c_k]}}{\sum_{i=1}^N I_{[w_i=1, c_{k-1} < \mathbf{x}'_i \hat{\beta} \leq c_k]}} - \frac{\sum_{i=1}^N y_i I_{[w_i=0, c_{k-1} < \mathbf{x}'_i \hat{\beta} \leq c_k]}}{\sum_{i=1}^N I_{[w_i=0, c_{k-1} < \mathbf{x}'_i \hat{\beta} \leq c_k]}},$$

where I_A is the indicator function that takes values one if event A is realized, and value zero otherwise. Alternatively, treatment effect estimates could be computed after controlling for a set of covariates using regression.

For example, Goldrick-Rab et al. (2011) in their study of the impact of a need-based grant regress college persistence on baseline characteristics using only observations from the control group, generate predicted probabilities of college persistence for all students, classify students into three equal-sized groups based on their ex-ante predicted probability, and then estimate treatment effects for each of the three groups.

This is a simple and direct approach to stratification, which has great intuitive appeal. Moreover, it is easy to show that under usual regularity conditions, $\hat{\tau}_k$ converges to

$$\tau_k = E[y|w = 1, c_{k-1} < \mathbf{x}'\beta \leq c_k] - E[y|w = 0, c_{k-1} < \mathbf{x}'\beta \leq c_k].$$

As we will see next, however, $\hat{\tau}_k$ is biased in finite samples, and the bias follows a predictable pattern.

To simplify the exposition, suppose that predicted outcomes are divided into three groups (low, medium, high). Let $\beta = (E[\mathbf{x}\mathbf{x}'|w = 0])^{-1}E[\mathbf{x}y|w = 0]$ be the population counterpart of $\hat{\beta}$, and let $e_i = y_i - \mathbf{x}'_i \beta$ be the regression error. In a finite sample, untreated observations with large negative values for e_i tend to be over-fitted, so we expect $\mathbf{x}'_i \hat{\beta} < \mathbf{x}'_i \beta$, which pushes these observations towards the lower interval of predicted outcomes. This creates a negative bias in the average outcome among control observations that fall into the lower interval for $\mathbf{x}'_i \hat{\beta}$ and, therefore, a positive bias in the average treatment effect estimated for that group. Analogously, average treatment effect estimators for the upper intervals of predicted outcomes are biased downward. Endogenous stratification results in a

predictable pattern: average treatment effect estimators are biased upward for individuals with low predicted outcomes and biased downward for individuals with high predicted outcomes. As we will demonstrate below, because the finite sample bias of the endogenous stratification estimator is created by over-fitting, this bias tends to be more pronounced when the number of observations is small and the dimensionality of \mathbf{x}_i is large.

The possibility of bias arising from endogenous stratification has been previously recognized in the evaluation literature (see, e.g. Peck, 2003), in statistics (Hansen, 2008), and in economics (Sanbonmatsu et al., 2006, and Giné et al., 2012). A deceptively comforting property of this bias is that it vanishes as the sample size increases, under weak regularity conditions. The present article shows, however, that in realistic scenarios endogenous stratification bias can be substantial and completely drive the quantitative and qualitative conclusions of an empirical study.

A natural solution to the over-fitting issue is provided by leave-one-out estimators.⁴ Let

$$\widehat{\boldsymbol{\beta}}_{(-i)} = \left(\sum_{j \neq i} \mathbf{x}_i (1 - w_i) \mathbf{x}'_i \right)^{-1} \sum_{j \neq i} \mathbf{x}_i (1 - w_i) y_j,$$

be the regression coefficients estimators that discard observation i . Over-fitting is precluded by not allowing the outcome, y_i , of each observation to contribute to the estimation of its own predicted value, $\mathbf{x}'_i \widehat{\boldsymbol{\beta}}_{(-i)}$. Because only untreated observations are employed in the estimation of $\widehat{\boldsymbol{\beta}}_{(-i)}$ and $\widehat{\boldsymbol{\beta}}$, if i is a treated observation then $\widehat{\boldsymbol{\beta}}_{(-i)} = \widehat{\boldsymbol{\beta}}$. We consider the following leave-one-out estimator of τ_k :

$$\widehat{\tau}_k^{LOO} = \frac{\sum_{i=1}^N y_i I_{[w_i=1, c_{k-1} < \mathbf{x}'_i \widehat{\boldsymbol{\beta}} \leq c_k]}}{\sum_{i=1}^N I_{[w_i=1, c_{k-1} < \mathbf{x}'_i \widehat{\boldsymbol{\beta}} \leq c_k]}} - \frac{\sum_{i=1}^N y_i I_{[w_i=0, c_{k-1} < \mathbf{x}'_i \widehat{\boldsymbol{\beta}}_{(-i)} \leq c_k]}}{\sum_{i=1}^N I_{[w_i=0, c_{k-1} < \mathbf{x}'_i \widehat{\boldsymbol{\beta}}_{(-i)} \leq c_k]}}.$$

Under weak assumptions, it can be seen that $\widehat{\boldsymbol{\beta}}$ is a consistent estimator of $\boldsymbol{\beta}$ and the maximal difference between $\widehat{\boldsymbol{\beta}}_{(-i)}$ and $\widehat{\boldsymbol{\beta}}$ converges in probability to zero. As a result, both $\widehat{\tau}_k$ and $\widehat{\tau}_k^{LOO}$ are consistent estimators of τ_k . Moreover, $\widehat{\tau}_k$ and $\widehat{\tau}_k^{LOO}$ have the same large

⁴This is the approach followed in Sanbonmatsu et al. (2006). Harvill et al. (2013) propose a variant of this approach based on 10-fold cross-validation.

sample distribution.⁵ However, we show in section 4 that $\hat{\tau}_k$ is substantially biased in two realistic scenarios, while $\hat{\tau}_k^{LOO}$ is not.⁶

Another way to avoid over-fitting is sample splitting. We consider a repeated split sample estimator. In each repetition, m , the untreated sample is randomly divided into two groups, which we will call the prediction and the estimation groups. Let $v_{im} = 0$ if untreated observation i is assigned the prediction group in repetition m , and $v_{im} = 1$ if it is assigned to the estimation group. In each repetition, m , we estimate β using only the observations in the prediction group:

$$\hat{\beta}_m = \left(\sum_{i=1}^N \mathbf{x}_i (1 - w_i) (1 - v_{im}) \mathbf{x}'_i \right)^{-1} \sum_{i=1}^N \mathbf{x}_i (1 - w_i) (1 - v_{im}) y_i.$$

For each repetition, m , the split sample estimator of τ_k is

$$\hat{\tau}_{km}^{SS} = \frac{\sum_{i=1}^N y_i I_{[w_i=1, c_{k-1} < \mathbf{x}'_i \hat{\beta}_m \leq c_k]}}{\sum_{i=1}^N I_{[w_i=1, c_{k-1} < \mathbf{x}'_i \hat{\beta}_m \leq c_k]}} - \frac{\sum_{i=1}^N y_i I_{[w_i=0, v_{im}=1, c_{k-1} < \mathbf{x}'_i \hat{\beta}_m \leq c_k]}}{\sum_{i=1}^N I_{[w_i=0, v_{im}=1, c_{k-1} < \mathbf{x}'_i \hat{\beta}_m \leq c_k]}}.$$

We then average $\hat{\tau}_{km}^{SS}$ over M repetitions to obtain the repeated split sample estimator:

$$\hat{\tau}_k^{RSS} = \frac{1}{M} \sum_{m=1}^M \hat{\tau}_{km}^{SS}.$$

The repeated split sample estimator is asymptotically unbiased and Normal but, unlike the leave-one-out estimator, its large sample distribution does not coincide with the large sample distribution of the full-sample endogenous stratification estimator.

In the next section, we apply the estimators described above to the analysis of data from two well-known experimental studies: the National JTPA Study and the Tennessee Project STAR experiment.

⁵Proofs of these and other formal statements made in this paper are provided in Appendix 1.

⁶A separate issue in the estimation of τ_k is that first step estimation of β affects the large sample distribution of the estimator (see Appendix 1 for a derivation of the large sample distribution of $\hat{\tau}_k$ and $\hat{\tau}_k^{LOO}$). The contribution of the estimation of β to the variance of $\hat{\tau}_k$ has been ignored in empirical practice.

3. Evidence of Large Biases in Two Actual Applications

To demonstrate the performance of the estimators described in the previous section and the magnitude of their biases in realistic scenarios we use data from two randomized evaluations: the National JTPA Study, an evaluation of a vocational training program in the U.S., and the kindergarten cohort of the Tennessee Project STAR class-size experiment.

3.1. *The National JTPA Experiment*

We first examine data from the National JTPA Study. The National JTPA Study was a large experimental evaluation of a job training program commissioned by the U.S. Department of Labor in the late 1980's. The National JTPA Study data have been extensively analyzed in Orr et al. (1996), Bloom, et al. (1997), and many others. The National JTPA Study randomized access to vocational training to applicants in 16 service delivery areas, or SDAs, across the U.S. Randomized assignment was done after applicants were deemed eligible for the program and were recommended to one of three possible JTPA service strategies: on the job training/job search assistance, classroom training, and other services. Individuals in the treatment group were provided with access to JTPA services, and individuals in the control group were excluded from JTPA services for an 18-month period after randomization. We use data for the sample of male applicants recommended to the job training/job search assistance service strategy, and discarded three SDAs with few observations. Our sample consists of 1681 treated observations and 849 untreated observations, for a total of 2530 observations in 13 SDAs.⁷ In this example, w_i is an indicator of a randomized offer of JTPA services, y_i is nominal 30-month earnings in U.S. dollars after randomization, and \mathbf{x}_i includes age, age squared, marital status, previous earnings, indicators for having worked less than 13 weeks during the year previous to randomization, having a high-school degree, being African-American, and being Hispanic, as well as SDA indicators.

Table 1 reports estimates for the JTPA sample. The first row reports two treatment

⁷See Appendix 2 for detailed information on sample selection and estimation methods.

effect estimates. The “unadjusted” estimate is the difference in outcome means between treated and controls. The “adjusted” estimate is the coefficient on the treatment indicator in a linear regression of the outcome variable, y_i , on the treatment indicator, w_i , and the covariates, \mathbf{x}_i , listed above. The unadjusted estimate suggests a \$1516 effect on 30-month earnings. This estimate is significant at the 10 percent level. Regression adjustment reduces the point estimate to \$1207, which becomes marginally non-significant at the 10 percent level. The rest of Table 1 reports average treatment effects by predicted outcome group. The first set of estimates correspond to $\hat{\tau}_k$, the full-sample endogenous stratification estimator. This estimator produces a large and significant effect for the low predicted outcome group. The unadjusted estimate is \$2380 and significant at the 5 percent level. This represents a 12.6 percent effect on 30-month earnings, once we divide it by the average value of 30-month earnings among the experimental controls. It also represents an effect that is 57 percent higher than the corresponding unadjusted estimate for the average treatment effect in the first row of the table. The adjusted estimate is \$2012, similarly large, and significant at the 10 percent level. For the high predicted outcome group, the estimates are also large, but not statistically significant at conventional test levels. For the middle predicted outcome group, the estimates are negative, but of moderate magnitude and not statistically significant. All in all, the full-sample endogenous stratification estimates provide a much more favorable picture of JTPA effectiveness relative to the average treatment effects reported on the first row. Now, the bulk of the effect seems to be concentrated on the low predicted outcome group, precisely the one in most need of help, with more diffuse effects estimated for the middle and high predicted outcome groups.

The next two sets of estimates reported in Table 1 correspond to the leave-one-out estimator, $\hat{\tau}_k^{LOO}$, and the repeated split sample estimator $\hat{\tau}_k^{RSS}$, with number of repetitions, M , equal to 100. These two estimators, which avoid over-fitting bias arising from the estimation of β , produce results that are substantially different than those obtained with the full-sample endogenous stratification estimator, $\hat{\tau}_k$. Relative to the $\hat{\tau}_k$ estimates, the $\hat{\tau}_k^{LOO}$ and $\hat{\tau}_k^{RSS}$ estimates are substantially smaller for the low predicted outcome group,

and substantially larger for the high predicted outcome group. For the high predicted outcome group we obtain unadjusted estimates of \$3647 (leave-one-out) and \$3569 (split sample) both significant at the 5 percent level, and adjusted estimates of \$3118 (leave-one-out) and \$2943 (split sample) significant at the 10 percent and 5 percent levels respectively. The estimates for the low and middle predicted outcome groups are small in magnitude and not statistically significant. These results place the bulk of the treatment effect on the high predicted outcome group and do not provide substantial statistical evidence of beneficial effects for the low and middle predicted outcome groups.⁸ The comparison of estimates produced with the full sample endogenous stratification estimator and the leave-one-out and split sample estimators suggests that the over-fitting bias of the full sample endogenous stratification estimator is of substantial magnitude and dramatically changes the qualitative and quantitative interpretations of the results.

As a further check on the magnitude of endogenous stratification biases in the analysis of the National JTPA Study data, Table 1 reports a last set of treatment effects estimates, which are stratified using data on earnings before randomization. The National JTPA Study data include individual earnings during the 12 months before randomization. We use the sorting of the experimental subjects in terms of pre-randomization earnings to approximate how the experimental subjects would have been sorted in terms of earnings in the absence of the treatment. We construct the estimator $\hat{\tau}_k^{PREV}$ in the same way as $\hat{\tau}_k$ but using previous earnings, instead of predicted earnings, to divide the individuals into three groups of approximately equal size. Notice that, because previous earnings is a baseline characteristic, $\hat{\tau}_k^{PREV}$ is not affected by over-fitting bias. As shown on the bottom of Table 1, stratification on previous earning produces results similar to those obtained with $\hat{\tau}_k^{LOO}$ and $\hat{\tau}_k^{RSS}$: large and significant effects for the high predicted outcome group and smaller and non-significant effects for the middle and low predicted outcome groups.

⁸This is loosely consistent with the findings in Abadie, Angrist, and Imbens (2002) who report large JTPA effects at the upper tail of the distribution of earnings for male trainees, and no discernible effects at the middle or lower parts of the distribution.

3.2. *The Tennessee Project STAR Experiment*

Our second example uses data from the Tennessee Project STAR class-size study. In the Project STAR experiment, students in 79 schools were randomly assigned to small, regular-size, and regular-size classes with a teacher's aide. Krueger (1999) analyzes the STAR data set and provides detailed explanations of the STAR experiment. For our analysis, we use the 3764 students who entered the study in kindergarten and were assigned to small classes or to regular-size classes (without a teacher's aide). Our outcome variable is standardized end-of-the-year kindergarten math test scores.⁹ The covariates are indicators for African-American, female, eligibility for the free lunch program, and school attended. We discard observations with missing values in any of these variables.

Results for the STAR experiment data are reported in Table 2. The adjusted and unadjusted estimators of the average treatment effect on the first row of Table 2 show positive and significant effects. Using a simple difference in means, the effect of small classes is estimated as 0.1659 of the regular class standard deviation in math test scores, and 0.1892 of the same standard deviation when we use a regression-adjusted estimator. In both cases, the estimates are significant at the 5 percent level. For the low and middle predicted outcomes groups, the full-sample endogenous stratification estimator, $\hat{\tau}_k$, produces estimates that are positive and of a magnitude that is roughly double the magnitude of the average treatment effects estimates on the first row of the table. Counter-intuitively, however, the full sample endogenous stratification estimates for the high predicted outcome group are negative and significant. They seem to suggest that being assigned to small classes was detrimental for students who would have performed well in terms of math scores if all students had remained in regular-size classes. We deem this result counter-intuitive because it implies that reductions in the student/teacher ratio have detrimental effects on average for a large group of students. Notice that the magnitudes of the negative effects estimated for high predicted outcome group are substantial: smaller, but not far from the positive average treatment effects reported in the first row of the table. We will see that the large

⁹Standardized test scores are computed dividing raw test scores by the standard deviation of the distribution of the scores in regular-size classes.

and significant negative effect for the high predicted outcome group disappears when the leave-one-out or the repeated split sample procedures are used for estimation. Indeed, the leave-one-out and repeated sample split estimates on the two bottom rows of Table 2 suggest positive, significant, and large effects on the low and middle predicted outcome groups and effects of small magnitude and not reaching statistical significance at conventional test levels for the high predicted outcome group. Like for the JTPA, the qualitative and quantitative interpretations of the STAR experiment results change dramatically when the leave-one-out or the repeated split sample estimators are used instead of the full-sample endogenous stratification estimator.

In this section, we have used data from two well-known and influential experimental studies to investigate the magnitude of the distortion that over-fitting may induce on endogenous stratification estimators. In the next section, we use Monte Carlo simulations to assess the magnitude of the biases of the different estimators considered Section 2. To keep the exercise as realistic as possible, in two of our simulations we choose data generating processes that mimic the features of the JTPA and STAR data sets.

4. Simulation Evidence on the Behavior of Endogenous Stratification Estimators

This section reports simulation evidence on the finite sample behavior of endogenous stratification estimators. We run Monte Carlo simulations in three different settings. In the first two Monte Carlo simulations, we make use of the JTPA and STAR data sets to assess the magnitudes of biases and of other distortions to inference in realistic scenarios. The third and fourth Monte Carlo simulations use computer-generated data to investigate how the bias of endogenous stratification estimators changes when the sample size or the number of covariates changes.

In the JTPA-based simulation, we first use the JTPA control units to estimate a two-part model for the distribution of earnings conditional on the covariates of the adjusted estimates in Table 1. The two-part model consists of a Logit specification for the probability

of zero earnings and a Box-Cox model for positive earnings.¹⁰ In each Monte Carlo iteration we draw 2530 observations, that is, the same number of observations as in the JTPA sample, from the empirical distribution of the covariates in the JTPA sample. Next, we use the estimated two-part model to generate earnings data for each observation in the Monte Carlo sample. Then, we randomly assign 1681 observations to the treatment group and 849 observations to the control group, to match the numbers of treated and control units in the original JTPA sample. Finally, in each Monte Carlo iteration, we compute the full-sample, leave-one-out, and repeated split sample endogenous stratification estimates. We also compute the value of the unfeasible estimator, $\hat{\tau}_k^{UNF}$, obtained by stratification on the population regression function (which can be calculated from the estimated parameters of the two-part model by simulation). We conduct a total of 10000 Monte Carlo iterations.

Figure 1 reports the Monte Carlo distributions of the endogenous stratification estimators that divide the experimental sample into three categories of predicted earnings of roughly equal size (bottom third, middle third, and top third). To economize space this figure shows only the distribution of the unadjusted estimators.¹¹ Because assignment to the treatment and control groups is randomized in our simulation and because the process that generates earnings data is the same for treated and controls, it follows that the average effect of the treatment in the simulations is equal to zero unconditionally as well as conditional on the covariates. As a result, unbiased estimators should generate Monte Carlo distributions centered around zero. The first plot of Figure 1 shows the Monte Carlo distribution of the full-sample endogenous stratification estimator of average treatment effects conditional on predicted earnings group. The pattern of the distribution of the average treatment effect estimator for the bottom, middle, and top third predicted earnings groups matches the directions of the biases discussed in Section 2. That is, $\hat{\tau}_k$ is biased upwards for the low predicted earnings group and downwards for the high predicted earnings group. The remaining three plots of Figure 1 do not provide evidence of substantial biases for the

¹⁰Additional details about the simulation models can be found in Appendix 2.

¹¹Simulation results for unadjusted and adjusted estimators are very similar, as reflected in Tables 3 to 6 below.

leave-one-out, repeated split sample, or unfeasible estimators. These three estimators produce Monte Carlo distributions that are centered around zero for each predicted earnings category.

Table 3 reports biases, coverage rates for nominal 0.05 confidence intervals based on the Normal approximation, and root mean square error (root-MSE) values for endogenous stratification estimators in the JTPA-based Monte Carlo simulation. In addition to the estimators considered in Figure 1, we compute a single split sample estimator, $\hat{\tau}_k^{SSS}$, which is defined like the repeated split sample estimator but with $M = 1$. The full-sample endogenous stratification estimator is subject to substantial distortions for the low and high predicted earnings group. The magnitude of the bias in each these two groups is more than \$1000, which is substantial compared to the \$1516 and \$1207 unadjusted and adjusted average effect estimates in the JTPA data. As reflected in Figure 1, the bias is positive for the low predicted earnings group and negative for the high predicted earnings group. Biases are uniformly small for the leave-one-out, repeated split sample, and unfeasible estimators, but the leave-one-out estimator has higher biases than the repeated split sample and the unfeasible estimator. Similar results emerge for coverage rates and mean square errors. The full-sample endogenous stratification estimator produces substantially higher than nominal coverage rates and substantially higher root-MSE than the leave-one-out and repeated split sample estimators for the low and high predicted income categories. The repeated split sample estimator dominates in terms of root-MSE. The single split sample estimators produce small biases and close to nominal coverage rates, but has a root-MSE values consistently higher than the full-sample endogenous stratification estimator.

Figure 2 and Table 4 report simulation results for the STAR-based Monte Carlo simulation. For this simulation, the data generating process is based on a linear model with Normal errors. The model is estimated using data for STAR students in regular size classes. The results are qualitatively identical to those obtained in the JTPA-based simulation. The biases of $\hat{\tau}_k$ are around 0.05 and -0.05 for the low and medium predicted test score groups, respectively. These are sizable magnitudes, compared to the STAR effect estimates in Table

2. Also, like in the JTPA-based simulation, for the low and high predicted outcome groups coverage rates of the full-sample endogenous stratification estimator are heavily distorted and root-MSE values are larger than for the leave-one-out and the repeated split sample estimators. The repeated split sample estimator has the lowest root-MSE, and single sample splits produce root-MSE values that are larger than for any other estimator with the exception of the full-sample endogenous stratification estimator.

The third and fourth Monte Carlo simulations use computer generated data only. The purpose of these simulations is to demonstrate how the bias of endogenous stratification estimators changes with changes in the sample size and in the number of covariates. The data generating model for the third simulation is

$$y_i = 1 + \sum_{l=1}^{40} z_{li} + v_i$$

for $i = 1, \dots, N$, where the variables z_{li} have independent Standard Normal distributions, and the variable v_i has an independent Normal distribution with variance equal to 60. As a result, the unconditional variance of y_i is equal to 100. In each Monte Carlo simulation the sample is divided at random into two equally-sized treated and control groups. Predicted outcomes are computed using data for the control group to estimate

$$y_i = \alpha + \mathbf{x}'_{Ki} \boldsymbol{\beta}_K + u_{Ki}$$

by least squares, where \mathbf{x}_{Ki} is the $(K \times 1)$ -vector $(z_{1i}, \dots, z_{Ki})'$, for $K \leq 40$. That is, \mathbf{x}_{Ki} contains the values of the first K regressors in z_{1i}, \dots, z_{40i} . The data generating process implies that α is equal to one, $\boldsymbol{\beta}_K$ is a $(K \times 1)$ -vector of ones, $u_{Ki} = z_{K+1i} + \dots + z_{40i} + v_i$ if $K < 40$ and $u_{40i} = v_i$. We run Monte Carlo simulations for samples sizes $N = 200$, $N = 1000$, and $N = 5000$, and numbers of included regressors $K = 10$, $K = 20$, and $K = 40$.

The results are reported in Table 5. To economize space, we omit results on the single split sample estimator and report bias results only. Coverage rate and root-MSE results are available upon request. The magnitude of the biases in the Table 5 are easily understood

when compared to the standard deviation of the outcome, which is equal to 10. As expected, the bias of the full-sample endogenous stratification estimator is particularly severe when the sample size is small or when the number of included regressors is large, because in both cases significant over-fitting may occur. The increase in bias resulting from increasing the number of regressors is particularly severe when the sample size is small, $N = 200$. The biases of the leave-one-out, repeated split sample, and unfeasible estimators are negligible in most cases and consistently smaller than the bias of the full-sample endogenous stratification estimators, although the leave-one-out estimator tends to produce larger biases than the repeated split sample and unfeasible estimators.

The bias of the full-sample endogenous stratification estimator increases with K in spite of the fact that, as K increases, each additional included regressor has the same explanatory power as each of the regressors included in simulations with smaller K . Our final simulation studies a setting where each additional included regressor has lower explanatory power than the previously included ones. Consider:

$$y_i = 1 + \sum_{l=1}^{40} \rho^{l-1} z_{li} + \tilde{v}_i,$$

where the variables z_{li} have independent Standard Normal distributions as before, and the variable \tilde{v}_i has an independent Normal distribution with a variance such that the variance of y_i is equal to 100. Table 6 reports the biases of the endogenous stratification estimators across Monte Carlo simulations under the new data generating process (with $\rho = 0.80$). The biases of the endogenous stratification estimator are larger than in the previous simulation. Their magnitudes increase faster than in the previous simulation when the number of included covariates increases, and decrease slower than in the previous simulation when the number of observations increases. The biases of the leave-one-out, repeated split sample, and unfeasible estimators are smaller and less sensitive to changes in the number of included covariates and sample size.

Overall, among the estimators that address the over-fitting problem of full-sample endogenous stratification, the repeated split sample estimator out-performs leave-one-out in the simulations. Moreover, the leave-one-out estimator can behave erratically in settings

where the regressors take on only a few values and the variance of e_i is large.¹² The single split sample estimator has low bias and produces close-to-nominal coverage rates, but also large dispersion induced by the reduction in sample size. The increased variance of the single split sample estimator can make root-MSE of this estimator larger than the root-MSE of the full-sample endogenous stratification estimator (Table 3). All in all, the repeated split sample estimator displays the best performance in our simulations. It has low bias, accurate coverage rates, and out-performs alternative estimators in terms of root-MSE.

5. Conclusions

In this paper, we have argued that the increasingly popular practice of stratifying experimental units on the basis of a prediction of the outcome without treatment estimated using full sample data from control group leads to substantially biased estimates of treatment effects. We illustrate the magnitude of this bias using data from two well-known social experiments: the National JTPA Study and the Tennessee STAR Class Size Experiment. The full-sample endogenous stratification approach is most problematic in studies with small sample sizes and many regressors, where the predictor of the outcome without treatment may be severely over-fitted in the control sample. We demonstrate that alternative endogenous stratification estimators based on leave-one-out and repeated split sample techniques display substantially improved small sample behavior in our simulations relative to the full-sample endogenous stratification estimator.

¹²This is the case, for example, in the Tennessee STAR experiment if school indicators are excluded from the vector \mathbf{x}_i . In that case, \mathbf{x}_i only includes three indicator variables for race, gender, and eligibility for a free lunch program. As a result, $\mathbf{x}_i'\hat{\beta}$ takes on only eight different values. In this setting, over-fitting is not an issue and the full-sample endogenous stratification estimator produces small biases in simulations. However, the leave-one-out estimator generates large biases. The reason is that, in this setting, choosing c_1 and c_2 to be the quantiles 1/3 and 2/3 of the distribution of the predicted outcomes results in a large number of observations being located exactly at the boundaries of the values of $\mathbf{x}_i'\hat{\beta}$ that define the predicted outcome groups. To be concrete, consider the untreated observations with $\mathbf{x}_i'\hat{\beta} = c_1$. These observations are classified by the full-sample endogenous stratification estimator as members of the low predicted outcome group. However, it is easy to see that if $\mathbf{x}_i'\hat{\beta} = c_1$, then $\mathbf{x}_i'\hat{\beta}_{(-i)} > c_1$ if $y_i < c_1$ and $\mathbf{x}_i'\hat{\beta}_{(-i)} \leq c_1$ if $y_i \geq c_1$, which induces biases in the leave-one-out estimator.

Appendix 1: Proofs

Suppose that we have data from an experiment where a fraction p of experimental units are assignment to a treatment group and a fraction $1 - p$ to a control group, with $0 < p < 1$. Let N_1 be the number of units assigned to the treatment group and N_0 the number of units assigned to the control group, with $N = N_0 + N_1$. We will derive the limit distributions of endogenous stratification estimators as $N \rightarrow 0$. For each experimental unit, i , we observe the triple $\mathbf{u}_i = (y_i, w_i, \mathbf{x}_i)$, where w_i is a binary indicator that takes value one if observation i is in the treatment group, and value zero otherwise, y_i is the outcome of interest for observation i , and \mathbf{x}_i is a vector of baseline characteristics for observation i . We conduct our analysis assuming that the experimental units are sampled at random from some large population of interest, so the values of (y_i, \mathbf{x}_i) for the treated and the non-treated can be regarded as independent i.i.d. samples of sizes N_1 and N_0 from some distributions P_1 and P_0 , respectively. Probability statements about $\mathbf{u} = (y, w, \mathbf{x})$ are understood to refer to the distribution induced by first sampling w at random from a Bernoulli with parameter p and then sampling (y, \mathbf{x}) from P_1 with probability p and from P_0 with probability $1 - p$. Let

$$\boldsymbol{\beta} = (E[\mathbf{x}\mathbf{x}'|w=0])^{-1}E[\mathbf{x}y|w=0].$$

That is, $\mathbf{x}'\boldsymbol{\beta}$ is the linear least-squares predictor of $E[y|\mathbf{x}, w=0]$. Let c be a known constant such that $\Pr(\mathbf{x}'\boldsymbol{\beta} \leq c) > 0$. We aim to estimate:

$$\tau = E[y|w=1, \mathbf{x}'\boldsymbol{\beta} \leq c] - E[y|w=0, \mathbf{x}'\boldsymbol{\beta} \leq c].$$

This is the average effect of the treatment for individuals with $\mathbf{x}'\boldsymbol{\beta} \leq c$. Consider the full-sample endogenous stratification estimator:

$$\hat{\tau}(\hat{\boldsymbol{\beta}}) = \frac{\sum_{i=1}^N y_i I_{[w_i=1, \mathbf{x}'_i \hat{\boldsymbol{\beta}} \leq c]}}{\sum_{i=1}^N I_{[w_i=1, \mathbf{x}'_i \hat{\boldsymbol{\beta}} \leq c]}} - \frac{\sum_{i=1}^N y_i I_{[w_i=0, \mathbf{x}'_i \hat{\boldsymbol{\beta}} \leq c]}}{\sum_{i=1}^N I_{[w_i=0, \mathbf{x}'_i \hat{\boldsymbol{\beta}} \leq c]}},$$

where $\hat{\boldsymbol{\beta}}$ is a first-step estimator of the linear regression parameters that uses the untreated sample only:

$$\hat{\boldsymbol{\beta}} = \left(\sum_{i=1}^N \mathbf{x}_i (1 - w_i) \mathbf{x}'_i \right)^{-1} \sum_{i=1}^N \mathbf{x}_i (1 - w_i) y_i.$$

We will assume that \mathbf{x} has bounded support and that $E[y^2] < 0$. Under these assumptions, $\hat{\boldsymbol{\beta}} \xrightarrow{p} \boldsymbol{\beta}$, and

$$\begin{aligned} \sqrt{N}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}) &= (E[\mathbf{x}(1 - w)\mathbf{x}'])^{-1} \frac{1}{\sqrt{N}} \sum_{i=1}^N \mathbf{x}_i (1 - w_i) (y_i - \mathbf{x}'_i \boldsymbol{\beta}) + \mathcal{O}_p(1) \\ &= \mathcal{O}_p(1). \end{aligned}$$

For $j, l \in \{0, 1\}$ let

$$m_{jl}(\mathbf{u}, \mathbf{b}) = y^l I_{[w=j, \mathbf{x}' \mathbf{b} \leq c]}, \quad \mu_{jl}(\mathbf{b}) = E[m_{jl}(\mathbf{u}, \mathbf{b})].$$

Notice that

$$\hat{\tau}(\hat{\beta}) = \frac{\frac{1}{N} \sum_{i=1}^N m_{11}(\mathbf{u}_i, \hat{\beta}) - \frac{1}{N} \sum_{i=1}^N m_{01}(\mathbf{u}_i, \hat{\beta})}{\frac{1}{N} \sum_{i=1}^N m_{10}(\mathbf{u}_i, \hat{\beta}) - \frac{1}{N} \sum_{i=1}^N m_{00}(\mathbf{u}_i, \hat{\beta})},$$

and

$$\tau = \frac{\mu_{11}(\beta)}{\mu_{10}(\beta)} - \frac{\mu_{01}(\beta)}{\mu_{00}(\beta)}.$$

Because $\Pr(\mathbf{x}'\beta \leq c) > 0$ and $0 < p < 1$, then $\mu_{10}(\beta) > 0$ and $\mu_{00}(\beta) > 0$, so τ is well defined. Let

$$\mathbf{m}(\mathbf{u}, \mathbf{b}) = \begin{pmatrix} m_{11}(\mathbf{u}, \mathbf{b}) \\ m_{10}(\mathbf{u}, \mathbf{b}) \\ m_{01}(\mathbf{u}, \mathbf{b}) \\ m_{00}(\mathbf{u}, \mathbf{b}) \end{pmatrix}, \quad \hat{\mu}(\mathbf{b}) = \frac{1}{N} \sum_{i=1}^N \mathbf{m}(\mathbf{u}_i, \mathbf{b}), \quad \mu(\mathbf{b}) = E[\mathbf{m}(\mathbf{u}, \mathbf{b})].$$

We will first prove that $\sqrt{N}(\hat{\mu}(\hat{\beta}) - \mu(\hat{\beta})) - \sqrt{N}(\hat{\mu}(\beta) - \mu(\beta)) \xrightarrow{p} 0$. The following material about asymptotic equicontinuity follows Andrews (1994) closely. Let $\mathcal{M} = \{\mathbf{m}(\cdot, \mathbf{b}) : \mathbf{b} \in \Theta\}$ be a class of \mathbb{R}^s -valued functions. Here Θ is a pseudo-metric space with pseudo-metric ρ . Let

$$\mathbf{v}_N(\mathbf{b}) = \frac{1}{\sqrt{N}} \sum_{i=1}^N [\mathbf{m}(\mathbf{u}_i, \mathbf{b}) - \mu(\mathbf{b})].$$

We say that \mathbf{v}_N is stochastically equicontinuous if for all $\varepsilon > 0$ and $\eta > 0$ there exists $\delta > 0$ such that :

$$\limsup_{N \rightarrow \infty} \Pr \left(\sup_{\mathbf{b}_1, \mathbf{b}_2 \in \Theta, \rho(\mathbf{b}_1, \mathbf{b}_2) < \delta} \|\mathbf{v}_N(\mathbf{b}_1) - \mathbf{v}_N(\mathbf{b}_2)\| > \eta \right) < \varepsilon.$$

We aim to prove that $\mathbf{v}_N(\hat{\beta}) - \mathbf{v}_N(\beta) \xrightarrow{p} 0$. Suppose that the stochastic equicontinuity condition holds and that $\hat{\beta} - \beta \xrightarrow{p} 0$. Then, for any $\eta > 0$ and $\varepsilon > 0$ there exists $\delta > 0$ such that:

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \Pr \left(\|\mathbf{v}_N(\hat{\beta}) - \mathbf{v}_N(\beta)\| > \eta \right) \\ & \leq \limsup_{N \rightarrow \infty} \Pr \left(\|\mathbf{v}_N(\hat{\beta}) - \mathbf{v}_N(\beta)\| > \eta, \rho(\hat{\beta}, \beta) \leq \delta \right) + \limsup_{N \rightarrow \infty} \Pr \left(\rho(\hat{\beta}, \beta) > \delta \right) \\ & \leq \limsup_{N \rightarrow \infty} \Pr \left(\sup_{\mathbf{b}_1, \mathbf{b}_2 \in \Theta, \rho(\mathbf{b}_1, \mathbf{b}_2) < \delta} \|\mathbf{v}_N(\mathbf{b}_1) - \mathbf{v}_N(\mathbf{b}_2)\| > \eta \right) < \varepsilon. \end{aligned}$$

Consider now $\mathcal{M}_{11} = \{ywI_{[\mathbf{x}'\mathbf{b} \leq c]} : \mathbf{b} \in \Theta\}$, $\mathcal{M}_{01} = \{y(1-w)I_{[\mathbf{x}'\mathbf{b} \leq c]} : \mathbf{b} \in \Theta\}$, $\mathcal{M}_{10} = \{wI_{[\mathbf{x}'\mathbf{b} \leq c]} : \mathbf{b} \in \Theta\}$, and $\mathcal{M}_{00} = \{(1-w)I_{[\mathbf{x}'\mathbf{b} \leq c]} : \mathbf{b} \in \Theta\}$. It follows from Andrews (1994, Theorems 2 and 3) that \mathcal{M}_{11} and \mathcal{M}_{01} satisfy Pollard's entropy condition with envelope $\max\{1, y\}$, while \mathcal{M}_{10} and \mathcal{M}_{00} satisfy Pollard's entropy condition with envelope 1. By Andrews (1994, Theorem 1), if $E[|y|^{2+\delta}] < \infty$ for some $\delta > 0$, we obtain that each element of \mathbf{v}_N is stochastically equicontinuous with pseudo-metric ρ_{jl} given by:

$$\rho_{jl}(\mathbf{b}_1, \mathbf{b}_2) = \left(E \left[(m_{jl}(\mathbf{u}, \mathbf{b}_1) - m_{jl}(\mathbf{u}, \mathbf{b}_2))^2 \right] \right)^{1/2}.$$

As a result we obtain:

$$\begin{aligned}\sqrt{N}(\widehat{\boldsymbol{\mu}}(\widehat{\boldsymbol{\beta}}) - \boldsymbol{\mu}(\boldsymbol{\beta})) &= \sqrt{N}(\widehat{\boldsymbol{\mu}}(\boldsymbol{\beta}) - \boldsymbol{\mu}(\boldsymbol{\beta})) + \sqrt{N}(\boldsymbol{\mu}(\widehat{\boldsymbol{\beta}}) - \boldsymbol{\mu}(\boldsymbol{\beta})) \\ &\quad + \sqrt{N}((\widehat{\boldsymbol{\mu}}(\widehat{\boldsymbol{\beta}}) - \boldsymbol{\mu}(\widehat{\boldsymbol{\beta}})) - (\widehat{\boldsymbol{\mu}}(\boldsymbol{\beta}) - \boldsymbol{\mu}(\boldsymbol{\beta}))) \\ &= \sqrt{N}(\widehat{\boldsymbol{\mu}}(\boldsymbol{\beta}) - \boldsymbol{\mu}(\boldsymbol{\beta})) + \sqrt{N}(\boldsymbol{\mu}(\widehat{\boldsymbol{\beta}}) - \boldsymbol{\mu}(\boldsymbol{\beta})) + o_p(1).\end{aligned}$$

We will assume that $\boldsymbol{\mu}(\mathbf{b})$ is differentiable at $\boldsymbol{\beta}$ (Kim and Pollard 1990 section 5 provides high-level sufficient conditions). Then,

$$\boldsymbol{\mu}(\boldsymbol{\beta} + \mathbf{h}) - \boldsymbol{\mu}(\boldsymbol{\beta}) - \mathbf{r}(\boldsymbol{\beta})' \mathbf{h} = o(\|\mathbf{h}\|)$$

where

$$\mathbf{r}(\boldsymbol{\beta}) = \frac{\partial \boldsymbol{\mu}(\mathbf{b})}{\partial \mathbf{b}}(\boldsymbol{\beta}).$$

As a result (see, e.g., Lemma 2.12 in van der Vaart, 1998),

$$\boldsymbol{\mu}(\widehat{\boldsymbol{\beta}}) - \boldsymbol{\mu}(\boldsymbol{\beta}) - \mathbf{r}(\boldsymbol{\beta})'(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}) = o_p(\|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|).$$

Therefore, because $\sqrt{N}\|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}\| = O_p(1)$,

$$\sqrt{N}(\boldsymbol{\mu}(\widehat{\boldsymbol{\beta}}) - \boldsymbol{\mu}(\boldsymbol{\beta}) - \mathbf{r}(\boldsymbol{\beta})'(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta})) = o_p(\sqrt{N}\|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|) = o_p(1).$$

Now, it follows that

$$\begin{aligned}\sqrt{N}(\widehat{\boldsymbol{\mu}}(\widehat{\boldsymbol{\beta}}) - \boldsymbol{\mu}(\boldsymbol{\beta})) &= \sqrt{N}(\widehat{\boldsymbol{\mu}}(\boldsymbol{\beta}) - \boldsymbol{\mu}(\boldsymbol{\beta})) + \mathbf{r}(\boldsymbol{\beta})' \sqrt{N}(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}) + o_p(1) \\ &= \sqrt{N}(\boldsymbol{\mu}(\boldsymbol{\beta}) - \boldsymbol{\mu}(\boldsymbol{\beta})) + \mathbf{r}(\boldsymbol{\beta})'(E[\mathbf{x}(1-w)\mathbf{x}'])^{-1} \frac{1}{\sqrt{N}} \sum_{i=1}^N \mathbf{x}_i(1-w_i)(y_i - \mathbf{x}'_i \boldsymbol{\beta}) + o_p(1).\end{aligned}$$

Applying the multivariate CLT (see, e.g., Billingsley, 1995 Theorem 29.5), we obtain that $\sqrt{N}(\widehat{\boldsymbol{\mu}}(\widehat{\boldsymbol{\beta}}) - \boldsymbol{\mu}(\boldsymbol{\beta}))$ is asymptotically Normal with mean zero and variance equal to

$$\begin{aligned}\mathbf{V} = E\left[&(\mathbf{m}(\mathbf{u}, \boldsymbol{\beta}) - \boldsymbol{\mu}(\boldsymbol{\beta})) + \mathbf{r}(\boldsymbol{\beta})' \mathbf{Q}^{-1} \mathbf{x}(1-w)(y - \mathbf{x}\boldsymbol{\beta}) \\ &(\mathbf{m}(\mathbf{u}, \boldsymbol{\beta}) - \boldsymbol{\mu}(\boldsymbol{\beta})) + \mathbf{r}(\boldsymbol{\beta})' \mathbf{Q}^{-1} \mathbf{x}(1-w)(y - \mathbf{x}\boldsymbol{\beta})' \right],\end{aligned}$$

where $\mathbf{Q} = E[\mathbf{x}(1-w)\mathbf{x}']$. Consider $\mathbf{D} = (1/\mu_{10}(\boldsymbol{\beta}), -\mu_{11}(\boldsymbol{\beta})/\mu_{10}(\boldsymbol{\beta})^2, -1/\mu_{00}(\boldsymbol{\beta}), \mu_{01}(\boldsymbol{\beta})/\mu_{00}(\boldsymbol{\beta})^2)'$. Using the delta method, we obtain:

$$\sqrt{N}(\widehat{\tau}(\widehat{\boldsymbol{\beta}}) - \tau) \xrightarrow{d} N(\mathbf{0}, \mathbf{D}' \mathbf{V} \mathbf{D}).$$

Notice that estimation of the derivative vector \mathbf{r} can be accomplished using numerical methods (see, e.g., Newey and McFadden 1994, Theorem 7.4).

The following intermediate lemma will be useful to derive the properties of the leave-one-out estimator.

LEMMA A.1: Let x_N be a sequence of random variables, a_N a sequence of real numbers, and I_{A_N} be the indicator function for the event A_N . Suppose that $a_N \Pr(I_{A_N} = 0) \rightarrow 0$ and $a_N \Pr(I_{A_N} x_N > \varepsilon) \rightarrow 0$ for some $\varepsilon > 0$. Then, $a_N \Pr(x_N > \varepsilon) \rightarrow 0$.

Proof:

$$\begin{aligned} a_N \Pr(x_N > \varepsilon) &= a_N \Pr(x_N > \varepsilon, I_{A_N} = 1) + a_N \Pr(x_N > \varepsilon, I_{A_N} = 0) \\ &= a_N \Pr(I_{A_N} x_N > \varepsilon, I_{A_N} = 1) + a_N \Pr(x_N > \varepsilon, I_{A_N} = 0) \\ &\leq a_N \Pr(I_{A_N} x_N > \varepsilon) + a_N \Pr(I_{A_N} = 0) \rightarrow 0. \end{aligned}$$

□

Consider now the leave-one-out estimator:

$$\begin{aligned} \hat{\tau}^{LOO} &= \frac{\frac{1}{N} \sum_{i=1}^N y_i I_{[w_i=1, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]} - \frac{1}{N} \sum_{i=1}^N y_i I_{[w_i=0, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]}}{\frac{1}{N} \sum_{i=1}^N I_{[w_i=1, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]} - \frac{1}{N} \sum_{i=1}^N I_{[w_i=0, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]}} \\ &= \frac{\frac{1}{N} \sum_{i=1}^N y_i I_{[w_i=1, \mathbf{x}'_i \hat{\beta} \leq c]} - \frac{1}{N} \sum_{i=1}^N y_i I_{[w_i=0, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]}}{\frac{1}{N} \sum_{i=1}^N I_{[w_i=1, \mathbf{x}'_i \hat{\beta} \leq c]} - \frac{1}{N} \sum_{i=1}^N I_{[w_i=0, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]}}. \end{aligned}$$

Therefore,

$$\begin{aligned} \sqrt{N}(\hat{\tau}(\hat{\beta}) - \hat{\tau}^{LOO}) &= \sqrt{N} \left(\frac{\frac{1}{N} \sum_{i=1}^N y_i I_{[w_i=0, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]} - \frac{1}{N} \sum_{i=1}^N y_i I_{[w_i=0, \mathbf{x}'_i \hat{\beta} \leq c]}}{\frac{1}{N} \sum_{i=1}^N I_{[w_i=0, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]} - \frac{1}{N} \sum_{i=1}^N I_{[w_i=0, \mathbf{x}'_i \hat{\beta} \leq c]}} \right) \\ &= \frac{\frac{1}{\sqrt{N}} \sum_{i=1}^N y_i (I_{[w_i=0, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]} - I_{[w_i=0, \mathbf{x}'_i \hat{\beta} \leq c]})}{\frac{1}{N} \sum_{i=1}^N I_{[w_i=0, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]}} \\ &\quad - \frac{\frac{1}{\sqrt{N}} \sum_{i=1}^N (I_{[w_i=0, \mathbf{x}'_i \hat{\beta}_{(-i)} \leq c]} - I_{[w_i=0, \mathbf{x}'_i \hat{\beta} \leq c]})}{\frac{1}{N} \sum_{i=1}^N I_{[w_i=0, \mathbf{x}'_i \hat{\beta} \leq c]}} \frac{1}{N} \sum_{i=1}^N y_i I_{[w_i=0, \mathbf{x}'_i \hat{\beta} \leq c]}. \end{aligned}$$

We will show that $\sqrt{N}(\hat{\tau}(\hat{\beta}) - \hat{\tau}^{LOO}) \xrightarrow{p} 0$. Suppose that the r -th moment of $|y|$ exists, where $r > 1$ (later we will strengthen this requirement to $r > 2$ and eventually to $r > 3$). Then, by

Holder's Inequality:

$$\frac{1}{\sqrt{N}} \sum_{i=1}^N |y_i| \left| I_{[\mathbf{x}_i' \widehat{\boldsymbol{\beta}} \leq c]} - I_{[\mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \leq c]} \right| \leq N^{1/2} \left(\frac{1}{N} \sum_{i=1}^N |y_i|^r \right)^{1/r} \left(\frac{1}{N} \sum_{i=1}^N \left| I_{[\mathbf{x}_i' \widehat{\boldsymbol{\beta}} \leq c]} - I_{[\mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \leq c]} \right| \right)^{(r-1)/r}.$$

The first sample average on the right hand side of last equation is bounded in probability. Now we need to show that the second sample average on the right hand side of last equation goes to zero fast enough to beat $N^{1/2}$ after taking the $(r-1)/r$ power. By exchangeability of $(\mathbf{x}_i' \widehat{\boldsymbol{\beta}}, \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)})$ (for $i = 1, \dots, N$):

$$\begin{aligned} E \left[\frac{1}{N} \sum_{i=1}^N \left| I_{[\mathbf{x}_i' \widehat{\boldsymbol{\beta}} \leq c]} - I_{[\mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \leq c]} \right| \right] &= E \left[\frac{1}{N} \sum_{i=1}^N I_{[\mathbf{x}_i' \widehat{\boldsymbol{\beta}} \leq c < \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \cup \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \leq c < \mathbf{x}_i' \widehat{\boldsymbol{\beta}}]} \right] \\ &= \Pr \left(\mathbf{x}_i' \widehat{\boldsymbol{\beta}} \leq c < \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \cup \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \leq c < \mathbf{x}_i' \widehat{\boldsymbol{\beta}} \right). \end{aligned}$$

Therefore, by Markov's inequality, it is enough to show that:

$$N^{\frac{r}{2(r-1)}} \Pr \left(\mathbf{x}_i' \widehat{\boldsymbol{\beta}} \leq c < \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \cup \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \leq c < \mathbf{x}_i' \widehat{\boldsymbol{\beta}} \right) \xrightarrow{p} 0.$$

Let $\zeta_N = N^\alpha$, where $\alpha > r/(2(r-1))$. Notice that,

$$\begin{aligned} &\Pr \left(\mathbf{x}_i' \widehat{\boldsymbol{\beta}} \leq c < \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \cup \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \leq c < \mathbf{x}_i' \widehat{\boldsymbol{\beta}} \right) \\ &\leq \Pr \left(\mathbf{x}_i' \widehat{\boldsymbol{\beta}} \leq c < \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \cup \mathbf{x}_i' \widehat{\boldsymbol{\beta}}_{(-i)} \leq c < \mathbf{x}_i' \widehat{\boldsymbol{\beta}}, |\mathbf{x}_i' \widehat{\boldsymbol{\beta}} - c| > 1/\zeta_N \right) \\ &\quad + \Pr \left(|\mathbf{x}_i' \widehat{\boldsymbol{\beta}} - c| \leq 1/\zeta_N \right) \\ &\leq \Pr \left(|\mathbf{x}_i' (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_{(-i)})| > 1/\zeta_N \right) + \Pr \left(|\mathbf{x}_i' \widehat{\boldsymbol{\beta}} - c| \leq 1/\zeta_N \right). \end{aligned}$$

Suppose that there exists $\epsilon > 0$ such that for $\mathbf{b} \in B(\boldsymbol{\beta}, \epsilon)$, the distribution of $\mathbf{x}' \mathbf{b}$ is absolutely continuous with density bounded (uniformly) by a constant C . Assume that $r > 2$. Consider a sequence $\epsilon_N = N^{-\gamma}$, where $0 < \gamma < (r-2)/4(r-1)$. Then, for large enough N (so $\epsilon_N < \epsilon$)

$$\begin{aligned} \Pr \left(|\mathbf{x}_i' \widehat{\boldsymbol{\beta}} - c| \leq 1/\zeta_N \right) &\leq \Pr \left(|\mathbf{x}_i' \widehat{\boldsymbol{\beta}} - c| \leq 1/\zeta_N, \widehat{\boldsymbol{\beta}} \in B(\boldsymbol{\beta}, \epsilon_N) \right) + \Pr \left(\widehat{\boldsymbol{\beta}} \notin B(\boldsymbol{\beta}, \epsilon_N) \right) \\ &\leq \sup_{\mathbf{b} \in B(\boldsymbol{\beta}, \epsilon_N)} \Pr \left(|\mathbf{x}_i' \mathbf{b} - c| \leq 1/\zeta_N \right) + \Pr \left(\|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}\| > \epsilon_N \right) \\ &\leq \frac{2C}{\zeta_N} + \Pr \left(N^\gamma \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}\| > 1 \right). \end{aligned}$$

The first term on the right hand side of last equation multiplied by $N^{r/(2(r-1))}$ converges to zero because $N^{r/(2(r-1))}/\zeta_N \rightarrow 0$. Assume $\lambda_{\min}(\mathbf{Q}) > 0$. Because $\|\mathbf{x}_i\|$ is bounded by some constant, M , it follows that

$$\lambda_{\max} \left(\frac{1}{N} \mathbf{x}_i (1 - w_i) \mathbf{x}_i' \right) = \max_{\|\mathbf{v}\|=1} \mathbf{v}' \mathbf{x}_i (1 - w_i) \mathbf{x}_i' \mathbf{v} / N \leq \frac{M^2}{N}.$$

Let

$$\mathbf{Q}_N = \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i (1 - w_i) \mathbf{x}_i'.$$

Now, Corollary 5.2 in Tropp (2012) implies

$$\Pr(\lambda_{\min}(\mathbf{Q}_N) \leq t\lambda_{\min}(\mathbf{Q})) \leq K e^{-\frac{N(1-t)^2\lambda_{\min}(\mathbf{Q})}{2M^2}}, \quad (\text{A.1})$$

where K is the length of \mathbf{x}_i and $t \in [0, 1]$. Define the event

$$A_N = \{\lambda_{\min}(\mathbf{Q}_N) \geq C_\lambda\},$$

for some $0 < C_\lambda < \lambda_{\min}(\mathbf{Q})$, and let I_{A_N} be the indicator function for the event A_N . The concentration inequality in (A.1) implies

$$N^{\frac{r}{2(r-1)}} \Pr(I_{A_N} = 0) \rightarrow 0. \quad (\text{A.2})$$

Let $e_i = y_i - \mathbf{x}'_i \boldsymbol{\beta}$. Notice that,

$$\begin{aligned} I_{A_N} N \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|^2 &= I_{A_N} N (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta})' (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}) \\ &= I_{A_N} \left(\frac{1}{\sqrt{N}} \sum_{i=1}^N e_i (1 - w_i) \mathbf{x}'_i \right) \left(\frac{1}{N} \sum_{i=1}^N \mathbf{x}_i (1 - w_i) \mathbf{x}'_i \right)^{-2} \left(\frac{1}{\sqrt{N}} \sum_{i=1}^N \mathbf{x}_i (1 - w_i) e_i \right) \\ &\leq I_{A_N} \lambda_{\min}^{-2}(\mathbf{Q}_N) \left(\frac{1}{\sqrt{N}} \sum_{i=1}^N e_i (1 - w_i) \mathbf{x}'_i \right) \left(\frac{1}{\sqrt{N}} \sum_{i=1}^N \mathbf{x}_i (1 - w_i) e_i \right) \\ &\leq C_\lambda^{-2} \left(\frac{1}{\sqrt{N}} \sum_{i=1}^N e_i (1 - w_i) \mathbf{x}'_i \right) \left(\frac{1}{\sqrt{N}} \sum_{i=1}^N \mathbf{x}_i (1 - w_i) e_i \right). \end{aligned}$$

Given that \mathbf{x} is bounded, $E[e_i^2] < \infty$ (which follows from $r > 2$), and given that $E[e_i(1 - w_i) \mathbf{x}'_i \mathbf{x}_j (1 - w_j) e_j] = E[e_i(1 - w_i) \mathbf{x}'_i] E[\mathbf{x}_j (1 - w_j) e_j] = 0$ for any $1 \leq i < j \leq N$, we obtain

$$\limsup_{N \rightarrow \infty} E[I_{A_N} N \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|^2] < \infty. \quad (\text{A.3})$$

By Markov's inequality, equation (A.3), and because $\gamma < (r-2)/4(r-1)$

$$N^{\frac{r}{2(r-1)}} \Pr(I_{A_N} N^\gamma \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}\| > 1) \leq N^{2\gamma-1+\frac{r}{2(r-1)}} E[I_{A_N} N \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|^2] \rightarrow 0. \quad (\text{A.4})$$

From equations (A.2), (A.4), and Lemma A.1, it follows that

$$N^{\frac{r}{2(r-1)}} \Pr(N^\gamma \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}\| > 1) \rightarrow 0.$$

Therefore,

$$N^{\frac{r}{2(r-1)}} \Pr(|\mathbf{x}'_i \widehat{\boldsymbol{\beta}} - c| \leq 1/\zeta_N) \rightarrow 0.$$

Notice that (see Hansen, 2012 sections 4.12 and 4.13)

$$\begin{aligned} |\mathbf{x}'_i (\widehat{\boldsymbol{\beta}} - \widehat{\boldsymbol{\beta}}_{(-i)})| &= \frac{h_{Ni}}{1 - h_{Ni}} |y_i - \mathbf{x}'_i \widehat{\boldsymbol{\beta}}| \\ &\leq \left(\frac{\max_{i=1,\dots,N} h_{Ni}}{1 - \max_{i=1,\dots,N} h_{Ni}} \right) |y_i - \mathbf{x}'_i \widehat{\boldsymbol{\beta}}|, \end{aligned}$$

where the leverage values h_{Ni} are

$$h_{Ni} = (1 - w_i) \mathbf{x}_i' \left(\sum_{i=1}^N \mathbf{x}_i (1 - w_i) \mathbf{x}_i' \right)^{-1} \mathbf{x}_i. \quad (\text{A.5})$$

Therefore,

$$\max_{1 \leq i \leq N} h_{Ni} \leq \lambda_{\min}^{-1}(\mathbf{Q}_N) \frac{M^2}{N}.$$

Notice also that,

$$\begin{aligned} & I_{A_N} \|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|^r \\ & \leq I_{A_N} \left(\left(\frac{1}{N} \sum_{i=1}^N (1 - w_i) e_i \mathbf{x}_i' \right) \left(\frac{1}{N} \sum_{i=1}^N \mathbf{x}_i (1 - w_i) \mathbf{x}_i' \right)^{-2} \left(\frac{1}{N} \sum_{i=1}^N \mathbf{x}_i e_i (1 - w_i) \right) \right)^{r/2} \\ & \leq I_{A_N} \lambda_{\min}^{-r}(\mathbf{Q}_N) \left(\left(\frac{1}{N} \sum_{i=1}^N (1 - w_i) e_i \mathbf{x}_i' \right) \left(\frac{1}{N} \sum_{i=1}^N \mathbf{x}_i e_i (1 - w_i) \right) \right)^{r/2} \\ & \leq C_{\lambda}^{-r} \left(\left(\frac{1}{N} \sum_{i=1}^N (1 - w_i) e_i \mathbf{x}_i' \right) \left(\frac{1}{N} \sum_{i=1}^N \mathbf{x}_i e_i (1 - w_i) \right) \right)^{r/2} \\ & = C_{\lambda}^{-r} M^r \left(\frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N |e_i e_j| \right)^{r/2}. \end{aligned}$$

If $r \geq 2$, by Minkowski's and Cauchy-Schwartz's inequalities:

$$\begin{aligned} & \left(E \left[\left(\frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N |e_i e_j| \right)^{r/2} \right] \right)^{2/r} \leq \frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \left(E [|e_i e_j|^{r/2}] \right)^{2/r} \\ & \leq \frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N (E [|e_1|^r])^{2/r}. \end{aligned}$$

Because $E[|e_i|^r] = E[|y_i - \mathbf{x}_i' \boldsymbol{\beta}|^r]$, $E[|y_i|^r] < \infty$ and $\|\mathbf{x}_i\|$ is bounded, we obtain

$$\limsup_{N \rightarrow \infty} E[I_{A_N} \|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|^r] < \infty,$$

which, by Minkowski's inequality, implies

$$\limsup_{N \rightarrow \infty} E[I_{A_N} |y_i - \mathbf{x}_i' \hat{\boldsymbol{\beta}}|^r] < \infty. \quad (\text{A.6})$$

By Markov's inequality:

$$N^{\frac{r}{2(r-1)}} \Pr \left(I_{A_N} |\mathbf{x}_i' (\hat{\boldsymbol{\beta}} - \hat{\boldsymbol{\beta}}_{(-i)})| > 1/\zeta_N \right) \leq N^{\frac{r}{2(r-1)} + r\alpha} E \left[I_{A_N} |\mathbf{x}_i' (\hat{\boldsymbol{\beta}} - \hat{\boldsymbol{\beta}}_{(-i)})|^r \right].$$

The condition $\alpha < (2r - 3)/(2(r - 1))$ implies $r/(2(r - 1)) + r\alpha < r$. So, under that condition, it is left to be proven that for any $\vartheta > 0$

$$N^{r-\vartheta} E \left[I_{A_N} |\mathbf{x}'_i(\hat{\boldsymbol{\beta}} - \hat{\boldsymbol{\beta}}_{(-i)})|^r \right] \rightarrow 0. \quad (\text{A.7})$$

Consider N large enough so that there is a positive constant C_d , such that $1/(1 - C_\lambda^{-1} M^2/N) < C_d$. Then,

$$I_{A_N} |\mathbf{x}'_i(\hat{\boldsymbol{\beta}} - \hat{\boldsymbol{\beta}}_{(-i)})| \leq C_d C_\lambda^{-1} \frac{M^2}{N} |y_i - \mathbf{x}'_i \hat{\boldsymbol{\beta}}|.$$

This result, along with equation (A.6) implies:

$$\limsup_{N \rightarrow \infty} E[I_{A_N} N^r |\mathbf{x}'_i(\hat{\boldsymbol{\beta}} - \hat{\boldsymbol{\beta}}_{(-i)})|^r] < \infty,$$

so equation (A.7) holds. Notice that for the condition

$$\frac{r}{2(r - 1)} < \alpha < \frac{2r - 3}{2(r - 1)}$$

to hold we need $r > 3$.

Similar derivations as for $\hat{\tau}_k$ can be used to find the large sample distribution of $\hat{\tau}_k^{RSS}$. We omit the details here.

Appendix 2: Estimation and Simulation Details

Leave-one-out predictions can be efficiently calculated using:

$$\mathbf{x}'_i \hat{\boldsymbol{\beta}}_{(-i)} = \mathbf{x}'_i \hat{\boldsymbol{\beta}} - \frac{h_{Ni}}{1 - h_{Ni}} (y_i - \mathbf{x}'_i \hat{\boldsymbol{\beta}}),$$

where h_{Ni} is the leverage value defined in equation (A.5). Let \hat{y}_i be generic notation for a predicted outcome without treatment. The prediction, \hat{y}_i , may come from full-sample, leave-one-out, or split sample endogenous stratification, stratification on previous earnings in the JTPA example of Section 3, or stratification on the true regression value in the simulations of Section 4. We group observations on the basis of predicted outcomes, \hat{y}_i , in the following way. First we sort the observations based on predicted outcomes: $\hat{y}_{(1)} \leq \hat{y}_{(2)} \leq \dots \leq \hat{y}_{(N)}$. We then define t_1 and t_2 as $N/3$ and $2N/3$ rounded to the nearest integer, respectively. We classify unit i in the low, medium, and high predicted outcome groups if $\hat{y}_i \leq \hat{y}_{(t_1)}$, $\hat{y}_{(t_1)} < \hat{y}_i \leq \hat{y}_{(t_2)}$, and $\hat{y}_{(t_2)} < \hat{y}_i$, respectively.

For the repeated split sample estimator estimation is as follows. For the JTPA data we randomly select 425 control observations and use them to estimate $\boldsymbol{\beta}$. We use the remaining 424 observations and all the treated JTPA units for the second step estimation of τ_k . For the STAR data, we use 1009 untreated observations to estimate $\boldsymbol{\beta}$ and the remaining 1008 and all the treated observations in the second step. We average the split sample estimators over 100 repetitions to obtain $\hat{\tau}_k^{RSS}$.

As explained in Section 3, the JTPA sample consists of male applicants assigned to on the job training/job search assistance. We discard three of the sixteen SDAs, Jersey City (21 observations), Butte (15 observations), and Oakland (5 observations) because of small sample sizes. The

STAR sample consists of use 3764 students who entered the study in kindergarten, were assigned to small classes or to regular-size classes without a teacher's aide, and for whom there is complete information on all the variables used in our analysis.

Standard errors in Tables 1 and 2 are calculated using the nonparametric bootstrap (conditioning on the number of treated and untreated observations in the original samples).

In the JTPA-based simulation we first estimate a Logit model, $p(\mathbf{x}, \boldsymbol{\gamma}) = e^{\mathbf{x}'\boldsymbol{\gamma}}/(1 + e^{\mathbf{x}'\boldsymbol{\gamma}})$, for the probability of employment, measured as positive labor market earnings, using the sample of experimental controls. Next, using only the experimental control with positive earnings, we estimate a Box-Cox regression model

$$\frac{y^\lambda - 1}{\lambda} = \mathbf{x}'\boldsymbol{\theta} + \sigma u,$$

where u has a Standard Normal distribution. We will use $\boldsymbol{\gamma}^*$ to refer to the estimate of $\boldsymbol{\gamma}$, and analogous notation for the estimated parameters of the Box-Cox model. We create each simulated data set in the following manner. We first resample 2530 observations from the empirical distribution of \mathbf{x} among all the JTPA sample units. We assign zero earnings with probability $1 - p(\mathbf{x}, \boldsymbol{\gamma}^*)$. With probability $p(\mathbf{x}, \boldsymbol{\gamma}^*)$ we assign earnings using

$$y = (\max\{1 + \lambda^*(\mathbf{x}'\boldsymbol{\theta}^* + \sigma^*u), 0\})^{1/\lambda^*},$$

where u has a Standard Normal distribution. We randomly label 1681 observations as treated and 849 as untreated. As a result, all treatment effects are equal to zero by construction. The coefficients of the regression function of y on \mathbf{x} under this data generating process, which are needed to compute $\hat{\tau}_k^{UNF}$, are calculated by simulation.

For the STAR-based simulations, we estimate the linear model

$$y = \mathbf{x}'\boldsymbol{\beta} + \sigma u,$$

where u has a Standard Normal distribution, using the sample of experimental controls. We use least squares to estimate $\boldsymbol{\beta}$ and the variance of the regression residuals corrected for degrees of freedom to estimate σ . To construct each simulated sample, we first randomly resample 3764 observations from the empirical distribution of \mathbf{x} in the STAR sample. We simulate math scores using

$$y = \mathbf{x}'\boldsymbol{\beta}^* + \sigma^*u,$$

where u has a Standard Normal distribution, and $\boldsymbol{\beta}^*$ and σ^* are the estimates of $\boldsymbol{\beta}$ and σ .

Section 4 contains detailed information on data generating processes for the simulations of Tables 5 and 6.

Bias, coverage rates, and root-MSE are calculated as follows. Because the simulations impose zero treatments effects, the bias and the MSE are calculated as the mean of the estimates and the mean of the square of the estimates, respectively, across all simulation repetitions. We calculate t -ratios dividing the estimates from each simulation repetition by the standard deviation of the estimates across repetitions. Coverage rates are the frequencies of the t -ratios falling outside the $[-1.96, 1.96]$ interval across repetitions. Root-MSE is the square-root of the MSE.

References

Abadie, Alberto, Joshua D. Angrist and Guido W. Imbens. 2002. "Instrumental Variables Estimates of the Effect of Subsidized Training on the Quantiles of Trainee Earnings." *Econometrica* 70(1): 91-117.

Andrews, Donald W. K. 1994. "Empirical Process Methods in Econometrics." In R.F. Engle and D.L. McFadden (Eds.), *Handbook of Econometrics*, Vol. IV. Amsterdam: Elsevier Science.

Angrist, Joshua D. and Victor Lavy. 2009. "The Effects of High Stakes High School Achievement Awards: Evidence from a Randomized Trial." *American Economic Review* 99(4): 1384-1414.

Angrist, Joshua D. and Jorn-Steffen Pischke. 2010. "The Credibility Revolution in Empirical Economics: How Better Research Design Is Taking the Con out of Econometrics." *Journal of Economic Perspectives* 24(2): 3-30.

Billingsley, Patrick. 1995. *Probability and Measure*, third edition. New York: Wiley.

Bloom, Howard S., Larry L. Orr, Stephen H. Bell, George Cave, Fred Doolittle, Winston Lin and Johannes M. Bos. 1997. "The Benefits and Costs of JTPA Title II-A Programs." *The Journal of Human Resources*, 32(3): 549-576.

Deming, David J. 2011. "Better Schools, Less Crime?" *Quarterly Journal of Economics* 126(4): 2063-2115.

Dynarski, Susan, Joshua M. Hyman and Diane Schanzenbach. 2011. "Experimental Evidence on the Effect of Childhood Investments on Post-secondary Attainment and Degree Completion." National Bureau of Economic Research Working Paper No. 17533.

Dynarski, Susan, Joshua M. Hyman and Diane Schanzenbach. 2013. "Experimental Evidence on the Effect of Childhood Investments on Postsecondary Attainment and Degree Completion." *Journal of Policy Analysis and Management* 32(4): 692-717.

Giné, Xavier, Jessica Goldberg and Dean Yang. 2012. "Credit Market Consequences of Improved Personal Identification." *American Economic Review* 102(6): 2923-54.

Goldrick-Rab, Sara, Douglas N. Harris, James Benson and Robert Kelchen. 2011. "Conditional cash transfers and college persistence: Evidence from a randomized need-based grant program." Institute for Research on Poverty Discussion Paper No. 1393-11.

Goldrick-Rab, Sara, Douglas N. Harris, James Benson and Robert Kelchen. 2012. "Need-based financial aid and college persistence: Experimental evidence from Wisconsin." Available at SSRN 1887826.

Hansen, Ben B. 2008. "The Prognostic Analogue of the Propensity Score." *Biometrika* 95(2): 481-488.

Hansen, Bruce E. 2012. *Econometrics*. Available at: <http://www.ssc.wisc.edu/~bhansen/>

Harris, Douglas N. and Sara Goldrick-Rab. 2012. "Improving the Productivity of Education Experiments: Lessons from a Randomized Study of Need-Based Financial Aid." *Education Finance and Policy* 7(2): 143-169.

Harvill, Eleanor L., Laura R. Peck and Stephen H. Bell. 2013. "On Overfitting in Analysis of Symmetrically Predicted Endogenous Subgroups From Randomized Experimental Samples: Part Three of a Method Note in Three Parts." *American Journal of Evaluation* 34(4): 545-566.

Heckman, James J., Jeffrey Smith and Nancy Clements. 1997. "Making the Most Out of Programme Evaluations and Social Experiments: Accounting for Heterogeneity in Programme Impacts." *Review of Economic Studies* 64(4): 487-535.

Hemelt, Steven W., Kimberly B. Roth and William W. Eaton. 2012. "Childhood Educational Interventions: Experimental Evidence on Post-secondary Outcomes." Paper presented at the Association for Education Finance and Policy Conference, Boston, MA.

Kent, David M. and Rodney A. Hayward. 2007. "Limitations of Applying Summary Results of Clinical Trials to Individual Patients: The Need for Risk Stratification." *Journal of the American Medical Association* 298(10): 1209-12.

Kim, Jeankyung and David Pollard. 1990. "Cube Root Asymptotics." *The Annals of Statistics* 18(1): 191-219.

Krueger, Alan B. 1999. "Experimental Estimates of Education Production Functions." *Quarterly Journal of Economics* 114(2): 497-532.

Murnane, Richard J. and John B. Willett. 2011. *Methods Matter: Improving Causal Inference in Educational and Social Science Research*. Oxford: Oxford University Press.

Newey, Whitney K. and Daniel McFadden. 1994. "Large Sample Estimation and Hypothesis Testing." In: Engle, R.F., McFadden, D.L. (Eds.), *Handbook of Econometrics, Vol. IV*. Amsterdam: Elsevier Science.

Orr, Larry L., Howard S. Bloom, Stephen H. Bell, Fred Doolittle, Winston Lin and George Cave. 1996. *Does Training for the Disadvantaged Work?* Washington, DC: The Urban Institute.

Pane, John F., Beth Ann Griffin, Daniel F. McCaffrey and Rita Karam. 2013. "Effectiveness of Cognitive Tutor Algebra I at Scale." RAND Education Working Paper.

Peck, Laura R. 2003. "Subgroup Analysis in Social Experiments: Measuring Program Impacts Based on Post-Treatment Choice." *American Journal of Evaluation* 24(2): 157-187.

Rodriguez-Planas, Nuria. 2012. "Schools and Drugs: Closing the Gap. Evidence from a Randomized Trial in the US." IZA Discussion Paper No. 6770.

Sanbonmatsu, Lisa, Jeffrey R. Kling, Greg J. Duncan and Jeanne Brooks-Gunn. 2006. "Neighborhoods and Academic Achievement: Results from the Moving to Opportunity Experiment." *Journal of Human Resources* 41(4): 649-691.

Tropp, Joel A. 2012. "User-Friendly Tail Bounds for Sums of Random Matrices." *Foundations of Computational Mathematics* 12: 389-434.

van der Vaart, Aad W. 1998. *Asymptotic Statistics*. New York: Cambridge University Press.




Figure 1
Distributions of the Estimators in the JTPA Simulation




Figure 2
Distributions of the Estimators in the STAR Simulation

Table 1
JTPA Estimation Results

Panel A: Average treatment effect

	unadjusted	adjusted
$\hat{\tau}$	1516.49* (807.27)	1207.22 (763.54)

Panel B: Average treatment effect by predicted outcome group

	unadjusted			adjusted		
	low	medium	high	low	medium	high
$\hat{\tau}_k$	2379.65** (1151.07)	-719.38 (1474.81)	2397.26 (1672.62)	2011.70* (1150.68)	-554.65 (1482.32)	1769.03 (1639.06)
$\hat{\tau}_k^{LOO}$	573.74 (1201.33)	35.31 (1509.30)	3646.53** (1727.08)	173.45 (1213.25)	172.28 (1513.70)	3118.17* (1679.62)
$\hat{\tau}_k^{RSS}$	788.75 (1027.47)	254.25 (1092.85)	3569.41** (1496.73)	412.01 (1042.17)	181.81 (1087.51)	2942.69** (1454.16)
$\hat{\tau}_k^{PREV}$	1278.88 (1221.96)	-67.95 (1284.77)	3972.21** (1497.47)	822.05 (1235.13)	-150.89 (1274.45)	3146.85** (1430.37)

Note: The JTPA sample includes 1681 treated observations and 849 untreated observations, for a total of 2530 observations. Bootstrap standard errors, based on 1000 bootstrap repetitions, are reported in parentheses. The repeated split sample estimator, $\hat{\tau}_k^{RSS}$, uses 100 repetitions. Each repetition randomly permutes the order of the untreated observations. Then, the first 425 untreated observations after re-ordering are used to estimate β . The remaining of 424 untreated observations and the 1681 treated observations are used in the second step estimation of τ_k . The “unadjusted” estimates are differences in mean outcomes between treated and non-treated. The “adjusted” estimates are regression coefficients on the treatment variable in a linear regression that includes the list of covariates detailed in Section 3.

* indicates statistical significance at the 0.10 level.

** indicates statistical significance at the 0.05 level.

Table 2
STAR Estimation Results

Panel A: Average treatment effect

	unadjusted			adjusted		
$\hat{\tau}$	0.1659** (0.0329)			0.1892** (0.0294)		

Panel B: Average treatment effect by predicted outcome group

	unadjusted			adjusted		
	low	medium	high	low	medium	high
$\hat{\tau}_k$	0.3705** (0.0521)	0.2688** (0.0655)	-0.1330** (0.0636)	0.3908** (0.0509)	0.3023** (0.0678)	-0.1242** (0.0614)
$\hat{\tau}_k^{LOO}$	0.3277** (0.0547)	0.2499** (0.0670)	-0.0486 (0.0654)	0.3440** (0.0519)	0.2730** (0.0696)	-0.0660 (0.0634)
$\hat{\tau}_k^{RSS}$	0.3152** (0.0467)	0.2617** (0.0505)	-0.0520 (0.0567)	0.3130** (0.0459)	0.3005** (0.0526)	-0.0374 (0.0552)

Note: The STAR sample includes 1747 treated observations and 2017 untreated observations, for a total of 3764 observations. Bootstrap standard errors, based on 1000 bootstrap repetitions, are reported in parentheses. The repeated split sample estimator, $\hat{\tau}_k^{RSS}$, uses 100 repetitions. Each repetition randomly permutes the order of the untreated observations. Then, the first 1009 untreated observations after re-ordering are used to estimate β . The remaining of 1008 untreated observations and the 1747 treated observations are used in the second step estimation of τ_k . The “unadjusted” estimates are differences in mean outcomes between treated and non-treated. The “adjusted” estimates are regression coefficients on the treatment variable in a linear regression that includes the list of covariates detailed in Section 3.

* indicates statistical significance at the 0.10 level.

** indicates statistical significance at the 0.05 level.

Table 3
JTPA Simulation Results

<i>Panel A: Bias</i>						
	unadjusted			adjusted		
	low	medium	high	low	medium	high
$\hat{\tau}_k$	1017.51	-4.81	-1082.42	1017.60	-0.98	-1062.39
$\hat{\tau}_k^{LOO}$	-88.23	-23.28	96.57	-59.08	-54.01	42.86
$\hat{\tau}_k^{RSS}$	-2.74	-2.34	-20.75	-3.30	-5.96	-17.56
$\hat{\tau}_k^{SSS}$	5.62	-9.88	6.62	1.34	-11.24	1.39
$\hat{\tau}_k^{UNF}$	-1.50	-8.56	-16.85	-2.50	-9.04	-11.67

<i>Panel B: Coverage rates for nominal 0.05 C.I.</i>						
	unadjusted			adjusted		
	low	medium	high	low	medium	high
$\hat{\tau}_k$	0.152	0.049	0.089	0.154	0.050	0.089
$\hat{\tau}_k^{LOO}$	0.051	0.048	0.050	0.051	0.049	0.051
$\hat{\tau}_k^{RSS}$	0.051	0.048	0.049	0.052	0.048	0.050
$\hat{\tau}_k^{SSS}$	0.050	0.048	0.051	0.052	0.049	0.050
$\hat{\tau}_k^{UNF}$	0.053	0.050	0.050	0.053	0.051	0.051

<i>Panel C: Root-MSE</i>						
	unadjusted			adjusted		
	low	medium	high	low	medium	high
$\hat{\tau}_k$	1492.26	1364.27	2145.78	1489.89	1375.04	2065.87
$\hat{\tau}_k^{LOO}$	1192.35	1399.74	1895.93	1180.13	1398.76	1800.86
$\hat{\tau}_k^{RSS}$	1031.61	1101.14	1751.43	1022.53	1103.53	1660.50
$\hat{\tau}_k^{SSS}$	1500.97	1797.74	2383.17	1493.40	1792.14	2271.90
$\hat{\tau}_k^{UNF}$	1119.34	1372.67	1867.51	1118.52	1383.76	1792.25

Note: Averages over 10000 simulations. See Section 4 and Appendix 2 for details.

Table 4
STAR Simulation Results

<i>Panel A: Bias</i>						
	unadjusted			adjusted		
	low	medium	high	low	medium	high
$\hat{\tau}_k$	0.0483	0.0006	-0.0511	0.0487	0.0010	-0.0506
$\hat{\tau}_k^{LOO}$	-0.0025	0.0005	0.0046	0.0028	-0.0025	-0.0075
$\hat{\tau}_k^{RSS}$	0.0001	-0.0000	-0.0012	0.0002	0.0001	-0.0010
$\hat{\tau}_k^{SSS}$	-0.0005	0.0004	-0.0017	-0.0002	0.0003	-0.0015
$\hat{\tau}_k^{UNF}$	0.0004	-0.0003	-0.0009	0.0002	-0.0002	-0.0006

<i>Panel B: Coverage rates for nominal 0.05 C.I.</i>						
	unadjusted			adjusted		
	low	medium	high	low	medium	high
$\hat{\tau}_k$	0.161	0.051	0.178	0.178	0.049	0.191
$\hat{\tau}_k^{LOO}$	0.048	0.050	0.051	0.050	0.049	0.056
$\hat{\tau}_k^{RSS}$	0.053	0.051	0.048	0.052	0.051	0.049
$\hat{\tau}_k^{SSS}$	0.051	0.050	0.052	0.052	0.052	0.050
$\hat{\tau}_k^{UNF}$	0.051	0.052	0.049	0.051	0.050	0.050

<i>Panel C: Root-MSE</i>						
	unadjusted			adjusted		
	low	medium	high	low	medium	high
$\hat{\tau}_k$	0.0695	0.0472	0.0716	0.0677	0.0471	0.0691
$\hat{\tau}_k^{LOO}$	0.0526	0.0509	0.0530	0.0492	0.0507	0.0494
$\hat{\tau}_k^{RSS}$	0.0473	0.0402	0.0470	0.0444	0.0399	0.0439
$\hat{\tau}_k^{SSS}$	0.0617	0.0589	0.0615	0.0577	0.0583	0.0571
$\hat{\tau}_k^{UNF}$	0.0501	0.0469	0.0503	0.0480	0.0475	0.0476

Note: Averages over 10000 simulations. See Section 4 and Appendix 2 for details.

Table 5
Bias in Simulations Using Artificial Data
(constant regression coefficients)

	K = 10						K = 20						K = 40					
	unadjusted			adjusted			unadjusted			adjusted			unadjusted			adjusted		
	low	med.	high	low	med.	high	low	med.	high	low	med.	high	low	med.	high	low	med.	high
N = 200																		
$\hat{\tau}_k$	2.24	-0.05	-2.28	2.23	-0.06	-2.27	3.15	-0.00	-3.13	3.12	-0.01	-3.08	4.09	0.02	-4.06	3.89	0.04	-3.88
$\hat{\tau}_k^{LOO}$	-0.32	-0.03	0.25	-0.07	-0.04	0.01	-0.18	0.00	0.17	0.30	0.01	-0.26	-0.09	0.00	0.13	0.89	0.02	-0.88
$\hat{\tau}_k^{RSS}$	-0.03	-0.04	-0.04	-0.02	-0.04	-0.05	0.01	-0.00	-0.01	0.03	0.01	0.01	0.03	0.01	-0.00	0.04	0.02	-0.01
$\hat{\tau}_k^{UNF}$	-0.04	-0.04	-0.03	-0.02	-0.03	-0.04	0.03	0.00	0.00	0.02	-0.01	0.03	0.02	0.00	0.01	-0.01	0.05	0.04
N = 1000																		
$\hat{\tau}_k$	0.55	0.01	-0.54	0.55	0.01	-0.54	0.71	-0.01	-0.71	0.70	-0.00	-0.70	0.82	0.01	-0.82	0.81	0.01	-0.82
$\hat{\tau}_k^{LOO}$	-0.05	0.01	0.07	-0.05	0.01	0.06	-0.05	-0.01	0.05	-0.05	-0.00	0.05	-0.01	0.01	0.01	-0.00	0.01	-0.01
$\hat{\tau}_k^{RSS}$	0.02	0.00	-0.00	0.02	0.01	-0.00	-0.01	-0.01	0.01	-0.01	-0.00	0.01	0.01	0.00	-0.01	0.01	0.00	-0.01
$\hat{\tau}_k^{UNF}$	0.01	0.01	0.00	0.00	0.01	0.00	-0.00	-0.01	-0.01	-0.00	-0.00	0.00	0.01	0.00	-0.01	0.01	0.01	-0.01
N = 5000																		
$\hat{\tau}_k$	0.12	-0.00	-0.12	0.12	-0.00	-0.12	0.15	0.00	-0.16	0.15	0.00	-0.15	0.16	-0.00	-0.16	0.16	-0.00	-0.16
$\hat{\tau}_k^{LOO}$	-0.01	-0.00	0.01	-0.01	-0.00	0.01	-0.01	0.00	0.00	-0.00	0.00	0.00	-0.01	-0.00	0.01	-0.01	-0.00	0.01
$\hat{\tau}_k^{RSS}$	0.01	-0.00	-0.01	0.01	-0.00	-0.01	0.00	-0.00	-0.01	0.01	-0.00	-0.01	-0.00	-0.00	0.00	0.00	-0.00	0.00
$\hat{\tau}_k^{UNF}$	0.01	-0.00	-0.01	0.01	-0.00	-0.01	0.01	-0.01	-0.01	0.01	-0.01	-0.00	-0.00	0.00	0.00	-0.00	0.00	0.01

Note: Averages over 10000 simulations. K is the number of included regressors. See Section 4 and Appendix 2 for details.

Table 6
 Bias in Simulations Using Artificial Data
 (decaying regression coefficients)

	K = 10						K = 20						K = 40					
	unadjusted			adjusted			unadjusted			adjusted			unadjusted			adjusted		
	low	med.	high	low	med.	high	low	med.	high	low	med.	high	low	med.	high	low	med.	high
N = 200																		
$\hat{\tau}_k$	2.94	-0.00	-2.97	2.97	-0.00	-2.99	4.57	-0.01	-4.60	4.63	-0.02	-4.69	6.86	0.04	-6.87	7.20	-0.01	-7.24
$\hat{\tau}_k^{LOO}$	-0.41	-0.01	0.40	0.01	-0.01	-0.02	-0.26	0.01	0.23	0.76	-0.04	-0.79	-0.14	-0.00	0.17	2.48	-0.00	-2.48
$\hat{\tau}_k^{RSS}$	-0.03	-0.01	0.01	-0.02	-0.01	0.01	-0.01	-0.01	-0.02	-0.02	-0.02	0.00	0.01	0.01	-0.02	0.01	0.02	0.02
$\hat{\tau}_k^{UNF}$	-0.01	-0.01	-0.01	-0.02	-0.01	0.00	-0.01	-0.03	0.02	0.00	-0.05	-0.01	-0.03	0.02	0.04	-0.03	0.02	0.01
N = 1000																		
$\hat{\tau}_k$	0.95	0.01	-0.94	0.95	0.01	-0.94	1.64	-0.01	-1.64	1.64	-0.00	-1.64	2.64	-0.01	-2.64	2.66	-0.01	-2.65
$\hat{\tau}_k^{LOO}$	-0.11	0.01	0.12	-0.10	0.01	0.11	-0.10	-0.01	0.11	-0.03	-0.01	0.04	-0.07	-0.00	0.08	0.14	0.00	-0.13
$\hat{\tau}_k^{RSS}$	0.01	0.00	0.01	0.01	0.00	0.01	-0.00	-0.00	0.00	-0.00	-0.00	0.01	0.00	0.00	-0.00	-0.00	0.01	0.01
$\hat{\tau}_k^{UNF}$	0.01	0.00	0.01	0.01	0.00	0.01	0.00	-0.00	-0.00	-0.00	-0.01	0.00	-0.01	-0.01	0.02	-0.01	-0.01	0.03
N = 5000																		
$\hat{\tau}_k$	0.23	-0.01	-0.22	0.23	-0.01	-0.22	0.43	-0.00	-0.43	0.43	-0.01	-0.43	0.81	-0.00	-0.81	0.81	-0.00	-0.81
$\hat{\tau}_k^{LOO}$	-0.02	-0.01	0.04	-0.02	-0.01	0.03	-0.03	-0.01	0.03	-0.03	-0.01	0.03	-0.03	-0.00	0.02	-0.02	-0.01	0.02
$\hat{\tau}_k^{RSS}$	0.00	-0.00	0.00	0.00	-0.00	0.00	-0.00	-0.00	0.00	-0.00	-0.00	0.00	-0.00	-0.00	0.00	-0.00	-0.00	-0.00
$\hat{\tau}_k^{UNF}$	0.00	-0.00	0.01	0.00	-0.00	0.01	0.00	-0.01	0.00	0.00	-0.01	0.00	-0.00	-0.00	0.00	-0.00	-0.00	0.00

Note: Averages over 10000 simulations. K is the number of included regressors. See Section 4 and Appendix 2 for details.